

ARTICLE

Open Access

# RhMYB108, an R2R3-MYB transcription factor, is involved in ethylene- and JA-induced petal senescence in rose plants

Shuai Zhang<sup>1,2</sup>, Qingcui Zhao<sup>1,2</sup>, Daxing Zeng<sup>1,2</sup>, Jiehua Xu<sup>1,2</sup>, Hougao Zhou<sup>3</sup>, Fenglan Wang<sup>3</sup>, Nan Ma<sup>4</sup> and Yonghong Li<sup>1,2</sup>

## Abstract

Rose (*Rosa hybrida*) plants are major ornamental species worldwide, and their commercial value greatly depends on their open flowers, as both the quality of fully open petals and long vase life are important. Petal senescence can be started and accelerated by various hormone signals, and ethylene is considered an accelerator of petal senescence in rose. To date, however, the underlying mechanism of signaling crosstalk between ethylene and other hormones such as JA in petal senescence remains largely unknown. Here, we isolated *RhMYB108*, an R2R3-MYB transcription factor, which is highly expressed in senescing petals as well as in petals treated with exogenous ethylene and JA. Applications of exogenous ethylene and JA markedly accelerated petal senescence, while the process was delayed in response to applications of 1-MCP, an ethylene action inhibitor. In addition, silencing of *RhMYB108* alter the expression of SAGs such as *RhNAC029*, *RhNAC053*, *RhNAC092*, *RhSAG12*, and *RhSAG113*, and finally block ethylene- and JA-induced petal senescence. Furthermore, *RhMYB108* was identified to target the promoters of *RhNAC053*, *RhNAC092*, and *RhSAG113*. Our results reveal a model in which *RhMYB108* functions as a receptor of ethylene and JA signals to modulate the onset of petal senescence by targeting and enhancing senescence-associated gene expression.

## Introduction

Petals are a key component of flowers, and by attracting pollinators, they play a vital role in ensuring successful pollination<sup>1</sup>. Petal senescence is a natural and irreversible process that occurs at the terminal stage of flower development and is tightly controlled by environmental and developmental signals. Many physiological activities occur together with petal senescence, such as the breaking down of intracellular structures, the degrading of membrane systems and macromolecules, and the recycling of nutrients, all of which similarly occur together with leaf

senescence<sup>2,3</sup>. Many excellent studies have reported that multiple vital activities are involved in petal senescence, such as programmed cell death (PCD)<sup>4–6</sup>, sugar metabolism<sup>7</sup>, hormone regulation<sup>8,9</sup>, and the regulatory activity of associated gene networks<sup>10–12</sup>.

The onset of petal senescence is initiated by external and internal cues, and phytohormones have been revealed to act as essential signals in initiating and modulating the senescence process. Among the hormones involved, ethylene and abscisic acid (ABA) function as inducers, and salicylic acid (SA), cytokinin (CTK), and gibberellic acid (GA) function as inhibitors<sup>13–16</sup>. The role of jasmonic acid (JA) and auxin in petal senescence is still unknown<sup>10</sup>.

Ethylene is considered a major endogenous signal of fruit ripening and leaf and petal senescence. Exogenous applications of ethylene can significantly accelerate petal senescence<sup>13</sup> and floral organ abscission<sup>17</sup>. *ETHYLENE INSENSITIVE 2* (*EIN2*), which functions as the core

Correspondence: Yonghong Li (liyongh@szpt.edu.cn)

<sup>1</sup>School of Applied Chemistry and Biological Technology, Postdoctoral Innovation Practice Base, Shenzhen Polytechnic, Shenzhen, Guangdong 518055, China

<sup>2</sup>Shenzhen Key Laboratory of Fermentation, Purification and Analysis, Shenzhen Polytechnic, Shenzhen 518055 Guangdong, China  
Full list of author information is available at the end of the article.

These authors contributed equally: Shuai Zhang, Qingcui Zhao

© The Author(s) 2019



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

member of ethylene signaling, acts as a key component in ethylene signaling. In *ein2-1* mutant plants, petal senescence and floral organ abscission are unaffected by exogenous ethylene applications. During petal senescence in rose, expression of the ethylene biosynthesis genes *1-AMINOCYCLOPROPANE-1-CARBOXYLIC ACID SYNTHASE 1* and *1-AMINOCYCLOPROPANE-1-CARBOXYLIC ACID SYNTHASE 2* (*ACS1* and *ACS2*, respectively) strongly increases, coupled with an accelerated increase in ethylene content<sup>18</sup>. It has been widely reported that JA functions in leaf senescence<sup>19–22</sup>, but its function in petal senescence is largely unclear. Some research has confirmed that applications of methyl jasmonate (MeJA), either as an aqueous solution or as a gas, accelerate petal senescence in *Petunia* and *Dendrobium*<sup>23</sup>; additional evidence has shown that, by completely suppressing tepal wilting, MeJA delays senescence in cut *Iris* flowers<sup>24</sup>.

Recent research on the ethylene and JA signaling pathways has shed light on the functional relationship of these two hormones in floral organ abscission<sup>17</sup> and leaf senescence<sup>25</sup>. Floral organ abscission is a typical characteristic that occurs in response to ethylene. In *ein2-1* mutants, floral organ abscission was shown to be unaffected by applications of exogenous ethylene, whereas applications of JA could rescue the ethylene response of those mutants<sup>17</sup>. These results suggest that interplay between JA and ethylene occurs. In Arabidopsis leaf senescence, the function of JA in accelerated senescence depends on the ethylene signaling pathway, especially the key molecular components *ETHYLENE INSENSITIVE 3* (*EIN3*) and *ETHYLENE INSENSITIVE 3-LIKE 1* (*EIL1*). Studies have shown that the *ein3* mutant displays delay JA-induced leaf senescence<sup>25</sup>. *EIN3/EIL1* are the direct targets of JAZ repressors, which inhibit *EIN3* and *EIL1* transcriptional activities via physical interactions. Moreover, JASMONATE-ZIM DOMAIN (JAZ) proteins recruit HISTONE DEACETYLASE 6 (HDA6), a positive regulator of JA signaling, which subsequently inhibits *EIN3/EIL1*-dependent transcription and JA signaling because of acetylation modification<sup>26</sup>. However, understanding of the signaling crosstalk between ethylene and JA during petal senescence is unclear. In *Dendrobium* and *Petunia*, the effects of JA-induced tepal/petal senescence are accompanied by a massive production of ethylene, and the effects on senescence can be abolished by treatment with antagonists of ethylene synthesis or ethylene action<sup>23</sup>. This evidence reveals that the JA signal appears to be dragged in the ethylene pathway to modulate senescence.

Typical signs of petal senescence include color change, wilting, and petal abscission<sup>8,10,18</sup>. In this study, we identified a MYB transcription factor, RhMYB108, as being involved in the interplay between ethylene and JA

signaling and thus regulating petal senescence in rose. According to their number of conserved MYB structural motifs, MYB proteins are classified into R1-MYB, R2R3-MYB, R3-MYB, and R4-MYB subfamilies<sup>27,28</sup> and function in the regulation of plant growth, hormone signal transduction, stress and disease resistance, and secondary metabolism<sup>29–33</sup>. Here, we show that RhMYB108, which belongs to the R2R3-MYB family, is intensely expressed in rose petals. Silencing of *RhMYB108* delayed petal senescence by directly regulating the expression of senescence-associated genes (SAGs), including *RhNAC053*, *RhNAC092*, and *RhSAG113*, by physically interacting with their promoter; thus, RhMYB108 functions in petal senescence. In brief, our findings highlight the importance of RhMYB108 in the onset of petal senescence and imply that its function may integrate ethylene and JA signaling crosstalk during petal senescence.

## Materials and methods

### Plant materials and treatments

*R. hybrida* cv. Samantha plantlets were propagated as previously described<sup>15,34</sup>. Rooted rose and tobacco (*Nicotiana benthamiana*) plants were transplanted into pots and then grown in a growth chamber whose conditions included a  $22 \pm 1$  °C temperature, a 16 h/8 h photoperiod and ~60% relative humidity.

With respect to JA treatment, flower buds of rose plantlets or petal discs were sprayed with 100  $\mu$ M MeJA, after which they were untouched for 24 h; the control samples were placed in 0.1% ethanol. With respect to ethylene or 1-MCP treatment, rose plantlets or petal discs were exposed to 10  $\mu$ L/L gaseous ethylene or 2  $\mu$ L/L 1-MCP, respectively, in an airtight growth container for 24 h, and NaOH solution (1 M) was also added to the container to prevent CO<sub>2</sub> accumulation, as previously described<sup>15,18,35</sup>.

### Measurement of ion leakage rates

Detached rose petals or petal discs were placed in 50 -mL tubes that contained 20 mL of deionized water. After the tubes were shaken at 28 °C for 30 min, the initial conductivity of the solution (C1) was measured using a conductivity meter (DDBJ-350, Precision & Scientific Instruments Co., Shanghai, China). The tubes were subsequently boiled in the same deionized water for 15 min, and the conductivity of the resulting solution (C2) was determined after the tubes cooled. The ion leakage rates were calculated by the C1:C2 ratios.

### RNA extraction and quantitative real-time PCR

RNA was extracted from the roots, stems, leaves, petals, sepals, and receptacles of rose plants as previously described<sup>36</sup>. The total RNA of the stamens and pistils was isolated using an RNA extraction kit (RN38-EASY, Aidlab

Biotech, Beijing, China). The RNA was then reverse-transcribed using QRT SuperMix for qPCR (R233-01, Vazyme Biotech Co., Nanjing, China) and oligo (dT) primers. In conjunction with 2  $\mu$ L of cDNA used as a template, qRT-PCR was performed via a Real-Time PCR System (Applied Biosystems, CA, USA) together with a Kapa SYBR Fast Universal qPCR Kit (Kapa Biosystems, Boston, MA, USA). *RhLUBI2* was used as an internal control.

#### Protein subcellular localization assays

With respect to subcellular localization assays, the open-reading frame (ORF) of *RhMYB108* without the stop codon was amplified by PCR and then fused into the N-terminal region of a GFP label; the fusion protein was driven by the Super promoter. On the 3rd day after infiltration of *Agrobacterium* carrying the corresponding plasmid, the tobacco leaves were monitored by laser confocal fluorescence microscopy (Olympus Fluoview FV1000). pSuper::GFP served as a negative control, and pSuper::NF-YA4-mCherry was used as a nuclear marker.

#### Transactivation and dual-luciferase reporter assays

Transcriptional activity assays of *RhMYB108* were performed in tobacco leaves. The ORF of *RhMYB108* was amplified and inserted into a pBD vector, which served as an effector. A double-reporter vector, which contained a GAL4-LUC and an internal control REN gene driven by the 35S promoter, was also used.

To analyze the interaction between *RhMYB108* and the SAG promoter in *N. benthamiana*, we constructed two vectors: pGreenII 0800-LUC and pGreenII 0029 62-SK<sup>37–39</sup>. The SAG promoters were amplified and inserted into the pGreenII 0800-LUC vector at the *Bam*HI and *Nco*I sites to drive the expression of the LUC reporter gene, and the *RhMYB108* ORF was inserted into the pGreenII 0029 62-SK vector at the *Bam*HI and *Kpn*I sites. The pGreenII 0800-LUC vector carries a *renilla luciferase* (REN) gene driven by the 35S promoter to serve as an internal control.

The effector construct and reporter plasmid were cotransformed into tobacco leaves by *Agrobacterium* infiltration as previously described<sup>40</sup>. At 48 h after transformation, the leaves were maintained in darkness for 5 min to measure LUC fluorescence.

#### Yeast one-hybrid assays

Y1H assays were performed as described in the Yeast Protocols Handbook (Clontech Laboratories, USA). Two vectors were constructed for use in this system: pLacZi and pJG4-5. SAG promoters were inserted into the pLacZi vector, and the *RhMYB108* ORF sequence was inserted into the pJG4-5 vector. The plasmid construct was subsequently transformed into yeast strain EGY48, which was

then grown on SD plates that lacked uracil and tryptophan, but contained X- $\beta$ -gal for blue/white colony screening.

#### Virus-induced gene silencing

VIGS of *RhMYB108* in rose petal discs and plantlets was performed as previously described<sup>15,41</sup>. A specific fragment of *RhMYB108* (437 bp in length) was amplified and inserted into a pTRV2 vector by *Xba*I and *Kpn*I. The vectors were transformed into *A. tumefaciens* strain GV3101, and the transformed *A. tumefaciens* cells were cultured in LB medium (supplemented with 50  $\mu$ g/mL kanamycin and 50  $\mu$ g/mL rifampicin) on a rocking platform (200 rpm) at 28 °C for 18 h. The cells were harvested and resuspended in infiltration buffer (10 mM MgCl<sub>2</sub>, 200 mM acetosyringone, 10 mM MES, pH 5.6) to a final OD<sub>600</sub> of ~1.5. The culture mixtures contained pTRV1- and pTRV2-*RhMYB108* at a ratio of 1:1 (v/v) or pTRV1 and pTRV2 (the negative control). The culture mixtures were incubated in the dark at room temperature for 3–4 h before infiltration.

With respect to VIGS in rose petal discs, the petals were collected at stage 2 of flower opening, and two 1 cm diameter discs were isolated from symmetrical locations of the petals using a hole punch. The plantlets were subsequently placed in deionized water for 1 day to equilibrate after multiplication by tissue culture as previously described<sup>15</sup>. The discs and plantlets were submerged in infiltration buffer and then exposed to a vacuum (–25 kPa) twice (each for 60 s). After the release of the vacuum, the discs and plantlets were briefly washed with deionized water and then incubated in the dark at 8 °C for 3 days. The phenotypes of the discs were observed daily until necrosis, and the flowers of the rose plantlets were monitored at various time points from stages 1 to 6.

#### Sequence analysis

With respect to multiple sequence alignment, the amino acid sequences were aligned using ClustalW with the default parameters (<https://www.genome.jp/tools-bin/clustalw>), and figures were created using the BioEdit software package (version 7.2.5)<sup>42</sup>. With respect to phylogenetic tree analysis, the alignment results were computed using MEGA software (version 5.05) based on the neighbor-joining algorithm and the following parameters: 1000 bootstrap replicates, Poisson correction, pairwise deletion, and uniform rates.

#### Statistical analyses

Statistical analysis of the data was performed via GraphPad Prism 8.0 (GraphPad Software Inc., USA; <http://www.graphpad.com/>). Two groups of data were compared using two-sided Student's *t* tests (\**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001; \*\*\*\**P* < 0.0001). The means of multiple groups


of data were compared via one-way ANOVA and Bonferroni's post hoc test, with  $P < 0.05$  considered significant.

## Results

### Ethylene and JA accelerate the senescence of rose petals

The function of ethylene and JA in the acceleration of leaf senescence has been widely reported<sup>21,25</sup>, but the interaction and crosstalk between these hormones during flower senescence has not been studied in detail. To determine the effects of ethylene and JA during flower senescence, we introduced a rose petal disc system<sup>13,43</sup>. As shown in Fig. 1a, the control petal discs at 10 days after incubation displayed slight color fading. After ethylene or JA treatment, the color fading started at 5 days, and nearly

all discs showed severe color loss after 10 days of incubation. The color fading rate and ion permeability were markedly higher in the treated petals than in the control petals (Fig. 1b, c). Furthermore, the transcript levels of SAG *RhSAG12*, a molecular marker of rose petal senescence progression, were also significantly higher in the hormone-treated petal discs than in the control discs (Fig. 1d). In addition, compared with those treated with ethylene or JA, the petal discs treated with 1-MCP, a powerful antagonist of ethylene action, presented a slight color loss (Fig. 1a), and this result was supported by the experimental data concerning the color fading rate (Fig. 1b), ion leakage rate (Fig. 1c), and *RhSAG12* expression (Fig. 1d). Briefly, these results indicate that ethylene and



JA accelerate the senescence process of rose petals and that the role of JA in petal senescence at least partly depends on ethylene signaling.


### Expression of *RhMYB108* increased with petal senescence and was induced by ethylene and JA

Analysis of the information within the expressed sequence tag (EST) database of rose petals treated with exogenous ethylene ([http://bioinfo.bti.cornell.edu/cgi-bin/rose\\_454/index.cgi](http://bioinfo.bti.cornell.edu/cgi-bin/rose_454/index.cgi))<sup>44</sup> revealed a transcript, RU04450, that encodes a putative MYB family transcription factor whose expression was markedly induced in response to the application of exogenous ethylene. The cDNA sequence of RU04450 is 1129 bp in length, with a 1032 bp ORF that encodes 343 amino acid residues. Phylogenetic analysis and sequence alignment revealed that RU04450 is closely related to *AtMYB108* from *Arabidopsis thaliana* (AT3G06490) (Supplementary Fig. 1a)

and belongs to the R2R3-MYB family (Supplementary Fig. 1b); hence, the gene corresponding to RU04450 was named *RhMYB108*.

To further investigate the function of *RhMYB108* during flower senescence, the transcripts of *RhMYB108* in all rose organs were measured by qRT-PCR. As shown in Fig. 2a, the expression of *RhMYB108* was greater in the rose petals than in the roots and stems, and was lowest in the leaves.


We further evaluated the expression of *RhMYB108* transcripts in different floral organs, including the petals, sepals, receptacles, stamens, and pistils, during rose flower opening. The expression of *RhMYB108* increased incrementally in the petals in parallel with flower opening (Fig. 2b), further implying that this gene may function in petal senescence. The *RhMYB108* expression pattern trends were similar between the sepals and stamens (Fig. 2d, e), and the transcript levels were higher in the




partially opened flower buds (stages 3–4) than at other stages. In the receptacle and pistil, *RhMYB108* expression increased throughout the flower-opening process (Fig. 2c, f). In addition, we confirmed that petal senescence could be accelerated by exogenous applications of ethylene and JA. Expression of the *RhMYB108* transcript was consistently induced by treatment with exogenous ethylene and JA, and compared with that in response to applications of the control solution, its expression in response to applications of the ethylene analog 1-MCP was not altered (Fig. 2g). These data provide evidence of a correlation between *RhMYB108* expression and petal senescence.

### Silencing of *RhMYB108* delays petal senescence

To assess the biological function of *RhMYB108* during the senescence of rose petals, we used VIGS specifically to silence *RhMYB108* transcripts in petal discs and rose plantlets; for this, we constructed a tobacco rattle virus vector (TRV-*RhMYB108*) from the 3' end fragment of *RhMYB108* (431 bp in length). The rose petal disc assay revealed that color fading started at 8 days in the TRV control, and the discs were nearly discolored on day 16 (Fig. 3a). By contrast, the discs whose *RhMYB108* gene was silenced exhibited a markedly delayed senescent phenotype and displayed a slight color fading at 16 days.



**Fig. 3** Silencing of *RhMYB108* delays senescence in rose petal discs. **a** Color fading phenotypes, **b** *RhMYB108* expression levels, **c** the color fading rate, **d** ion leakage rates, **e** *RhSAG12* expression levels were analyzed in *RhMYB108*-silenced and TRV control petal discs. The error bars represent the means of three biological replicates  $\pm$  SDs, and the asterisks indicate significant differences according to Student's *t* test (\* $P < 0.05$ ; \*\* $P < 0.01$ ; \*\*\* $P < 0.001$ ; \*\*\*\* $P < 0.0001$ ).



The expression of *RhMYB108* was significantly lower in the *RhMYB108*-silenced discs than in the TRV control discs (Fig. 3b). Consistent with this pattern, the silenced discs also presented significantly less color fading, ion leakage, and lower expression of *RhSAG12* than did the TRV control discs (Fig. 3c, d, e). With respect to the petal senescence process in rose plantlets, the *RhMYB108*-silenced plants also displayed a distinctly slower senescent phenotype than did the control plants (Fig. 4a); expression of *RhMYB108* and *RhSAG12* was markedly lower in the *RhMYB108*-silenced plants than in the TRV control plants (Fig. 4c, d). The flower life cycle of the *RhMYB108*-silenced plants was  $11.1 \pm 1.0$  days, whereas that of the TRV control plants was  $7.9 \pm 0.9$  days (Fig. 4b). These results indicate that *RhMYB108* functions as a positive regulator of the petal senescence process.

#### Ethylene- and JA-triggered petal senescence is *RhMYB108* dependent


We previously found that ethylene and JA could accelerate the senescence process of rose petals and that a reduction in *RhMYB108* expression led to a delayed-petal-senescence phenotype of *RhMYB108*-silenced plants. We therefore postulated that *RhMYB108* could

also function in ethylene- and JA-induced petal senescence.

To address this hypothesis further, we observed the phenotypes of *RhMYB108*-silenced petal discs or rose plantlets in response to applications of exogenous ethylene or JA. Compared with the TRV control discs, the *RhMYB108*-silenced discs presented a slightly senescent phenotype, which included mild color loss, markedly less ion leakage, and reduced *SAG12* expression, after the discs were in culture dishes for 8 days after ethylene or JA treatment. By day 12 after hormone treatment, a more extreme difference in the senescent phenotype emerged between the silenced discs and the TRV control discs (Fig. 5). With respect to the petal senescence process in rose plantlets, the phenotype of the *RhMYB108*-silenced plants revealed delayed ethylene- and JA-induced petal senescence (Fig. 6), which is consistent with the role of *RhMYB108* in the petal disc assay. These results indicate that *RhMYB108* accelerates the petal senescence process mediated by both ethylene and JA signaling.


#### *RhMYB108* directly targets senescence-related genes

To determine the subcellular localization of *RhMYB108*, we expressed an *RhMYB108*-GFP protein



**Fig. 5** Effects of exogenous ethylene and JA treatment on the senescence of *RhMYB108*-silenced rose petal discs. **a** The phenotypes of rose petal discs treated with 10  $\mu$ L/L ethylene or 100  $\mu$ M MeJA were recorded. **b, c, d, e** Color fading and ion leakage rates (**b, c** ethylene treatment; **d, e** JA treatment) of *RhMYB108*-silenced and TRV control petal discs. **d** transcript levels of *RhSAG12* in *RhMYB108*-silenced and TRV control petal discs. Three biological replicates were tested, and the error bars represent the SDs. The asterisks indicate statistically significant differences (Student's *t* test, \**P* < 0.05; \*\**P* < 0.01; \*\*\**P* < 0.001; \*\*\*\**P* < 0.0001).





together with NF-YA4-mCherry (an mCherry-labeled nuclear marker protein) in *N. benthamiana* leaves, and both proteins were driven by the pSuper promoter. We observed a GFP fluorescence signal pattern that was consistent with the localization of the nuclear marker protein (mCherry) in the nucleus, indicating that RhMYB108 is a nuclear protein (Supplementary Fig. 2a).

To confirm whether RhMYB108 acts as a transcriptional regulator, we performed a transactivation assay in *N. benthamiana* leaves using the dual-luciferase reporter system. Compared with the pBD-negative control, the full-length (MYB108), middle-length (MYB108<sup>1-284</sup>), and short-length (MYB108<sup>1-204</sup>) RhMYB108 variants intensely activated the reporter gene (Supplementary Fig. 2b), and the relative LUC activity was 69.7-fold, 32.5-fold, and 10.7-fold higher than that of the negative control, respectively (Supplementary Fig. 2c). These results indicate that RhMYB108 functions as a transcriptional activator and that the C-terminus has high transactivation activity.

SAGs usually serve as markers of the senescence process. Here, a series of SAG expression levels were detected


using qRT-PCR in *RhMYB108*-silenced plants and TRV control plants. As expected, the expression of only *NAC083* was obviously upregulated, acting as a senescence-inhibiting regulator<sup>45</sup> when *RhMYB108* was silenced. The expression trend of *WRKY22* was markedly similar in both the *RhMYB108*-silenced plants and the TRV control plants, while the expression of the transcripts of *NAC029*, *NAC053*, *NAC092*, *WRKY40*, *WRKY53*, *SAG12*, and *SAG113* was notably inhibited in the *RhMYB108*-silenced plants (Fig. 7a). These data confirmed that by regulating the expression of SAGs, *RhMYB108* may act as a transcriptional activator to accelerate petal senescence.

To better understand the possible regulatory mechanism that governs how *RhMYB108* regulates petal senescence in rose plants, the promoter region of SAGs, including the 1.5-kb region upstream of the initiation codon, was amplified and inserted into a pLacZi vector, and the ORF of *RhMYB108* was amplified and inserted into a pJG4-5 plasmid, yielding an expression vector. Y1H assays revealed that the RhMYB108 protein could bind to the promoter region of *NAC053*, *NAC092*, and *SAG113*, triggering the expression of the LacZ reporter gene (Fig. 7b). Additional evidence from a dual-luciferase reporter assay in *N. benthamiana* revealed that the activity of the promoters of *NAC053*, *NAC092*, and *SAG113* could be induced by overexpression of *RhMYB108* driven by the CaMV 35S promoter (Fig. 7c, d). Together, these results clarify that by directly altering SAG expression, RhMYB108 functions in the acceleration of petal senescence.

## Discussion

### RhMYB108 positively regulates petal senescence in rose

Petal senescence is a highly programmed event that follows visible signs, including color fading, wilting, and abscission<sup>13</sup>. At the transcriptional level, the expression of many transcription factors is altered during senescence, implying that complex regulatory networks are involved in this process<sup>46,47</sup>. As members of one of the most important families of transcription factors in plants, MYB proteins play pivotal roles in modulating multiple developmental and physiological processes. In Arabidopsis, only a few MYB transcription factors, such as MYBL<sup>48</sup>, MYBH<sup>49</sup>, and MYB2<sup>50</sup>, are reportedly involved in the modulation of the senescence process. We found that *RhMYB108* transcripts were more highly expressed in the petals than in other organs, including the roots, stems, and leaves (Fig. 2a). In addition, *RhMYB108* expression was higher in senescent petals than in young petals during flower opening (Fig. 2b). These data implied that *RhMYB108* may play a role in petal senescence. Furthermore, compared with the TRV control plants, the *RhMYB108*-silenced plants exhibited delayed petal



senescence (Figs. 3, 4), along with less ion leakage and lower *SAG12* expression. The expression of several SAGs, such as *SAG12*, *SAG113*, *NAC029*, *NAC053*, and *NAC092*, which serve as molecular markers of the senescence process, was downregulated in the *RhMYB108*-silenced lines (Fig. 7a). Additional research showed that these SAGs contain some MYB-specific cis-elements in their promoters, and our Y1H and dual-luciferase reporter assays demonstrated that these SAGs (*NAC053*, *NAC092*, and *SAG113*) were direct targets of *RhMYB108* (Fig. 7b, c, d).

Several previous studies have demonstrated that, by acting downstream of *AtMYB21*, *AtMYB108* can function at the transcriptional level in *Arabidopsis* stamen and pollen maturation in response to JA<sup>51</sup>. *AtMYB108* (*AtBOS1*) also acts as a negative regulator in controlling wound-induced cell death mediated by ABA signaling<sup>52</sup>. Recently, by targeting the *AtNAC003* promoter during dark-stressed leaf senescence, *AtMYB108* was shown to directly control *AtNAC003* expression<sup>53</sup>. These reports reveal that the MYB108 transcription factor participates in complex signaling and regulatory networks mediated by multiple phytohormone pathways and stress responses.

#### **RhMYB108 mediates ethylene and JA signaling during senescence**

Multiple phytohormones play a role in regulating the process of petal senescence. Among these hormones,

ethylene is generally considered a senescence accelerator on the basis of increased ethylene production during petal senescence. The function of JA in leaf senescence has been widely investigated<sup>19,21,22,54</sup>; however, its role in petal senescence is currently unclear, and the detailed mechanism by which signaling crosstalk between ethylene and JA modulates petal senescence is still poorly understood.

Studies have shown that endogenous JA levels are stable during flower opening or senescence in orchids and in *Lilium*<sup>8,55,56</sup>. Another study showed that applications of endogenous JA could delay tepal senescence in *Iris* flowers by 2 days<sup>57</sup>. In contrast, treatment of *Dendrobium* and *Petunia* flowers with JA reportedly accelerated petal senescence by promoting ethylene production. In addition, applications of silver thiosulfate, an inhibitor of ethylene action, completely inhibited the effects of JA function on accelerating petal senescence<sup>23</sup>. In conclusion, the role of JA in petal senescence seems contradictory, and additional investigations are needed to reveal the definite mechanisms underlying JA-mediated petal senescence. Here, our results are consistent with those of the study by Porat et al.<sup>23</sup> and further verify the positive effects of JA on petal senescence. In addition, the function of JA in accelerating the senescence process was abolished by applying 1-MCP, an ethylene action inhibitor (Fig. 1). Our data imply that JA may act as a promoter of petal

senescence via the ethylene signaling pathway. The signaling crosstalk between ethylene and JA may depend on the interaction of JAZ-EIN3 and JAZ-EIL1. A previous paper revealed that JA could enhance the transcriptional activity of EIN3/EIL1 by removing JAZ proteins, which physically interact with and inhibit *EIN3/EIL1* transcription and JA signaling by recruiting the corepressor HDA6, an RPD3-type histone deacetylase that functions in modulating histone acetylation<sup>26</sup>.

qRT-PCR analysis revealed that the expression of *RhMYB108* transcripts increased in response to exogenous ethylene and JA treatment (Fig. 2g), and the results implied that *RhMYB108* could be involved in the ethylene and JA signaling pathways at the transcriptional level. Overall, *RhMYB108* may function as a member of the regulatory network that integrates ethylene and JA signaling to modulate the onset and progression of petal senescence. In the future, it will be interesting to investigate how ethylene and JA signaling regulates *RhMYB108* transcription, which would provide further insights into the detailed mechanism that governs signaling crosstalk between ethylene and JA during petal senescence.

In this study, we identified a MYB transcription factor, *RhMYB108*, as being highly expressed in senescing petals as well as in petals treated with exogenous ethylene and JA. *RhMYB108* belongs to the R2R3-MYB family and is closely related to *AtMYB108* from *Arabidopsis thaliana* (AT3G06490). Silencing of *RhMYB108* results in delayed petal senescence and blocks ethylene- and JA-induced petal senescence by altering the expression of SAGs, including *RhNAC029*, *RhNAC053*, *RhNAC092*, *RhSAG12*, and *RhSAG113*. Furthermore, *RhMYB108* was identified as targeting the promoters of *RhNAC053*, *RhNAC092*, and *SAG113*. In brief, our findings highlight the importance of *RhMYB108* in the onset of petal senescence and imply that its function may integrate ethylene- and JA signaling crosstalk during petal senescence.

#### Acknowledgements

We thank AJE (<https://www.aje.cn/>) for the careful editing of this article. This work was supported by the National Natural Science Foundation of China (Grant Nos. 31572162 and 31902054) and the School Project of Shenzhen Polytechnic (No. 601822K27003).

#### Author details

<sup>1</sup>School of Applied Chemistry and Biological Technology, Postdoctoral Innovation Practice Base, Shenzhen Polytechnic, Shenzhen, Guangdong 518055, China. <sup>2</sup>Shenzhen Key Laboratory of Fermentation, Purification and Analysis, Shenzhen Polytechnic, Shenzhen 518055 Guangdong, China. <sup>3</sup>College of Horticulture and Landscape Architecture, Zhongkai University of Agriculture and Engineering, Guangzhou, Guangdong 510642, China. <sup>4</sup>China Beijing Key Laboratory of Development and Quality Control of Ornamental Crops, Department of Ornamental Horticulture, China Agricultural University, Beijing, China

#### Authors' contributions

S.Z. and Q.Z. performed the experiments. N.M. and Y.L. designed the research. D.Z., J.X., H.Z. and F.W. provided technical support and conceptual advice and analyzed the data. S.Z. and N.M. wrote the paper.

#### Data availability

The GenBank accession number of the *RhMYB108* rose gene used in this study is MK606453.

#### Conflict of interest

The authors declare that they have no conflict of interest.

**Supplementary Information** accompanies this paper at (<https://doi.org/10.1038/s41438-019-0221-8>).

Received: 1 September 2019 Revised: 18 October 2019 Accepted: 3 November 2019

Published online: 01 December 2019

#### References

- Ma, N. et al. Petal senescence: a hormone view. *J. Exp. Bot.* **69**, 719–732 (2018).
- Xu, Y. & Hanson, M. R. Programmed cell death during pollination-induced petal senescence in petunia. *Plant Physiol.* **122**, 1323–1334 (2000).
- Price, A. M. et al. A comparison of leaf and petal senescence in wallflower reveals common and distinct patterns of gene expression and physiology. *Plant Physiol.* **147**, 1898–1912 (2008).
- Rogers, H. J. Programmed cell death in floral organs: how and why do flowers die? *Ann. Bot.* **97**, 309–315 (2006).
- Van Doorn, W. G. & Woltering, E. J. Senescence and programmed cell death: substance or semantics? *J. Exp. Bot.* **55**, 2147–2153 (2004).
- Van Doorn, W. et al. Morphological classification of plant cell deaths. *Cell Death Differ.* **18**, 1241 (2011).
- Van Doorn, W. G. Is petal senescence due to sugar starvation? *Plant Physiol.* **134**, 35–42 (2004).
- Tripathi, S. K. & Tuteja, N. Integrated signaling in flower senescence: an overview. *Plant Signal. Behav.* **2**, 437–445 (2007).
- Kumar, N., Srivastava, G. C. & Dixit, K. Flower bud opening and senescence in roses (*Rosa hybrida* L.). *Plant Growth Regul.* **55**, 81–99 (2008).
- Van Doorn, W. G. & Woltering, E. J. Physiology and molecular biology of petal senescence. *J. Exp. Bot.* **59**, 453–480 (2008).
- Rogers, H. J. From models to ornamentals: how is flower senescence regulated? *Plant Mol. Biol.* **82**, 563–574 (2013).
- Shahri, W. & Tahir, I. Flower senescence: some molecular aspects. *Planta* **239**, 277–297 (2014).
- Lü, P. et al. *RhHB1* mediates the antagonism of gibberellins to ABA and ethylene during rose (*Rosa hybrida*) petal senescence. *Plant J.* **78**, 578–590 (2014).
- Trivellini, A., Cocetta, G., Vernieri, P., Mensuali-Sodi, A. & Ferrante, A. J. Effect of cytokinins on delaying petunia flower senescence: a transcriptome study approach. *Plant Mol. Biol.* **87**, 169–180 (2015).
- Wu, L. et al. An ethylene-induced regulatory module delays flower senescence by regulating cytokinin content. *Plant Physiol.* **173**, 853–862 (2017).
- Shabaniyan, S., Esfahani, M. N., Karamian, R. & Tran, L. S. P. Salicylic acid modulates cutting-induced physiological and biochemical responses to delay senescence in two gerbera cultivars. *Plant Growth Regul.* **87**, 245–246 (2018).
- Kim, J., Patterson, S. E. & Binder, B. M. Reducing jasmonic acid levels causes *ein2* mutants to become ethylene responsive. *FEBS Lett.* **587**, 226–230 (2013).
- Ma, N., Cai, L., Lu, W. J., Tan, H. & Gao, J. P. Exogenous ethylene influences flower opening of cut roses (*Rosa hybrida*) by regulating the genes encoding ethylene biosynthesis enzymes. *Sci. China Ser. C.* **48**, 434 (2005).
- He, Y., Fukushige, H., Hildebrand, D. F. & Gan, S. Evidence supporting a role of jasmonic acid in *Arabidopsis* leaf senescence. *Plant Physiol.* **128**, 876–884 (2002).
- Shibuya, K., Shimizu, K., Niki, T. & Ichimura, K. Identification of a NAC transcription factor, EPHEMERAL1, that controls petal senescence in Japanese morning glory. *Plant J.* **79**, 1044–1051 (2014).
- Hu, Y. et al. Jasmonate regulates leaf senescence and tolerance to cold stress: crosstalk with other phytohormones. *J. Exp. Bot.* **68**, 1361–1369 (2017).
- Zhang, Y. et al. Circadian evening complex represses jasmonate-induced leaf senescence in *Arabidopsis*. *Mol. Plant* **11**, 326–337 (2018).
- Porat, R., Borochov, A. & Halevy, A. H. Enhancement of petunia and *Andromeda* flower senescence by jasmonic acid methyl ester is via the promotion of ethylene production. *Plant Growth Regul.* **13**, 297–301 (1993).

24. Van Doorn, W. G., Çelikel, F. G., Pak, C. & Harkema, H. Delay of Iris flower senescence by cytokinins and jasmonates. *Physiol. Plant.* **148**, 105–120 (2013).
25. Li, Z., Peng, J., Wen, X. & Guo, H. ETHYLENE-INSENSITIVE3 is a senescence-associated gene that accelerates age-dependent leaf senescence by directly repressing miR164 transcription in Arabidopsis. *Plant Cell* **25**, 3311–3328 (2013).
26. Zhu, Z. et al. Derepression of ethylene-stabilized transcription factors (EIN3/EL1) mediates jasmonate and ethylene signaling synergy in Arabidopsis. *Proc. Natl Acad. Sci. USA* **108**, 12539–12544 (2011).
27. Dubos, C. et al. MYB transcription factors in Arabidopsis. *Trends Plant Sci.* **15**, 573–581 (2010).
28. Li, H. L., Guo, D. & Peng, S. Q. Molecular characterization of the *Jatropha curcas* JcR1MYB1 gene encoding a putative R1-MYB transcription factor. *Genet. Mol. Biol.* **37**, 549–555 (2014).
29. Kranz, H. D. et al. Towards functional characterisation of the members of the R2R3-MYB gene family from Arabidopsis thaliana. *Plant J.* **16**, 263–276 (1998).
30. Stracke, R., Werber, M. & Weisshaar, B. J. The R2R3-MYB gene family in Arabidopsis thaliana. *Curr. Opin. Plant Biol.* **4**, 447–456 (2001).
31. Jiang, C., Gu, J., Chopra, S., Gu, X. & Peterson, T. J. Ordered origin of the typical two- and three-repeat Myb genes. *Gene* **326**, 13–22 (2004).
32. Cheng, H. Q. et al. The cotton MYB108 forms a positive feedback regulation loop with CML11 and participates in the defense response against *Verticillium dahliae* infection. *J. Exp. Bot.* **67**, 1935–1950 (2016).
33. Du, Y. T. et al. Identification and characterization of GmMYB118 responses to drought and salt stress. *BMC Plant Biol.* **18**, 320 (2018).
34. Zhang, S. et al. In rose, transcription factor PTM balances growth and drought survival via PIP2;1 aquaporin. *Nat. Plants* **5**, 290–299 (2019).
35. Xue, J., Yang, F. & Gao, J. Isolation of Rh-TIP1;1, an aquaporin gene and its expression in rose flowers in response to ethylene and water deficit. *Postharvest. Biol. Tec.* **51**, 407–413 (2009).
36. Wan, C. Y. & Wilkins, T. A. A modified hot borate method significantly enhances the yield of high-quality RNA from cotton (*Gossypium hirsutum* L.). *Anal. Biochem.* **223**, 7–12 (1994).
37. Hellens, R. P. et al. Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants. *Plant Methods* **1**, 13 (2005).
38. Yin, X., Allan, A. C., Chen, K., Ferguson, I. B. & Kiwifruit, E. I. L. and ERF genes involved in regulating fruit ripening. *Plant Physiol.* **153**, 1280–1292 (2010).
39. Wei, Q. et al. Control of chrysanthemum flowering through integration with an aging pathway. *Nat. Commun.* **8**, 829 (2017).
40. Han, Y. et al. Banana transcription factor MaERF11 recruits histone deacetylase MaHDA1 and represses the expression of MaACO1 and expansins during fruit ripening. *Plant Physiol.* **171**, 1070–1084 (2016).
41. Tian, J. et al. TRV-GFP: a modified Tobacco rattle virus vector for efficient and visualizable analysis of gene function. *J. Exp. Bot.* **65**, 311–322 (2013).
42. Hall, T. A. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41**, 95–98 (1999).
43. Liu, J. et al. A RhABF2/Ferritin module affects rose (*Rosa hybrida*) petal dehydration tolerance and senescence by modulating iron levels. *Plant J.* **92**, 1157–1169 (2017).
44. Pei, H. et al. Integrative analysis of miRNA and mRNA profiles in response to ethylene in rose petals during flower opening. *PLoS ONE* **8**, e64290 (2013).
45. Yang, S. D., Seo, P. J., Yoon, H. K. & Park, C. M. The Arabidopsis N. A. C. transcription factor VNI2 integrates abscisic acid signals into leaf senescence via the COR/RD genes. *Plant Cell* **23**, 2155–2168 (2011).
46. Liu, X. et al. LSD: a leaf senescence database. *Nucleic Acids Res.* **39**, D1103–D1107 (2010).
47. Breeze, E. et al. High-resolution temporal profiling of transcripts during Arabidopsis leaf senescence reveals a distinct chronology of processes and regulation. *Plant Cell* **23**, 873–894 (2011).
48. Zhang, X. et al. The RR-type MYB-like transcription factor, AtMYBL, is involved in promoting leaf senescence and modulates an abiotic stress response in Arabidopsis. *Plant Cell Physiol.* **52**, 138–148 (2010).
49. Huang, C. K. et al. A single-repeat MYB transcription repressor, MYBH, participates in regulation of leaf senescence in Arabidopsis. *Plant Mol. Biol.* **88**, 269–286 (2015).
50. Guo, Y. & Gan, S. AtMYB2 regulates whole plant senescence by inhibiting cytokinin-mediated branching at late stages of development in Arabidopsis. *Plant Physiol.* **156**, 1612–1619 (2011).
51. Mandaokar, A. & Browse, J. MYB108 acts together with MYB24 to regulate jasmonate-mediated stamen maturation in Arabidopsis. *Plant Physiol.* **149**, 851–862 (2008).
52. Cui, F., Brosché, M., Sipari, N., Tang, S. & Overmyer, K. Regulation of ABA dependent wound induced spreading cell death by MYB108. *N. Phytol.* **200**, 634–640 (2013).
53. Chou, M. L. et al. The direct involvement of dark-induced Tic55 protein in chlorophyll catabolism and its indirect role in the MYB108-NAC signaling pathway during leaf senescence in Arabidopsis thaliana. *Int. J. Mol. Sci.* **19**, 1854 (2018).
54. Jiang, Y., Liang, G., Yang, S. & Yu, D. Arabidopsis WRKY57 functions as a node of convergence for jasmonic acid- and auxin-mediated signaling in jasmonic acid-induced leaf senescence. *Plant Cell* **26**, 230–245 (2014).
55. Porat, R., Halevy, A., Serek, M. & Borochoy, A. J. An increase in ethylene sensitivity following pollination is the initial event triggering an increase in ethylene production and enhanced senescence of Phalaenopsis orchid flowers. *Physiol. Plant* **93**, 778–784 (1995).
56. Arron, L. & Munné-Bosch, S. Hormonal changes during flower development in floral tissues of Liliun. *Planta* **236**, 343–354 (2012).
57. Imsabai, W. & Van Doorn, W. G. Effects of auxin, gibberellin, and cytokinin on petal blackening and flower opening in cut lotus flowers (*Nelumbo nucifera*). *Postharvest. Biol. Tech.* **75**, 54–57 (2013).