ARTICLE

Open Access

A temporal gene expression map of *Chrysanthemum* leaves infected with *Alternaria alternata* reveals different stages of defense mechanisms

Ye Liu¹, Jingjing Xin¹, Lina Liu¹, Aiping Song¹, Zhiyong Guan¹, Weimin Fang¹ and Fadi Chen¹

Abstract

Chrysanthemum (*Chrysanthemum morifolium*) black spot disease (CBS) poses a major threat to *Chrysanthemum* cultivation owing to suitable climate conditions and current lack of resistant cultivars for greenhouse cultivation. In this study, we identified a number of genes that respond to *Alternaria alternata* infection in resistant and susceptible *Chrysanthemum* cultivars. Based on RNA sequencing technology and a weighted gene coexpression network analysis (WGCNA), we constructed a model to elucidate the response of *Chrysanthemum* leaves to *A. alternata* infection at different stages and compared the mapped response of the resistant cultivar 'Jinba' to that of the susceptible cultivar 'Zaoyihong'. In the early stage of infection, when lesions had not yet formed, abscisic acid (ABA), salicylic acid (SA) and EDS1-mediated resistance played important roles in the *Chrysanthemum* defense system. With the formation of necrotic lesions, ethylene (ET) metabolism and the Ca^{2+} signal transduction pathway strongly responded to *A. alternata* infection. During the late stage, when necrotic lesions continued to expand, members of the multidrug and toxic compound extrusion (MATE) gene family were highly expressed, and their products may be involved in defense against *A. alternata* invasion by exporting toxins produced by the pathogen, which plays important roles in the pathogenicity of *A. alternata*. Furthermore, the function of hub genes was verified by qPCR and transgenic assays. The identification of hub genes at different stages, the comparison of hub genes between the two cultivars and the highly expressed genes in the resistant cultivar 'Jinba' provide a theoretical basis for breeding cultivars resistant to CBS.

Introduction

The production of cut *Chrysanthemum* flowers growing under greenhouse conditions is threatened by many diseases. *Alternaria alternata*, a common necrotrophic fungus that is ubiquitously present on various plant species, damages plant tissues by producing toxins^{1,2}. Warm and humid environmental conditions lead to the spread of diseases caused by this fungus, which is a year-round problem for *Chrysanthemum* production in greenhouses.

Although the early stage of the infection is symptomless, small round black spots form at the site of *A. alternata* invasion, and these spots eventually expand into round, round-like, or irregular spots that are covered with a dark mildew layer. The widespread occurrence of this disease severely affects the yield and ornamental value of *Chrysanthemum*, causing substantial losses in flower production.

Currently, *Chrysanthemum* black spot disease (CBS) is controlled by fungicides; however, residues and severe environmental pollution caused by fungicides contradict the green, sustainable development of agricultural production. Therefore, screening and breeding resistant *Chrysanthemum* cultivars is the most environmentally

Correspondence: Fadi Chen (chenfd@njau.edu.cn)

¹State Key Laboratory of Crop Genetics and Germplasm Enhancement, Key Laboratory of Landscaping, Ministry of Agriculture and Rural Affairs, College of Horticulture, Nanjing Agricultural University, Nanjing, China
These authors contributed equally: Ye Liu, Jingjing Xin

© The Author(s) 2020

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

friendly approach to controlling CBS. Furthermore, revealing the molecular mechanism of CBS resistance will provide a strong theoretical basis for formulating disease resistance breeding strategies and preventing black spot disease. In our laboratory, a variety of Chrysanthemum germplasm resources with different levels of resistance to black spot disease are being used to identify key genes involved in disease resistance.

The response of the plant immune system to pathogen infection involves two branches: PTI (pathogen-associated molecular pattern-triggered immunity) and ETI (effector-triggered immunity)³. PTI is mainly triggered by the perception of microbial proteins, leading to the deposition of callose in the cell wall, MAP kinase cascade signal transduction, changes in gene expression, closure of stomata, and/or an oxidative burst^{4–6}. ETI is regulated by resistance genes (R genes). Activation of R genes leads to reactive oxygen production and the hypersensitive response (HR), which is considered a programmed cell death (PCD) response to impede expansion of the pathogen^{7–9}. During the interaction between plants and pathogens, R genes play significant roles in the HR; the SA-dependent defense signaling pathway is invoked when the effector is recognized by an R gene^{10,11}. SA signaling is also an important aspect of the PAMP response and R gene-mediated resistance^{12,13}. Both ET and jasmonic acid, which are plant hormones, are crucial for the PCD response^{14–17}. Additionally, abscisic acid (ABA) plays a positive role in early-acting defenses, such as stomatal closure, but plays a negative role in later-acting defense mechanisms mediated by SA or JA^{18–20}. These immune responses are controlled by the reprogramming of a considerable number of defense-related genes, including those encoding transcription factors (TFs)^{21,22}.

The plant immune response is closely related to the calcium ion (Ca^{2+}) signal transduction pathway via the decoding of the Ca^{2+} signatures triggered by pathogens^{23–27}. Different Ca^{2+} sensors have been reported to recognize Ca^{2+} signatures, such as calmodulin-like proteins (CMLs) and calmodulin (CaM), demonstrating that the conversion of Ca^{2+} signals to plant immune responses is mediated by different channels and machineries²⁸. As a general secondary messenger, Ca^{2+} participates in different types of cellular processes and has important functions in sensing microbe signals to establish appropriate immune and symbiotic responses, including the production of ROS and NO as well as PCD²⁶. Moreover, Ca^{2+} signaling is connected to other hormone-mediated pathways. In guard cells, stomatal closure in response to increased levels of Ca^{2+} is impaired in *cpk5/6/11/23* mutants treated with ABA²⁹. Research also shows that Ca^{2+} is involved in the ET-mediated pathogenesis response³⁰. Moreover, many TFs have been identified as being involved in calcium signaling. One class of TFs comprising Ca^{2+} /CaM-

regulated members is the WRKY family; WRKY43, WRKY53, and all members of the WRKYIIId subfamily have been found to interact with Ca^{2+} /CaM³¹. Although the Ca^{2+} signal transduction pathway has been a popular research topic for many years, it remains unknown whether Ca^{2+} is involved in Chrysanthemum-*A. alternata* interaction.

RNA sequencing technology is an efficient method for revealing genetic diversity and comparing variation in global expression of coding genes among genotypes/cultivars. Several transcriptome studies involving various genotypes/cultivars have been performed to understand stress responses³². Nevertheless, no such detailed analysis has been performed to elucidate the molecular regulatory mechanisms underlying the response of both resistant and susceptible Chrysanthemum cultivars to *A. alternata* infection.

Here, we dissect RNA sequencing time series data to reveal the transcriptional regulatory network in Chrysanthemum leaves after *A. alternata* inoculation. Our integrated analysis revealed that EDS1-mediated disease resistance and hormone and Ca^{2+} signal transduction pathways are of vital importance in the response of Chrysanthemum to infection of this fungus. Differentially expressed gene (DEG) and coexpression network analyses revealed a series of hub genes that play key roles at different stages of *A. alternata* infection. To verify the function of these hub genes, we performed transgenic experiments with *CmWRKY33.1* and acquired sense-expression genetic transformation lines. Inoculation assays showed that three lines overexpressing *CmWRKY33.1* exhibited enhanced susceptibility to black spot disease compared with that of wild-type 'Jinba'. Our research presents insight into the molecular regulatory mechanisms underlying Chrysanthemum responses to *A. alternata* inoculation.

Results

Evaluation of Chrysanthemum cultivars after inoculation with *A. alternata*

We selected two Chrysanthemum cultivars, 'Jinba' (resistant cultivar) and 'Zaoyihong' (susceptible cultivar), which differ significantly in their symptoms after *A. alternata* inoculation. The black leaf spot disease symptoms of the two Chrysanthemum cultivars were investigated at 3 days post inoculation. 'Jinba (SM)' displayed necrotic lesions that were mostly limited to the inoculated leaf, whereas 'Zaoyihong (ZYH)' exhibited necrotic lesions on other leaves, including the leaf inoculated with *A. alternata* (Fig. 1a). The disease symptoms were evaluated based on the proportion of diseased leaves at 3 days post inoculation; as shown in Fig. 1b, the proportions of diseased 'Jinba' and 'Zaoyihong' leaves were 0.08–0.10 and 0.45–0.56, respectively.

Leaves inoculated with *A. alternata* were sampled at six time points (0 h, 1 h, 6 h, 12 h, 24 h, and 36 h) based on the phenotypes to analyze differences in transcript levels between 'Jinba' and 'Zaoyihong', with three biological replicates sampled for each cultivar at each time point.

Generation of transcriptomic time series data via RNA sequencing

RNA sequencing technology is an effective method for revealing genetic diversity and comparing the variation in global expression of coding genes among various genotypes/cultivars. To further investigate the transcriptome dynamics and regulatory mechanism of Chrysanthemum leaf responses to *A. alternata*, we performed a comparative transcriptome-wide time series analysis of the two Chrysanthemum cultivars, 'Zaoyihong (ZYH)' and 'Jinba (SM)', with three independent biological replicates for all samples at each time point. In total, 36 cDNA libraries from the leaf tissues of the two cultivars were generated: SM_0_1, SM_0_2, SM_0_3, SM_1_1, SM_1_2, SM_1_3, SM_6_1, SM_6_2, SM_6_3, SM_12_1, SM_12_2, SM_12_3, SM_24_1, SM_24_2, SM_24_3, SM_36_1, SM_36_2, SM_36_3, ZYH_0_1, ZYH_0_2, ZYH_0_3, ZYH_1_1, ZYH_1_2, ZYH_1_3, ZYH_6_1, ZYH_6_2, ZYH_6_3, ZYH_12_1, ZYH_12_2, ZYH_12_3, ZYH_24_1, ZYH_24_2, ZYH_24_3, ZYH_36_1, ZYH_36_2, and ZYH_36_3. The raw data were then converted to clean reads, which were subsequently used for further analysis (Table S1). As a result, 1.7 billion clean reads (average of ~23.3 million reads from each sample) were obtained. All clean data can be downloaded from NCBI (BioProjects: PRJNA596929).

Analysis of DEGs in the time series samples

RSEM and BOWTIE were used to process the clean reads for each sample to obtain the expected FPKM (fragments per kilobase of transcript length per million

mapped reads) number for each gene. The DESeq R package was used to identify differentially expressed genes. Pearson correlation analysis was then applied to compare the RNA sequencing data of the three biological replicates, which varied from 0.87 to 0.93, indicating high consistency among the replicates (Fig. S1).

To investigate the temporal dynamics of transcriptional reprogramming between the two Chrysanthemum cultivars following challenge with *A. alternata*, we plotted the number of DEGs ($\text{padj} < 0.05$) against those of non-inoculated plants (SM_0 or ZYH_0) at each inoculation time point. In the 'Jinba' cultivar, 5740 (2835 upregulated; 2905 downregulated), 19,691 (12,176 upregulated; 7515 downregulated), 13,805 (8868 upregulated; 4937 downregulated), 25,549 (16,315 upregulated; 9234 downregulated) and 25,375 (15,543 upregulated; 9732 downregulated) DEGs were found in the samples at the five inoculation times; 7104 (3352 upregulated; 3752 downregulated), 12,010 (7129 upregulated; 4881 downregulated), 17,662 (11,277 upregulated; 6385 downregulated), 9813 (7172 upregulated; 2641 downregulated) and 18,826 (11,816 upregulated; 7010 downregulated) DEGs were identified in 'Zaoyihong' (Fig. 2a). The largest number of stage-specific genes (25,549 in 'Jinba' and 18,826 in 'Zaoyihong') was identified in the comparison of SM_24 and ZYH_36 compared with that of SM_0 and ZYH_0 (Fig. 2a). In total, 39,802 DEGs were obtained from the two cultivars; of these, 2084 DEGs were shared among the five time series stages in 'Jinba', and 3144 DEGs were shared in 'Zaoyihong'. We also identified DEGs uniquely expressed at each time series stage, as shown in Fig. 2b, and the results suggest that each stage has its own independent defense mechanisms.

The results of our study showed that a large number of DEGs, up to 33,343 in the resistant cultivar 'Jinba' and 26,173 in the susceptible cultivar 'Zaoyihong', respond to *A. alternata* infection. This is a considerably greater

Fig. 2 Statistical and GO analyses of DEGs in 'Jinba' and 'Zaoyihong' after infection with *A. alternata*. **a** Statistical analysis of upregulated and downregulated differentially expressed genes (DEGs) in response to *A. alternata* infection. **b** Venn diagrams of upregulated and downregulated differentially expressed genes (DEGs) in response to *A. alternata* infection in 'Zaoyihong' (top) and 'Jinba' (bottom). **c** GO analysis of DEG responses to *A. alternata* infection in 'Jinba' and 'Zaoyihong'. The enriched categories are represented by circles, and the size of each circle represents the number of genes

number of genes than previously reported in similar transcriptomic studies. By using BINGO, we next performed GO analysis to further characterize the identified DEGs. As shown in Fig. 2c, the DEGs were distributed among the three GO categories: "biological process", "molecular function", and "cellular component". The DEGs enriched in the "cellular component" category are mainly involved in "cell wall", "peroxisome", "nucleoplasm", and "thylakoid". Enriched terms in the "biological process" category mainly included "response to stress", "response to endogenous stimulus", and "carbohydrate metabolic process". The GO terms "hydrolase activity", "kinase activity", "DNA-binding transcription factor activity", "carbohydrate binding", and "lipid binding" were enriched in the "molecular function" category.

Pathway analysis of Chrysanthemum in response to *A. alternata* infection

To facilitate our analysis of the plant and fungal networks, we also mapped the pathways of the DEGs involved in Chrysanthemum-*A. alternata* interactions using the KEGG database (www.genome.jp/kegg). Plant hormones are crucial for helping plants adapt to adverse environmental conditions. Our data showed that the expression of genes involved in hormone signaling, especially that of ET and ABA, was upregulated when Chrysanthemum leaves

were challenged with *A. alternata* (Fig. S2b and c); the expression of a small proportion of genes involved in other hormone signal transduction pathways was also upregulated (Fig. S3). Traditionally, it has been thought that mainly ET and JA respond to necrotrophic fungi, but SA mediates defense signaling against both hemibiotrophic and biotrophic fungi. Based on our data, SA might also play a significant role in local immunity against necrotrophic fungi (Fig. S3). Moreover, plant-pathogen interaction pathway analysis indicated that the expression of genes involved in the Ca^{2+} -dependent signaling pathway was upregulated by more than 2-fold (Fig. S2a).

The transcriptional changes in the leaves of resistant and susceptible Chrysanthemum cultivars infected with *A. alternata* were also visualized via MapMan software. Figure S4 depicts the general overview of the cellular response, confirming that the number of DEGs responding to biotic stress was greater than the number responding to other stresses. Figure S5 illustrates a general overview of the biotic stress response. MapMan visualization revealed that many DEGs are involved in cell wall strengthening, transcriptional regulation, the oxidative respiratory burst, cellular signaling, and hormone metabolism. Upregulated genes play important roles in the response to pathogen infection. Moreover, the expression of most of the DEGs related to ET metabolism,

the respiratory burst and mitogen-activated protein kinases (MPKs) was upregulated at different times after *A. alternata* inoculation, which suggests the importance of these pathways for the resistance of Chrysanthemum to infection by this fungus.

In total, the involvement of multiple signal transduction pathway genes suggests that the response of Chrysanthemum leaves to *A. alternata* infection is not simply linear cascades but rather is a combination of multiple signaling pathways. Previous studies have shown that ET and the H₂O₂-mediated HR and PCD play vital roles in sand pear–*A. alternata* interaction³³. In our study, we identified that ET, ABA, and Ca²⁺ are important in the defense response to *A. alternata* inoculation and highlight the regulatory mechanism occurring in Chrysanthemum leaves.

Identification of gene coexpression modules during *A. alternata* infection

WGCNA is a powerful tool for identifying a set of genes associated with phenotypes. To investigate the gene regulatory network of Chrysanthemum leaves in response to *A. alternata* infection, we performed a WGCNA by using RNA sequencing profiles. After the expression data were preprocessed, a total of 36,403 genes were subjected to WGCNA, and we ultimately discovered 14 gene coexpression modules. The number of genes in these modules ranged from 62 to 8235.

The eigengenes for the 14 modules correlated with different infection stages (Fig. 3). We studied the preservation of coexpression modules between ‘Jinba’ and ‘Zaoyihong’ at different stages of *A. alternata* infection via module-trait relationship and module-sample relationship analyses. In most of the modules, the trend of genes was nearly the same between the two cultivars at different inoculation times, as indicated by the yellow, tan, and brown colors in the figure. Interestingly, a few modules differed in response to *A. alternata* infection. Notably, we identified one module of ‘Jinba’, i.e., that associated with the turquoise color in the figure, that was cultivar specific.

Transcriptional regulatory modules associated with the expansion of necrotic lesions

We investigated the transcriptome of Chrysanthemum leaves infected with *A. alternata* and collected samples at six time points. The early stage of infection was symptomless up to ~1 h post inoculation (hpi). The first signs of necrosis were observed at 6 hpi, and round or irregular lesions formed at 12 hpi at the inoculation sites, which was considered the middle stage/period of infection. During the late stage of infection, the lesions expanded to the outside of the inoculation site at 24 hpi, and half of the Chrysanthemum leaves showed black necrotic spots at 36 hpi. Plant defense genes play crucial roles during fungal

infection, and the expression of many is regulated by TFs. Therefore, we further analyzed the important defense genes and TFs with a strong regulatory relationship with the coexpressed defense genes in these notable modules.

Combined with our WGCNA, we identified key genes (also called hub genes) at different infection stages, as shown in Fig. 4. The expression of the genes in the yellow module were most upregulated at 1 hpi, corresponding to the early stage of the response of Chrysanthemum leaves to *A. alternata* infection. At 6–12 hpi, there was one tan module in which the gene expression showed peak upregulation at 6 hpi or 12 hpi. Many genes with a transcriptional peak at 24 hpi or 36 hpi were classified into the black or brown module; at 24 hpi or 36 hpi, the late stage, the lesions expanded outside of the inoculation site. In the yellow module, three hub genes, e.g., an *EDS1* homolog (*CmEDS1*; *DN78955c0_g1*), an *MLO1* homolog (*CmMLO1*; *DN86381c0_g1*), and an *NPR4* homolog (*CmNPR4*; *DN80377c1_g1*), related to defense were identified. Because TFs play important roles in gene regulation and plant immunity, we further investigated their regulatory network. TFs showed a strong regulatory relationship (weight/topological overlap matrix (TOM)>0.3) with other genes in the module and were used for further analysis. The identified *WRKY4* homolog showed a regulatory relationship with the three hub defense response genes. In addition, the expressions of the *RPS4* homolog and *PYL4* homolog were upregulated in the yellow module (Fig. 4a and Fig. S2c). In the tan module, we identified two hub genes related to plant-pathogen interactions: a *WRKY33* homolog (*CmWRKY33*; *DN63096c1_g1*) and a *CML45* homolog (*CmCML45*; *DN69937c2_g1*). An *ERF2* homolog (*CmERF2*; *DN90778c3_g1*), whose product plays important roles in the ET signal transduction pathway, was found to have a strong regulatory relationship with the two hub genes (*CmWRKY33* and *CmCML45*) (Fig. 4b). In the late stage of the Chrysanthemum response to *A. alternata* infection, we identified two hub genes belonging to the MATE family, which are involved in multiple drug and toxic compound extrusion; a *DTX30* homolog (*CmDTX30*; *DN96458c3_g7*); and a *DTX32* homolog (*CmDTX32*; *DN73836c0_g2*). One TF, a *WRKY6* homolog (*CmWRKY6*; *DN87555c0_g2*), was also identified as a hub gene. Furthermore, an *HSF24* homolog (*DN82662c0_g1*) and a *NAC029* homolog (*DN73063c1_g1*) exhibited strong regulatory relationships with other genes in the brown module, which also had regulatory relationships with two hub genes (*CmDTX30* and *CmDTX32*) (Fig. 4c). These results suggest a previously unknown function for MATE family genes in response to *A. alternata* infection.

Coexpression networks reveal that the highly resistant cultivar ‘Jinba’ uniquely responds to *A. alternata* infection

The Chrysanthemum cultivar ‘Jinba’ was more resistant and had smaller necrotic lesions than ‘Zaoyihong’ did during infection with *A. alternata* (Fig. 5a). The

Fig. 3 WGCNA of the transcripts in ‘Jinba’ and ‘Zaoyihong’. **a** Dendrogram of genes based on a coexpression analysis in ‘Jinba’ and ‘Zaoyihong’. Each leaf in the tree corresponds to an individual gene. Fourteen modules, each associated with a different color, formed from the major tree branches. **b** Association between modules and traits. The color of each module is the same as that in **a**. Each column represents a leaf necrotic lesion trait. The legend indicates the correlation. Each cell contains the number of corresponding correlations and *P* values

Fig. 4 (See legend on next page.)

(see figure on previous page)
Fig. 4 DEGs at different stages of the response of Chrysanthemum leaves to *A. alternata* infection. **a** Coexpression network of the yellow module. **b** Coexpression network of the tan module. **c** Coexpression network of the brown module. The coexpressed genes in each module are shown in heatmaps and bar graphs (left). The red rectangles in the heatmap represent high expression; the green rectangles represent low expression. The network of top hub genes is indicated by larger circles and the red color in the network (middle). Transcription factors with strong regulatory relationships (weight >0.3) in the yellow modules are indicated by larger circles and green color in the network. The dots around the circle represent other coexpressed genes indicated by small dots, and the relationships are connected by lines. The heatmaps (right) show the expression of the top hub genes and transcription factors that present strong regulatory relationships with the top hub genes

Fig. 5 Analysis of highly expressed genes in the resistant cultivar 'Jinba'. **a** Phenotype of Chrysanthemum leaves at 12 h post inoculation in 'Jinba' and 'Zaoyihong'. **b** The coexpressed genes are shown in heatmaps and bar graphs in the turquoise module. The red rectangles in the heatmap represent high expression; the green rectangles represent low expression. **c** The turquoise module was visualized by Cytoscape. Differently expressed transcription factors are indicated in larger red, blue, or purple quadrangles. Larger purple circles, triangles, or quadrangles represent differentially expressed genes involved in the plant-pathogen interaction pathway. Differentially expressed genes involved in the hormone signal transduction pathway are indicated in larger blue or purple triangles and larger blue quadrangles. **d** Heatmaps showing the expression profile of DEGs involved in PPI and HST and transcription factors in the turquoise module. The color represents the expression levels of genes in the module as shown in the figure legend

‘turquoise’ module ($r = 0.83$, $p = 9e-04$), which included 8228 genes, was cultivar specific and upregulated after *A. alternata* infection (Fig. 5b), with a possible association with the resistant phenotype.

The genes of this module were examined for further analysis. Based on the DEG analysis, we found 10 hub genes belonging to different TF families, e.g., *AP2-EREBP*, *AUX*, *bHLH*, *MADS*, *MYB*, *NAC*, *PLATZ*, and *TRAF*, to be differentially expressed (Fig. 5d). A total of 52 genes that interact with TFs according to the WGCNA are involved in plant-pathogen interactions, including *DN80856c2_g1*. Interestingly, *DN80856c2_g1*, called *PTI6*, which belongs to the *AP2-EREBP* TF family, also plays significant roles in plant-pathogen interactions. The results showed that the hub TFs may interact with genes involved in plant-pathogen interactions. Additionally, among the genes interacting with the hub TFs, 43 are involved in the hormone signal transduction pathway, which may be crucial in resistant cultivars challenged with *A. alternata*. The regulatory relationship was visualized with Cytoscape software, as depicted in Fig. 5c. Analysis of the DEGs interacting with TFs revealed one ABCA3 family gene (*DN91413c1_g3*) and four MATE family genes (*DN94014c1_g1*; *DN90910c1_g1*; *DN95310c2_g2* and *DN82887c0_g1*) that were distinctly highly expressed in the resistant cultivar ‘Jinba’ (Fig. 5d and Table S3), suggesting a previously unknown function in resistance to pathogen infection.

Validation of DEGs

To verify the RNA sequencing data, we selected 10 DEGs from different modules for quantitative real-time

PCR (qPCR) assays; the housekeeping gene *EF1α* was used as an internal control. The DEGs included *CmCML45* (*DN69937c2_g1*), *CmWRKY33* (*DN63096c1_g1*), *CmERF2* (*DN90778c3_g1*), *CmNAC029* (*DN73063c1_g1*), *CmMLO6* (*DN77454c0_g1*), *CmMYB15* (*DN77779c0_g1*), *CmHSF24* (*DN82662c0_g1*), *CmDTX24* (*DN81968c0_g1*), *CmFTSH* (*DN87341c1_g3*), and *CmVTE3* (*DN93367c1_g4*). The final qPCR data for the 10 DEGs were in accordance with the results of the transcriptome analysis, as presented in Fig. S6. Additionally, the qPCR and RNA sequencing results showed consistency according to a correlation analysis. The Pearson correlation coefficients of the results between the qPCR and RNA sequencing were 0.80–1 and 0.77–1 for the genes of ‘Jinba’ and ‘Zaoyihong’, respectively, at different stages of disease lesion development (Fig. S7), which also indicates that the data produced via RNA sequencing are reliable.

Overexpression of *CmWRKY33.1* resulted in enhanced susceptibility to black spot disease

In our study, we found that *WRKY* family genes were highly expressed as the lesion spread. Based on the *WRKY33* homologous sequences from our transcriptome data, we cloned one *WRKY33* homologous gene: *CmWRKY33.1*. To further verify the function of *CmWRKY33.1*, we constructed a pMDC43-*CmWRKY33.1* expression vector for genetic transformation of Chrysanthemum and acquired sense-expression transformed lines. Electrophoresis detection assays showed that the vector was successfully transformed into wild-type ‘Jinba’ (Fig. 6a), and according to qPCR assays, the expression of *CmWRKY33.1* in the transformed lines significantly

Fig. 6 Detection of *CmWRKY33.1* transgenic lines and phenotypes after inoculation with *A. alternata*. **a** Electrophoresis detection of transgenic lines. 3–68: Overexpressing lines; + Positive control (Plasmid DNA), – Negative control (H_2O), WT Wild-type ‘Jinba’. Up: PCR primers *CmWRKY33.1*-ID-F1/R1 were used for amplification of the pMDC43-*CmWRKY33.1* vector in the transformation lines. Down: PCR primers *CmWRKY33.1*-ID-F2/R2 were used for amplification of the pMDC43-*CmWRKY33.1* vector in the transformation lines. **b** Relative expression of *CmWRKY33.1* in the overexpression lines. **c** Phenotypic differences of wild-type and *CmWRKY33.1*-overexpressing lines. Bar = 10 mm

increased (Fig. 6b). Moreover, inoculation assays demonstrated that, compared with wild-type 'Jinba', the three transformation lines overexpressing *CmWRKY33.1* displayed enhanced susceptibility to black spot disease (Fig. 6c).

Discussion

The molecular mechanisms underlying the defense system of Chrysanthemum leaves in response to biotic stress are poorly understood. In our research, we identified a series of genes that respond to *A. alternata* at different times after inoculation in both susceptible and resistant Chrysanthemum cultivars and aimed to elucidate the mechanism by which the resistant cultivar 'Jinba' specifically engages its defense systems. RNA sequencing was applied to detect dynamic changes in the transcriptome of two Chrysanthemum cultivars (the resistant cultivar 'Jinba' and the susceptible cultivar 'Zaoyihong') and to investigate the molecular mechanism underlying lesion expansion. Analysis of the transcriptome time series and WGCNA revealed that pathogen-induced stress triggers defense responses in Chrysanthemum. Our research thus elucidated how Chrysanthemum conveys various signals in response to *A. alternata* infection.

When no lesions formed on Chrysanthemum leaves after *A. alternata* inoculation, the defense regulator *CmEDS1* plays significant roles in the response to infection, which is consistent with the involvement of *CmRPS4*, *CmNPR4*, and *CmPYL4*. In *Arabidopsis*, *EDS1* together with *RPS4* activates the TIR-NB-LRR signal transducer for defense across cellular compartments^{34–36}. Moreover, *EDS1* is an upstream regulator of the SA signaling pathway, and the *NPR4* gene codes for a protein that acts as an SA receptor^{37,38}. Our results showed that the *EDS1*-, *ABA*-, and *SA*-mediated disease resistance pathways in Chrysanthemum have crucial roles in the early response to *A. alternata* infection. In *Arabidopsis*, *WRKY4* acts as a positive regulator of plant resistance to necrotrophic pathogens³⁹. Among the *WRKY* target genes, *EDS1* and *NPR4* have been identified⁴⁰. *CmWRKY4* showed high expression during the early infection stage, which indicates that Chrysanthemum *CmWRKY4* may act as an upstream regulator of *CmEDS1*, collectively contributing to the early defense response of Chrysanthemum leaves to *A. alternata* infection.

From the display of no symptoms to the expansion of necrotic lesions on Chrysanthemum leaves after *A. alternata* inoculation, ET plays a central role in defense. The expression levels of all the ET signal transduction pathway genes (*CmETR1*, *CmCTR1*, *CmEIN2*, and *CmERF2*) were upregulated during this transition. WGCNA also demonstrated that another major transcription factor involved in Ca^{2+} signaling, *CmWRKY33*, has a strong regulatory relationship with *CmERF2*.

CmWRKY33 and *CmERF2* were highly expressed in the stage of necrotic lesion formation at inoculation sites and may act as important regulators in response to *A. alternata* infection. Ca^{2+} participates in different types of cellular processes as a general secondary messenger in plants, with important roles of sensing a series of extracellular stimuli and responding quickly²⁸. We also identified the gene encoding the calcium-binding EF-hand protein *CmCML45*, which is involved in the Ca^{2+} signaling pathway. In *Arabidopsis*, *WRKY33* is a significant regulator involved in the transcriptional regulation of metabolic and hormone responses to *Botrytis cinerea* infection⁴¹. Compared with that which occurred in the resistant wild type, SA levels were elevated and JA-associated responses were downregulated in susceptible *WRKY33* mutants after infection with *B. cinerea*⁴². As reported, *ERF2* appears to be positively controlled by *WRKY33*⁴³. Both *ChiB* and *PDF1.2* can be activated by *ERF2*, which has been shown to lead to resistance to a number of pathogens^{44,45}. In addition, *ACS2*, together with *ACS6*, is strongly correlated with the induction of ET, and *WRKY33* can bind to the W-box region in the promoters of *ACS2* and *ACS6*, the process of which is directly associated with MPK-induced expression of *ACS2* and *ACS6*⁴⁶. However, whether *CmWRKY33* is involved in ET signal transduction is still unknown. Our WGCNA showed that *CmCML45* and *CmWRKY33* have regulatory relationships with *CmERF2*, which was upregulated at 6 or 12 h after inoculation. *ERF2* is an important TF for the ET signal transduction pathway. ACC synthases (ACSs) convert S-adenosyl methionine to ACC, which is the direct precursor of ET⁴⁷. The results of our study also showed that two ACS homologous genes (*DN61354c0_g1* and *DN93750c1_g1*) were upregulated at the early stage of *A. alternata* infection. Ca^{2+} influx depends on plasma membrane-localized CNGC family proteins, which are involved in the uptake of Ca^{2+} and initiate immune signaling in plant cells^{31,48}. Two CNGC homologs (*DN96030c0_g1* and *DN93811c1_g2*) were also upregulated in the early stage. The above results show that Ca^{2+} signal transduction and the ET pathway play important roles in the early and middle stages of *A. alternata* infection in Chrysanthemum leaves.

In the late stage of *A. alternata* infection, necrotic lesions of Chrysanthemum leaves expanded, and a brown module was identified. However, no defense genes were identified among the hub genes. Interestingly, *CmDTX30* and *CmDTX32*, belonging to the MATE family, were identified hub genes in the brown module. Members of the MATE gene family play predominant roles in exporting toxins and other substrates^{49,50}, and MATE proteins involved in the response to abiotic stress have been reported^{51,52}. For example, *AtDTX50*, belonging to the MATE family, enhances drought tolerance in

*Arabidopsis thaliana*⁵³. However, the roles that members of the MATE gene family play in response to biotic stress remain unknown. In the brown module, *CmDTX30* and *CmDTX32* were highly upregulated at 24 or 36 h after inoculation, which indicated that some MATE family genes respond to *A. alternata* infection in the late stage. *Alternaria alternata* is a filamentous necrotrophic fungus that extracts nutrients from dead cells and produces different host-selective toxins that damage plant tissues^{54,55}; seven pathogenic variants of *A. alternata* have been identified, which can cause disease in a variety of plants by producing host-specific toxins². Based on the above studies, we speculate that Chrysanthemum leaves may transport toxins out of cells through multidrug resistance proteins, thereby reducing damage. Based on our results, we speculate that Chrysanthemum defends itself against *A. alternata* infection by reducing host-selective toxins via MATE family proteins. Additionally, ET signal transduction pathway genes exhibited peak upregulation at 24 hpi or 36 hpi, which indicated the important roles of ET in the late stage of the Chrysanthemum response to *A. alternata* infection.

Furthermore, 10 differentially expressed TFs were identified from the turquoise module, indicating that these proteins may play significant roles in the resistant cultivar 'Jinba'. Among the TFs expressed in the 'Jinba' cultivar, a PTI6 homolog (*CmPTI6*; *DN80856c2_g1*) was highly expressed in the resistant cultivar 'Jinba' and differentially expressed after inoculation. PTI6, an important ERF transcription factor, specifically interacts with Pto kinase and binds to the GCC-box cis element, which is an important regulator in mediating the expression of

pathogenesis-related (PR) genes⁵⁶. We investigated the expression of PR homologous genes in Chrysanthemum, and the results showed that *PR3*-like genes (*DN75226c1_g1*, *DN87821c0_g1*, and *DN88995c0_g2*) were upregulated in 'Jinba' after *A. alternata* inoculation, with higher expression in 'Jinba' than in 'Zaoyihong' at most of the times assessed (Table S3). The results indicated that the PTI6 homolog and *PR3*-like genes play important roles in the response to *A. alternata* infection in 'Jinba'.

Collectively, we mapped a model to elucidate the response of Chrysanthemum leaves to *A. alternata* infection at different stages and compared the model of the resistant cultivar 'Jinba' to that of the susceptible cultivar 'Zaoyihong' (Fig. 7). In the early stage, the calcium signal transduction pathway was activated in response to infection. With the transmission of calcium signals, more plant immune responses were activated, especially those involving the ET signaling pathway. Because *A. alternata* can produce a large number of toxins, we speculate that Chrysanthemum leaves may resist damage through MDR protein detoxification. In addition, we identified a series of hub genes including *PTI6*, and *PR3*-like genes, that were highly expressed specifically in response to *A. alternata* infection in the resistant cultivar 'Jinba'. Our study systematically identified the immunity map of Chrysanthemum in response to *A. alternata* infection, revealed the defense strategies of Chrysanthemum in different infection periods, and added to our understanding of the plant immune response. The identification of the hub genes of Chrysanthemum lays a foundation for elucidating the mechanism of plant responses to pathogen infection at

different periods. The hub genes identified from the resistant cultivar 'Jinba' provide a theoretical basis for breeding cultivars resistant to CBS.

Methods

Plant materials and *A. alternata* inoculation

Two Chrysanthemum cultivars ('Jinba' and 'Zaoyihong') and *A. alternata* were obtained from the Chrysanthemum Germplasm Resource Preserving Centre (Nanjing Agricultural University, China). Rooted seedlings were cultured in a 2:1 mixture of vermiculite and limestone. The seedlings were grown under a 16-h photoperiod with a day/night temperature of 25 °C and 18 °C, respectively, and the relative humidity was maintained at ~70%⁵⁷. Mycelia of *A. alternata* were cultured in liquid media. Mycelia collected from 1 mL of liquid media were placed on a Chrysanthemum leaf. After inoculation, the Chrysanthemum seedlings were grown in the dark for 36 h. Chrysanthemum leaves were subsequently sampled at 1, 6, 12, 24, and 36 h post inoculation. Uninoculated leaves of Chrysanthemum were used as the control (0 h), which were collected at the beginning of the experiment before inoculation. Leaves infected by *A. alternata* collected at defined time points were used for RNA sequencing, with three replicates for each treatment.

RNA extraction and sequencing

The total RNA from 36 samples was extracted according to previously reported methods⁵⁷. All 36 libraries were constructed and sequenced using the Illumina platform at Beijing Novogene (Tianjin, China) to generate 150-nucleotide-long paired-end sequence reads. Clean reads were obtained from the raw reads by removing adapter sequences and low-quality reads. Chrysanthemum unigenes were acquired using Trinity (v2.4.0) for de novo assembly of the transcriptome⁵⁸. The clean reads were then matched to the Chrysanthemum unigenes using RSEM software⁵⁹. The frequency of individual reads was normalized to FPKM⁶⁰.

DEG identification and GO enrichment analysis

DESeq was used for DEG analysis⁶¹. The padj (adjusted *P*-value) provides a criterion to determine the *P*-value threshold in multiple tests and analyses by manipulating the adjusted *P*-value. In our study, differential expression was considered when the absolute value of the log2 induction ratios of the treated samples compared with the mock treatment or control samples (0 h) was ≥ 1.0 and when padj < 0.05 . To obtain comprehensive information on the function of the unigenes, seven major databases, including the nonredundant, nucleotide, Pfam, COG, Swiss-Prot, GO, and KEGG databases, were used for gene annotations. GO enrichment analysis of the DEG sets was performed using BINGO⁶². The *P*-value providing a criterion to determine enrichment for each represented GO

term was calculated and corrected by the Benjamini and Hochberg false discovery rate (FDR). GO terms that showed corrected *P* values < 0.05 were considered significantly enriched. Pathway enrichment analysis was then performed to identify the series of genes involved in the response of 'Jinba' and 'Zaoyihong' to *A. alternata* infection using MapMan (version 3.5.1)⁶³.

Coexpression network analysis for the construction of modules

The R package for WGCNA was used for coexpression network analysis to evaluate modules in which the genes showed high correlations⁶⁴. An adjacency matrix was constructed based on normalized FPKM values. To identify modules related to phenotypic traits in the reconstruction network, the adjacency matrix was further converted to a topological overlap matrix (TOM) using the WGCNA package. After building the network, transcripts with similar expression patterns were classified into one module, and eigengenes (also called hub/key genes) for these modules were calculated. The TOM and degree were subsequently calculated; the top 20 genes according to degree were considered hub genes. Additionally, TOM > 0.3 was considered highly correlated and strongly regulatory. Coexpression networks were visualized using Cytoscape software⁶⁵.

qPCR validation

Ten genes from different modules were selected for quantitative real-time polymerase chain reaction (qPCR) analysis. Gene-specific primer pairs of the selected genes were designed using the primer design website (<https://www.yeastgenome.org/primer3>) (Table S2), and qPCR assays were performed as previously described⁶⁶. The Chrysanthemum *EF1 α* gene was used as a reference. Transcript abundances are given as the means $\pm$ SEs of three replicates. Relative transcription levels were calculated by using the $2^{-\Delta\Delta CT}$ method⁶⁷. Three biological replicates were performed for each sample.

Chrysanthemum *CmWRKY33.1* genetic transformation

Chrysanthemum plants were transformed with *Agrobacterium tumefaciens* strain EHA105 according to previously reported methods⁶⁸. The coding region of *CmWRKY33.1* was cloned by polymerase chain reaction (PCR). According to the multiple cloning sites within the Chrysanthemum genetic transformation expression vector pENTRTM 1A, two restriction endonuclease sites (*SalI* and *NotI*) were selected for vector construction. *CmWRKY33.1* was inserted into pMDC43 by homologous recombination. The resultant pMDC43-*CmWRKY33.1* vector was then used for genetic transformation, and hygromycin was used to select stable transformants. DNA and RNA were isolated from the wild-type and transformant lines, which were used

in PCR and qPCR experiments with constructed primers (Table S2) to identify the successfully transformed lines containing *CmWRKY33.1*.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (31700620), the Natural Science Fund of Jiangsu Province (BK20170722), and the Fundamental Research Funds for the Central Universities (KJQN201812).

Author contributions

Y.L. and F.C. designed the experiment. Y.L. and J.X. performed the experiment. Y.L. analyzed the data and wrote the manuscript, and F.C. revised the manuscript. All authors read and approved the final manuscript.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-0245-0>).

Received: 19 July 2019 Revised: 24 December 2019 Accepted: 4 January 2020

Published online: 01 March 2020

References

- Thomma, B. P. *Alternaria* spp. from general saprophyte to specific parasite. *Mol. Plant Pathol.* **4**, 225–236 (2003).
- Takaoka, S. et al. Complex regulation of secondary metabolism controlling pathogenicity in the phytopathogenic fungus *Alternaria alternata*. *N. Phytol.* **202**, 1297–1309 (2014).
- Jones, J. D. & Dangl, J. L. The plant immune system. *Nature* **444**, 323–329 (2006).
- Zipfel, C. Early molecular events in PAMP-triggered immunity. *Curr. Opin. Plant Biol.* **12**, 414–420 (2009).
- Tsuda, K. & Katagiri, F. Comparing signaling mechanisms engaged in pattern-triggered and effector-triggered immunity. *Curr. Opin. Plant Biol.* **13**, 459–465 (2010).
- Li, B., Meng, X., Shan, L. & He, P. Transcriptional regulation of pattern-triggered immunity in plants. *Cell Host Microbe* **19**, 641–650 (2016).
- Nimchuk, Z., Eulgem, T., Holt, B. F. 3rd & Dangl, J. L. Recognition and response in the plant immune system. *Annu. Rev. Genet.* **37**, 579–609 (2003).
- Collier, S. M. & Moffett, P. NB-LRRs work a "bait and switch" on pathogens. *Trends Plant Sci.* **14**, 521–529 (2009).
- Le Roux, C. et al. A receptor pair with an integrated decoy converts pathogen disabling of transcription factors to immunity. *Cell* **161**, 1074–1088 (2015).
- Vlot, A. C., Dempsey, D. A. & Klessig, D. F. Salicylic acid, a multifaceted hormone to combat disease. *Annu. Rev. Phytopathol.* **47**, 177–206 (2009).
- Baggs, E., Dagdas, G. & Krasileva, K. V. NLR diversity, helpers and integrated domains: making sense of the NLR identity. *Curr. Opin. Plant Biol.* **38**, 59–67 (2017).
- Zipfel, C. et al. Bacterial disease resistance in *Arabidopsis* through flagellin perception. *Nature* **428**, 764–767 (2004).
- Wang, L. et al. *Arabidopsis* CaM binding protein CBP60g contributes to MAMP-induced SA accumulation and is involved in disease resistance against *Pseudomonas syringae*. *PLoS Pathog.* **5**, e1000301 (2009).
- van Loon, L. C., Geraats, B. P. & Linthorst, H. J. Ethylene as a modulator of disease resistance in plants. *Trends Plant Sci.* **11**, 184–191 (2006).
- Zhang, L. & Xing, D. Methyl jasmonate induces production of reactive oxygen species and alterations in mitochondrial dynamics that precede photosynthetic dysfunction and subsequent cell death. *Plant Cell Physiol.* **49**, 1092–1111 (2008).
- Broekgaarden, C., Caarls, L., Vos, I. A., Pieterse, C. M. & Van Wees, S. C. Ethylene: traffic controller on hormonal crossroads to defense. *Plant Physiol.* **169**, 2371–2379 (2015).
- Luo, J. et al. Integrated transcriptome analysis reveals plant hormones jasmonic acid and salicylic acid coordinate growth and defense responses upon fungal infection in poplar. *Biomolecules* **9**, <https://doi.org/10.3390/biom9010012> (2019).
- Adie, B. A. et al. ABA is an essential signal for plant resistance to pathogens affecting JA biosynthesis and the activation of defenses in *Arabidopsis*. *Plant Cell* **19**, 1665–1681 (2007).
- Ton, J., Flors, V. & Mauch-Mani, B. The multifaceted role of ABA in disease resistance. *Trends Plant Sci.* **14**, 310–317 (2009).
- Cai, S. et al. Evolutionary conservation of ABA signaling for stomatal closure. *Plant Physiol.* **174**, 732–747 (2017).
- Ohme-Takagi, M., Suzuki, K. & Shinshi, H. Regulation of ethylene-induced transcription of defense genes. *Plant Cell Physiol.* **41**, 1187–1192 (2000).
- Singh, K., Foley, R. C. & Onate-Sanchez, L. Transcription factors in plant defense and stress responses. *Curr. Opin. Plant Biol.* **5**, 430–436 (2002).
- Dolmetsch, R. E., Xu, K. & Lewis, R. S. Calcium oscillations increase the efficiency and specificity of gene expression. *Nature* **392**, 933–936 (1998).
- Poovaiah, B. W., Du, L., Wang, H. & Yang, T. Recent advances in calcium/calmodulin-mediated signaling with an emphasis on plant-microbe interactions. *Plant Physiol.* **163**, 531–542 (2013).
- Gust, A. A. & Felix, G. Receptor like proteins associate with SOBIR1-type of adaptors to form bimolecular receptor kinases. *Curr. Opin. Plant Biol.* **21**, 104–111 (2014).
- Yuan, P., Jauregui, E., Du, L., Tanaka, K. & Poovaiah, B. W. Calcium signatures and signaling events orchestrate plant-microbe interactions. *Curr. Opin. Plant Biol.* **38**, 173–183 (2017).
- Zipfel, C. & Oldroyd, G. E. Plant signalling in symbiosis and immunity. *Nature* **543**, 328–336 (2017).
- Edel, K. H., Marchadier, E., Brownlee, C., Kudla, J. & Hetherington, A. M. The evolution of calcium-based signalling in plants. *Curr. Biol.* **27**, R667–R679 (2017).
- Brandt, B. et al. Calcium specificity signaling mechanisms in abscisic acid signal transduction in *Arabidopsis* guard cells. *Elife* **4**, <https://doi.org/10.7554/eLife.03599> (2015).
- Raz, V. & Fluhr, R. Calcium requirement for ethylene-dependent responses. *Plant cell* **4**, 1123–1130 (1992).
- Reddy, A. S., Ali, G. S., Celesnik, H. & Day, I. S. Coping with stresses: roles of calcium- and calcium/calmodulin-regulated gene expression. *Plant Cell* **23**, 2010–2032 (2011).
- Zhao, N. et al. Comparative Transcriptome analysis of waterlogging-sensitive and waterlogging-tolerant *Chrysanthemum morifolium* cultivars under waterlogging stress and reoxygenation conditions. *Int. J. Mol. Sci.* **19**, <https://doi.org/10.3390/ijms19051455> (2018).
- Wang, H., Lin, J., Chang, Y. & Jiang, C. Z. Comparative transcriptomic analysis reveals that ethylene/H₂O₂-mediated hypersensitive response and programmed cell death determine the compatible interaction of sand pear and *Alternaria alternata*. *Front. Plant Sci.* **8**, 195 (2017).
- Wirthmueller, L., Zhang, Y., Jones, J. D. & Parker, J. E. Nuclear accumulation of the *Arabidopsis* immune receptor RPS4 is necessary for triggering EDS1-dependent defense. *Curr. Biol.* **17**, 2023–2029 (2007).
- Heidrich, K. et al. *Arabidopsis* EDS1 connects pathogen effector recognition to cell compartment-specific immune responses. *Science* **334**, 1401–1404 (2011).
- Huang, P. Y., Catinot, J. & Zimmerli, L. Ethylene response factors in *Arabidopsis* immunity. *J. Exp. Bot.* **67**, 1231–1241 (2016).
- Tsuda, K., Sato, M., Stoddard, T., Glazebrook, J. & Katagiri, F. Network properties of robust immunity in plants. *PLoS Genet.* **5**, e1000772 (2009).
- Cui, H. et al. Antagonism of transcription factor MYC2 by EDS1/PAD4 complexes bolsters salicylic acid defense in *Arabidopsis* effector-triggered immunity. *Mol. Plant* **11**, 1053–1066 (2018).
- Lai, Z., Vinod, K., Zheng, Z., Fan, B. & Chen, Z. Roles of *Arabidopsis* WRKY3 and WRKY4 transcription factors in plant responses to pathogens. *BMC Plant Biol.* **8**, 68 (2008).
- Birkenbihl, R. P., Kracher, B., Roccaro, M. & Somssich, I. E. Induced genome-wide binding of three *Arabidopsis* WRKY transcription factors during early MAMP-triggered immunity. *Plant Cell* **29**, 20–38 (2017).
- Liu, S., Kracher, B., Ziegler, J., Birkenbihl, R. P. & Somssich, I. E. Negative regulation of ABA signaling by WRKY33 is critical for *Arabidopsis* immunity towards *Botrytis cinerea* 2100. *Elife* **4**, e07295 (2015).
- Liu, S., Ziegler, J., Zeier, J., Birkenbihl, R. P. & Somssich, I. E. *Botrytis cinerea* B05.10 promotes disease development in *Arabidopsis* by suppressing WRKY33-mediated host immunity. *Plant Cell Environ.* **40**, 2189–2206 (2017).

43. Birkenbihl, R. P., Diezel, C. & Somssich, I. E. *Arabidopsis* WRKY33 is a key transcriptional regulator of hormonal and metabolic responses toward *Botrytis cinerea* infection. *Plant Physiol.* **159**, 266–285 (2012).
44. Brown, R. L., Kazan, K., McGrath, K. C., Maclean, D. J. & Manners, J. M. A role for the GCC-box in jasmonate-mediated activation of the PDF1.2 gene of *Arabidopsis*. *Plant Physiol.* **132**, 1020–1032 (2003).
45. Meng, X. et al. Phosphorylation of an ERF transcription factor by *Arabidopsis* MPK3/MPK6 regulates plant defense gene induction and fungal resistance. *Plant Cell* **25**, 1126–1142 (2013).
46. Li, G. et al. Dual-level regulation of ACC synthase activity by MPK3/MPK6 cascade and its downstream WRKY transcription factor during ethylene induction in *Arabidopsis*. *PLoS Genet.* **8**, e1002767 (2012).
47. Dubois, M., Van den Broeck, L. & Inze, D. The pivotal role of ethylene in plant growth. *Trends Plant Sci.* **23**, 311–323 (2018).
48. Ma, W. et al. Ca²⁺, cAMP, and transduction of non-self perception during plant immune responses. *Proc. Natl Acad. Sci. USA* **106**, 20995–21000 (2009).
49. He, X. et al. Structure of a cation-bound multidrug and toxic compound extrusion transporter. *Nature* **467**, 991–994 (2010).
50. Lu, P. et al. Genome-wide analysis of multidrug and toxic compound extrusion (MATE) family in *Gossypium raimondii* and *Gossypium arboreum* and its expression analysis under salt, cadmium, and drought stress. *G3* **8**, 2483–2500 (2018).
51. Liu, J., Li, Y., Wang, W., Gai, J. & Li, Y. Genome-wide analysis of MATE transporters and expression patterns of a subgroup of MATE genes in response to aluminum toxicity in soybean. *BMC Genomics* **17**, 223 (2016).
52. Li, N. et al. Genome-wide analysis of MATE transporters and molecular characterization of aluminum resistance in *Populus*. *J. Exp. Bot.* **68**, 5669–5683 (2017).
53. Zhang, H. et al. A DTX/MATE-type transporter facilitates abscisic acid efflux and modulates ABA sensitivity and drought tolerance in *Arabidopsis*. *Mol. Plant* **7**, 1522–1532 (2014).
54. Estiarte, N., Lawrence, C. B., Sanchis, V., Ramos, A. J. & Crespo-Sempere, A. LaeA and VeA are involved in growth morphology, asexual development, and mycotoxin production in *Alternaria alternata*. *Int. J. Food Microbiol.* **238**, 153–164 (2016).
55. Sanchez-Lopez, A. M. et al. *Arabidopsis* responds to *Alternaria alternata* volatiles by triggering plastid phosphoglucose isomerase-independent mechanisms. *Plant Physiol.* **172**, 1989–2001 (2016).
56. Gu, Y. Q. et al. Tomato transcription factors *pti4*, *pti5*, and *pti6* activate defense responses when expressed in *Arabidopsis*. *Plant Cell* **14**, 817–831 (2002).
57. Li, H. et al. RNA-Seq derived identification of differential transcription in the Chrysanthemum leaf following inoculation with *Alternaria tenuissima*. *BMC Genomics* **15**, 9 (2014).
58. Grabherr, M. G. et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nat. Biotechnol.* **29**, 644–652 (2011).
59. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC Bioinform.* **12**, 323 (2011).
60. Trapnell, C. et al. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**, 511–515 (2010).
61. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol.* **11**, R106 (2010).
62. Maere, S., Heymans, K. & Kuiper, M. BiNGO: a Cytoscape plugin to assess overrepresentation of gene ontology categories in biological networks. *Bioinformatics* **21**, 3448–3449 (2005).
63. Thimm, O. et al. MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic pathways and other biological processes. *Plant J.* **37**, 914–939 (2004).
64. Zhang, B. & Horvath, S. A general framework for weighted gene co-expression network analysis. *Stat. Appl. Genet. Mol. Biol.* **4**, Article17 (2005).
65. Stuart, J. M., Segal, E., Koller, D. & Kim, S. K. A gene-coexpression network for global discovery of conserved genetic modules. *Science* **302**, 249–255 (2003).
66. Song, A. et al. Comprehensive analysis of mitogen-activated protein kinase cascades in Chrysanthemum. *PeerJ* **6**, e5037 (2018).
67. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* **25**, 402–408 (2001).
68. Li, P. et al. The over-expression of a Chrysanthemum WRKY transcription factor enhances aphid resistance. *Plant Physiol. Biochem.* **95**, 26–34 (2015).