ARTICLE

Open Access

UGT74AF3 enzymes specifically catalyze the glucosylation of 4-hydroxy-2,5-dimethylfuran-3(2H)-one, an important volatile compound in *Camellia sinensis*

Yongxian Chen¹, Xiangyang Guo¹, Ting Gao¹, Na Zhang¹, Xiaochun Wan¹, Wilfried Schwab^{1,2} and Chuankui Song¹

Abstract

4-Hydroxy-2,5-dimethylfuran-3(2H)-one (HDMF) is an important odorant in some fruits, and is proposed to play a crucial role in the caramel-like notes of some teas. However, its biosynthesis and metabolism in tea plants are still unknown. Here, HDMF glucoside was unambiguously identified as a native metabolite in tea plants. A novel glucosyltransferase UGT74AF3a and its allelic protein UGT74AF3b specifically catalyzed the glucosylation of HDMF and the commercially important structural homologues 2 (or 5)-ethyl-4-hydroxy-5 (or 2)-methylfuran-3(2H)-one (EHMF) and 4-hydroxy-5-methylfuran-3(2H)-one (HMF) to their corresponding β -D-glucosides. Site-directed mutagenesis of UGT74AF3b to introduce a single A456V mutation resulted in improved HDMF and EHMF glucosylation activity and affected the sugar donor preference compared with that of the wild-type control enzyme. The accumulation of HDMF glucoside was consistent with the transcript levels of UGT74AF3 in different tea cultivars. In addition, transient *UGT74AF3a* overexpression in tobacco significantly increased the HDMF glucoside contents, and downregulation of *UGT74AF3* transcripts in tea leaves significantly reduced the concentration of HDMF glucoside compared with the levels in the controls. The identification of HDMF glucoside in the tea plant and the discovery of a novel-specific UDP-glucose:HDMF glucosyltransferase in tea plants provide the foundation for improvement of tea flavor and the biotechnological production of HDMF glucoside.

Introduction

Plants produce a multitude of volatile compounds that play key roles in biological interactions^{1,2}. Tea plants synthesize, accumulate, and emit many volatile compounds, which are characteristic metabolites and play an important role in the quality of tea. The formation of volatile constituents in various teas during the manufacturing process has been intensively studied, and ~700 volatiles have been identified, including esters, aldehydes, ketones, alcohols, terpenes, and furanones^{3,4}. Among

them, 4-hydroxy-2,5-dimethylfuran-3(2H)-one (HDMF) is considered an exceptional volatile because of its low-odor threshold in some fruits, attractive odor qualities, and remarkable odor-enhancement effects^{5–8}. HDMF has a strong caramel-like aroma similar to that of its structural homologues 4-hydroxy-5-methyl-3(2H)-furanone (HMF, norfuranol) and 2 (or 5)-ethyl-4-hydroxy-5 (or 2)-methyl-3(2H)-furanone (EHMF, homofuranol). HDMF has been identified as a prominent volatile in pineapples⁹, strawberries¹⁰, grapes¹¹, and tomatoes¹², and it has also been found in black tea, where it contributes to the sweet and caramel-like odors^{4,13,14}.

Glycosylation is a major plant metabolite modification that plays an important role in many aspects, such as detoxification, transportation, and storage¹⁵. In addition, glycosylation also ensures the chemical stability and water

Correspondence: Chuankui Song (sckfriend@163.com)

¹State Key Laboratory of Tea Plant Biology and Utilization, International Joint Laboratory on Tea Chemistry and Health Effects, Anhui Agricultural University, 230036 Hefei, Anhui, P. R. China

²Biotechnology of Natural Products, Technische Universität München, Liesel-Beckmann-Str. 1, 85354 Freising, Germany

© The Author(s) 2020

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

solubility of plant metabolites while reducing chemical reactivity and toxicity¹⁶. Glycosylation also contributes to the storage of secondary metabolites and transport in plants¹⁷. Glycosylation is strictly controlled by glycosyltransferases (UGTs), which catalyze the transfer of activated sugar molecules to receptors to form corresponding glycosides. Although the study of glycosyltransferases is of great significance, there are few studies on glycosyltransferases, especially those related to volatiles¹⁵. In mature strawberry fruits, a large amount of HDMF was transformed into HDMF β -D-glucoside and accumulated^{18,19}. The glucosylation of HDMF affects the development of plant volatiles, because HDMF glucoside is less volatile than HDMF and odorless^{7,20}. Several *UGT* genes were isolated from strawberries whose encoded proteins catalyze the glucosylation of HDMF, including *FaGT2*²¹, *UGT71K3*²², and *UGT85K16*²³. One glycosyltransferase gene, *UGT85K14*, was isolated from a grapevine cultivar¹¹.

More than 300 UGTs were found in the tea plant genome^{24,25}, but the function of most of them is unknown, except for that of four UGTs (*UGT78A14*, *UGT78A15*, *UGT82A22*, and *UGT73A20*), which exhibited catalytic activity toward phenolic acids and flavonoid^{26,27}, and three UGTs involved in the glycosylation of geraniol²⁸ and (Z)-3-hexen-1-ol². However, the genes coding for HDMF glycosyltransferases in tea plants are not yet known.

In this study, HDMF glucoside was unambiguously identified as a native metabolite in tea plants by comparison of its retention time and accurate mass with those of an authentic standard. In addition, a novel glycosyltransferase *UGT74AF3a* and its allelic protein *UGT74AF3b* were identified to catalyze the glucosylation of HDMF and EHMF. The site-directed mutagenesis of *UGT74AF3b* led to the identification of one amino acid at the C-terminal end that influenced HDMF and EHMF glucosylation activity and sugar-donor preference compared with that of the wild-type control enzyme. In addition, the correlation between *UGT74AF3* transcript level and HDMF glucoside accumulation in different tea cultivars was studied. The *in planta* function of *UGT74AF3* was assessed by overexpression and silencing in tobacco and tea leaves, respectively. The identification of HDMF glucoside in the tea plant and the discovery of a novel-specific UDP-glucose:HDMF glycosyltransferase provides the foundation for improvement of tea flavor and the biotechnological production of HDMF glucoside.

Materials and methods

Plant material

Several cultivars of tea plants, including *Camellia sinensis* var. *sinensis* “Shuchazao”, “Yingshuang”, “Minxuan213”, “Zhenghedabai”, “Huangqi”, “Mingshanbaihao”, and “Fudingdabai”, were collected from the Tea

Plant Cultivar and Germplasm Resource Garden of Anhui Agricultural University (Guohe Town) and frozen in liquid nitrogen. All of the tea samples were stored at -80°C until use.

Chemicals and reagents

4-Hydroxy-2,5-dimethylfuran-3(2H)-one (HDMF), 2 (or 5)-ethyl-4-hydroxy-5 (or 2)-methylfuran-3(2H)-one (EHMF), 4-hydroxy-5-methylfuran-3(2H)-one (HMF), naringenin, eugenol, vanillic acid, salicylic acid, 1-naphthol, farnesol, sorbic acid, nerolidol, geraniol, fisetin, vanillin, chlorogenic acid, 2-naphthol, caffeic acid, methyl salicylate, ferulic acid, 1-octanol, and myricetin were purchased from Sigma-Aldrich (Shanghai, China). UDP-glucose, UDP-galactose, and UDP-glucuronic acid were purchased from Promega (Madison, USA). UDP-Xylose was purchased from Angfei Biotech Co., Ltd. (Guangzhou, China).

RNA isolation, cDNA cloning, and sequence analysis

For analysis of gene expression, the leaves of *Camellia sinensis* var. *sinensis* “Shuchazao” were collected for RNA extraction. The total RNA was isolated by RNAiso-mate for plant tissue (Takara, Dalian, China) and RNAiso Plus (Takara, Dalian, China). The cDNAs were amplified using Phusion® High-Fidelity DNA Polymerase (New England Biolabs, MA, USA), and the PCR products were purified via a MiniBEST agarose gel extraction kit (Takara, Dalian, China). The amplified PCR product was ligated into the pGEX-4T1 vector and transformed into TransT1-competent cells for sequencing.

Quantitative real-time PCR analysis

Real-time PCR was performed according to our published protocols² with gene-specific primers (Supplementary Table S1). qRT-PCR was carried out with the CFX96™ real-time system (Bio-Rad, USA). The two-step temperature program was 95°C for 3 min, 40 cycles of 95°C for 10 s, and 62°C for 30 s in 96-well optical reaction plates. The glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene was used as an internal reference gene², and the relative gene expression was calculated using the $2^{-\Delta\Delta\text{CT}}$ method²⁹.

Heterologous protein expression and purification

The full-length sequence of *UGT74AF3* was digested with BamH1 and Sma1, and the resulting gene fragments were subcloned into the expression vector pGEX-4T-1. The recombinant plasmids were transformed into *E. coli* strain BL21 (DE3) pLysS cells. The empty expression vector pGEX-4T-1 was transformed into *E. coli* BL21 (DE3) pLysS cells, which served as the negative control. Protein expression was induced by adding IPTG (isopropyl- β -D-thio-galactopyranoside) at a final concentration of 1 mM

when the OD₆₀₀ of the culture was between 0.6 and 0.8. The cultures were incubated at 16 °C with oscillation at 150 rpm overnight. The next day, the proteins were purified using GST-binding resin (Novagen, Darmstadt, Germany) following the manufacturer's protocol. The protein concentration was determined by the Bradford method, and bovine serum albumin served as the standard protein. The protein purity was further confirmed by SDS-PAGE.

Enzymatic activity assay

The enzyme assays were carried out according to the methods described previously^{2,21}. In the initial screening, the total volume of the enzyme reaction mixture (25 µL) contained 20 µL of Tris-HCl buffer (50 mM, pH 7.5, 10% glycerol, and 10 mM 2-mercaptoethanol), 1 µL of 2.5 mM UDP-glucose, 1 µL of 20 µM substrate solution, 1 µL of 50 mM DL-dithiothreitol, and 2 µL of purified protein (3 µg per reaction). The recombinant enzyme assay was carried out using the UDP-GloTM glycosyltransferase assay kit³⁰. The optimum reaction temperature was determined to be in the range of 10–45 °C at pH 7.0. The optimum reaction pH was tested in the range from pH 3 to pH 10. Citrate, phosphate, and Tris-HCl buffers were used for the pH ranges 3–6, 6–8, and 8–10, respectively. To determine kinetic parameters, at least seven different substrate concentrations covering the range from 1 to 500 µM were used at the optimized conditions as previously described^{2,21}.

Identification of the products by LC–MS

The standard assays (200 µL) included 5 mM UDP-glucose, 200 µM substrate, and 10 to 20 µg purified protein. The enzyme assays were incubated at 30 °C for 3 h. The reaction was stopped and extracted twice with 200 mL of ethyl acetate. After vaporizing the organic solvent, the residue was further dissolved in 50 µL methanol/water (1:1, vol) for LC–MS analysis. LC–MS was performed with a reverse-phase C18 column (1.8 µm, 100 × 2.1 mm) at 40 °C. A DIONEX Ultimate 3000 UHPLC system (Thermo Fisher Scientific, Waltham, MA, USA) with an autosampler was utilized for all experiments. The solvent selection and LC parameters for product identification were set according to our published paper². The products were identified by comparing their UV and MS spectra with those in the literature and reference material.

Site-directed mutagenesis

A single A456V mutation of UGT74AF3b was conducted. The primers were designed based on sequence data of UGT74AF3b (Supplementary Table S3). PCR amplifications were carried out using the following conditions: 1 cycle of 2 min at 98 °C; 25 cycles at 98 °C for

10 s, 58 °C for 30 s, and 72 °C for 3 min; and a final extension at 72 °C for 5 min. The mutant gene was confirmed by sequencing. Mutant proteins were expressed and purified using the same procedure described above for native proteins.

Construction and transformation of the overexpression plasmid

Full-length cDNA of *UGT74AF3a* from tea leaves was amplified by PCR using specific primers (Supplementary Table S2), and transferred to the expression vector pBI121. pBI121 with an intron containing the GUS gene was treated as the control. *Agrobacterium tumefaciens* strains AGL0 with pBI121-UGT74AF3a and pBI121 vectors were cultured at 28 °C in the LB medium containing appropriate antibiotics. When the bacteria reached an OD₆₀₀ of 0.8, the cells were harvested and resuspended in modified MMA medium (MS salts, 10 mM MES pH 5.6, and 20 g L⁻¹ sucrose). Transient experiments in tobacco leaves have been described previously².

Gene suppression of *UGT74AF3a* in tea using AsODNs

The previously described procedure for gene suppression in tea leaves was modified³¹. Candidate sequences of antisense oligonucleotides (AsODN) containing complementarity to the segment of the target gene were selected by using Soligo software³². AsODNs and special primers (Supplementary Table S4) were synthesized by Beijing Genomics Institute (BGI). To validate the suppression of *UGT74AF3a*, AsODN was used, and random nonsense ODNs were treated as the control. To determine the effects of *UGT74AF3a* gene suppression, naturally growing tender tea shoots containing two leaves were selected and put into Eppendorf tubes with 1 ml of a 20 µM AsODN-*UGT74AF3a* solution in this study.

Metabolite analysis

HDMF glycosides were extracted and measured using the protocol described previously with slight modification². For metabolite analysis, 50 mg of samples were extracted with 1 ml 75% (v/v) methanol twice. A chlorophenylalanine solution (3 µg mL⁻¹) was added as an internal standard. The metabolites were sonicated for 20 min at 4 °C. After that, the mixture was centrifuged at 12,000 rpm and 4 °C for 10 min. The supernatants were used for HDMF glycoside analysis by LC–MS as mentioned above. For metabolite analysis, at least three independent biological replicates were examined.

Phylogenetic tree and accession numbers

A phylogenetic tree was generated using a previously described method². The following accession numbers were utilized: UGT85K14 (LC021362), UGT85K11 (ABB847092), UGT85K16 (LC312711), UGT76D1

(At2g26480), CsUGT78A14 (ALO19888.1), CsUGT78A15 (ALO1989.1), CsUGT84A22 (ALO19890), FaGT2 (AY663785), CsUGT73A20 (ALO19886.1), UGT94P1 (ABB847093), UGT71K3 (XP_004294260.1), UGT71C3 (At1g07260), UGT74F1 (At2g43840), UGT74F2 (At2g43820), UGT74D1 (At2g31750), and UGT74E2 (At1g05680).

Results

Identification of HDMF glucoside in tea plants

To study whether HDMF glucosides were present in tea plants, the metabolites in young tissues of tea plants (*Camellia sinensis* var. *sinensis* “Shuchazao”) were extracted and analyzed by LC–MS. A peak with a retention time of 6.5 min was identified as HDMF glucoside by comparison of the retention time and accurate mass with those of an authentic standard (Fig. 1a, d). To prove that HDMF can be glucosylated in tea plants, undamaged tea leaves were exposed to 20 mM authentic HDMF in a closed glass vessel according to the method used in our previous research². LC–MS analysis confirmed that the concentration of HDMF glucoside significantly increased twofold after exposure to HDMF for 24 h (Fig. 1b). Furthermore, when 20 mM authentic HDMF was directly injected into the tea plant, the concentration of HDMF glucoside rapidly increased to more than tenfold that of the control (Fig. 1c). All these data proved that HDMF glucoside is a native metabolite in tea plants; the tea plants could glucosylate HDMF, whereby HDMF glucoside was produced by an as-yet unknown UGT in the tea plants.

UGT74AF3a and b specifically catalyze the glucosylation of HDMF

In *Arabidopsis*, 27 GTs, from ~107 enzymes, exhibited catalytic activity toward volatile compounds. The activities occurred in groups D, E, G, H, and L of the phylogenetic tree of the GT1 multigene family of *Arabidopsis*^{26,33}. To study UGTs related to HDMF glycosylation, the candidate genes TEA012191, TEA016995, TEA031670, and TEA016084 were randomly selected from groups E, G, H, and L, respectively. Genes from group D were not selected because some genes in this group have been reported to encode flavonoid glucosyltransferases in tea plants²⁷. TEA031670 could not be isolated as a full-length gene. The full-length sequences of TEA012191 and TEA016995 and two alleles of TEA016084 were obtained from the leaves of Shuchazao. The encoded UGTs were successfully expressed in *E. coli* with an N-terminal glutathione S-transferase (GST) tag.

The activities of recombinant proteins were first tested using the UDP-GLO glycosyltransferase assay with HDMF as the substrate. The recombinant protein encoded by TEA016084 catalyzed the formation of HDMF glucoside from HDMF and UDP-glucose, while the control vector

Fig. 1 HDMF glucoside was identified in tea plants. The identity of HDMF glucoside in untreated leaves (a), tea leaves exposed to 20 mM HDMF for 24 h (b), and tea leaves injected with HDMF (c) was confirmed by LC–MS. DMSO was used as a control

proteins and the proteins encoded by TEA012191 and TEA016995 could not form HDMF glucoside. Two alleles of TEA016084 were isolated. The encoded proteins were subsequently assigned CsUGT74AF3a and b by the UGT Nomenclature Committee³⁴ (Supplementary Fig. S2). The HDMF glucoside formed by both CsUGT74AF3a and b was identified by LC–MS (Fig. 2). The enzymatic activity of CsUGT74AF3a and b was further tested with 19 acceptor substrates (Fig. 3). The results indicated that the allelic UGT74AF3a and b proteins exhibited specific glucosylation activity toward HDMF (Figs. 2 and 3).

UGT74AF3a and b exhibit different sugar-donor preferences

In addition to HDMF, its structural homologues (EHMF and HMF) were also used as substrates for UGT74AF3a and b, which exhibited the ability to use HDMF, EHMF, and HMF as substrates; however, UGT74AF3a exhibited a higher glucosylation activity toward HDMF, EHMF, and HMF than that of UGT74AF3b when UDP-glucose was used as a donor substrate (Supplementary Fig. S3).

Sugar-donor specificity was further analyzed. UGT74AF3a preferred UDP-Glc (100%) as a sugar donor over UDP-Gal (70–85%) and UDP-GA (70–80%) when HDMF and EHMF were used as donor substrates (Supplementary Fig. S3). However, when HMF was used as the substrate, the sugar preference was less obvious (Supplementary Fig. S3). Interestingly, UGT74AF3b also preferred UDP-Glc (38–42%) as a sugar donor, but the catalytic activities toward UDP-Gal and UDP-GA were very low (<20%) compared with that of UGT74AF3a when HDMF, EHMF, and HMF were used as substrates (Supplementary Fig. S3). Both UGT74AF3a and b could not use UDP-UDP-Xylose as a sugar donor when HDMF, EHMF, and HMF were used as substrates (Supplementary Fig. S3).

Kinetic properties of the recombinant CsUGT74AF3a and b

The highest activity of both UGT74AF3a and UGT74AF3b was detected at pH 8.5 and 30 °C (Supplementary Fig. S1). The kinetic parameters of UGT74AF3a and b for HDMF were detected under the optimized conditions. The apparent K_M values of UGT74AF3a and b for HDMF were 1.59 and 5.14 μM , respectively (Table 1). The k_{cat}/K_M ratios of UGT74AF3a and b for HDMF were 40.66 and 28.77 $\mu\text{M}^{-1} \text{s}^{-1}$, respectively (Table 1).

Mutagenesis of UGT74AF3b at residue 456

Although the similarity of the amino acid sequences of UGT74AF3a and UGT74AF3b was 99.43% (Supplementary Fig. S2), their activities toward HDMF and their sugar-donor preferences were significantly different. To

Table 1 Kinetic data of recombinant UGT74AF3 for HDMF

Proteins	Substrate	K_M (μ M)	V_{max} (nKat/mg)	K_{cat}/K_M (μ M ⁻¹ s ⁻¹)
UGT74AF3a	HDMF	1.59 ± 0.19	4.23 ± 0.02	40.6
UGT74AF3b	HDMF	5.14 ± 1.58	11.38 ± 2.18	28.7

investigate the role of amino acids in sugar-donor preference, an A456V mutant of UGT74AF3b was produced. The A456V mutant exhibited enhanced glucosylation activity toward HDMF, EHMF, and HMF in comparison with that of the wild-type UGT74AF3b enzyme when UDP-gal and UDP-GA were used as sugar donors (Fig. 4). In addition, the results also revealed that the A456V mutant protein exhibited a stronger preference toward UDP-gal than UDP-glc (Fig. 4). This result indicated that the residue at A456V played a key role in the sugar-donor preference of UGT74AF3.

Transient overexpression of CsUGT74AF3a in tobacco

To further investigate the role of UGT74AF3a in tobacco, transgenic tobacco studies were carried out as previously described². Leaves were infiltrated using a suspension of *Agrobacterium tumefaciens* with empty pBI121-UGT74AF3a or pBI121 vectors, and were treated as the control. Relative mRNA levels of UGT74AF3a in *N. benthamiana* leaves were measured by two primers (Fig. 5a, b). The reaction products in both UGT74AF3a-overexpressing and pBI-intron control leaves were detected using LC-MS (Fig. 5d). The content of HDMF glucoside was markedly increased when *UGT74AF3a* was overexpressed (Fig. 5c), suggesting that HDMF can be formed in tobacco leaves, but the enzymes responsible for its glucosylation are not expressed. In addition, *UGT74AF3a* expression levels in individual tobacco leaves correlated very well with the concentration of HDMF glucoside (Fig. 5), showing that UGT74AF3 catalyzes HDMF glucoside production in tobacco.

Functional assessment of UGT74AF3 in tea plants

To determine the function of UGT74AF3 in HDMF catabolism in tea plants, the expression level of UGT74AF3 was transiently downregulated in *C. sinensis* leaves by gene-specific antisense oligodeoxynucleotide suppression according to Zhao et al.³¹. Three antisense oligodeoxynucleotides were designed (designated as 1, 2, and 3; Fig. 6). Tea plant leaves treated with sense oligodeoxynucleotides were used as a control. The expression level of *UGT74AF3* in tea leaves treated with AsODN-UGT74AF3 was significantly reduced compared with that in leaves of the control group, except for one sample treated with AsODN 3-1 (Fig. 6a). The content of HDMF

Fig. 4 The relative enzymatic activities analysis of UGT74AF3 and A456V mutant. Comparison of the relative enzymatic activities of UGT74AF3a and b and the A456V mutant with HDMF (a), EHMF (b), and HMF (c) as substrates. Mean separation analysis of substrates was calculated using one-way ANOVA test and Duncan's multiple-range test by SPSS 17.0. Means with different letters in each graph are significantly different ($P < 0.05$)

glucoside in *CsUGT74AF3*-silenced tea leaves was determined by LC-MS. The HDMF glucoside content was significantly reduced in the *CsUGT74AF3*-silenced tea leaves compared with that in the control (Fig. 6b). The results indicated that UGT74AF3 can catalyze HDMF glucosylation in tea plants.

Fig. 5 Overexpression of UGT74AF3a in tobacco. The relative mRNA levels of UGT74AF3a were measured by primer-1 (a) and primer-2 (b), and the HDMF glucoside product was analyzed by LC-MS (c, d). Duncan's multiple-range test was carried out, and statistical significance was calculated with one-way ANOVA using SPSS 17.0 ($P < 0.05$). CK empty vector control, OE overexpression of UGT74AF3a-OE

To further investigate the relationship between *UGT74AF3* gene expression and the concentration of glycosylation products of UGT74AF3a in different tea plant cultivars, several common tea cultivars were selected and analyzed. *UGT74AF3a* expression was detected in all cultivars, but the expression level of *UGT74AF3a* in Fudingdabai (FDDDB) was the lowest (Fig. 6c). The concentration of HDMF glucoside was measured by LC-MS. The compound was detected in all cultivars, but not in FDDDB (Fig. 6d). These results showed that there was a strong correlation between the mRNA levels of *UGT74AF3a* and the concentration of HDMF glucosides in different tea plant cultivars.

Discussion

Glycosylation of HDMF and phylogenetic analysis

HDMF is a key odorant in many fruits and was unambiguously identified as a native metabolite in tea plants in this study. In many aromatic plants, volatile compounds are stored as glycoside conjugates, which are water-soluble and odorless^{15,28}. With regard to the biosynthesis of HDMF glucoside, some genes coding for HDMF UGTs have already been isolated and identified. The UGT85 family members UGT85K14 and UGT85K16 possess substrate specificity for HDMF compounds, and were isolated from the hybrid grapevine cultivar Muscat Bailey

A (*Vitis labrusca* × *V. vinifera*) and strawberries (*Fragaria* × *ananassa*), respectively^{11,23}. Although structurally distinct from the UGT85K family, UGT71K3 in strawberries could also catalyze the glucosylation of HDMF and EDMF (identical to EHMF)²², in addition to acylphloroglucinol, quercetin, 1-naphthol, and pelargonidin^{22,35}. In tea plants, several UGTs have been cloned and functionally characterized, but only a few of them are involved in the glucosylation of volatiles. UGT85K11 from *Camellia sinensis* was recently reported to catalyze the formation of geranyl glucoside, and UGT94P1 specifically catalyzes the xylosylation of the 6'-hydroxy group of the sugar moiety of geranyl β-D-glucopyranoside and forms geranyl β-primeveroside²⁸. CsUGT85A53 catalyzes the formation of (Z)-3-hexen-1-ol glucoside in tea plants and might be involved in plant defense and plant-plant interaction². However, UGTs related to the glucosylation of HDMF in tea plants are still unknown. In this study, it was shown that UGT74AF3a and UGT74AF3b, which are members of the new UGT74AF subfamily, exhibit specific glucosylation activity toward HDMF, although they are structurally distinct from the UGT85K and UGT71 families. A previous study indicated that UGT85C2 and UGT74G1 could catalyze the glucosylation of steviol³⁶. The apparent K_M values of UGT74AF3a (1.59 μM) and b (5.14 μM) for HDMF were similar to that of UGT85K16

Fig. 6 Functional characterization of UGT74AF3 in tea plants. The expression levels of UGT74AF3 (**a, c**) and the accumulation of HDMF glucoside (**b, d**) in tea leaves after treatment with AsODN-UGT74AF3 (**a and b**) and in different genotypes of the tea plant (**c, d**). Tea leaves treated with sense oligonucleotides were used as a control (CK). Three antisense oligodeoxynucleotides were designated as 1, 2, and 3. ND not detected. Statistical significance was calculated with one-way ANOVA using SPSS 17.0 ($P < 0.05$)

(0.32 μ M) from strawberries²³, but significantly lower than that of UGT71k3a (900 μ M) and UGT71K3b (406 μ M) from strawberries²² and UGT85K14 (156 μ M) from a hybrid grapevine cultivar¹¹, suggesting that UGT74AF3a/b from tea plants showed a higher affinity for HDMF.

Although phylogenetic analysis can predict the function of UGTs, some enzymes of the same branch in the phylogenetic tree may have different functions (Fig. 7). In our study, UGT74AF3 preferentially glucosylated HDMF (Fig. 2), while UGT74F1, UGT74F2, and UGT74D1 from *Arabidopsis thaliana* glucosylated flavonols, salicylic acid, and auxin, respectively^{37–39}. In addition, a previous study indicated that UGT74E2 was involved in the modulation of the plant water-stress response⁴⁰. Identification of other HDMF glucosyltransferases in a new family will provide new insight into the substrate specificity of UGTs and the molecular evolution of HDMF glucosyltransferases.

Residue 456 determines sugar-donor preference of UGT74AF3

Diversity in nucleotide sequences of glycosyltransferases leads to diversity in their functions⁴¹. The structural

information of glycosyltransferases plays an important role in studying the glycosyltransferase structure or mechanism. Although 3D structures of UGTs have already been reported^{41–43}, the molecular mechanism of glycosyltransferase binding to substrates is still unknown. Previous studies indicated that the N-terminal amino acid residues of glycosyltransferase-encoded proteins are related to acceptor binding, while the C-terminal amino acid residues are closely related to that of donors^{22,41,44–47}. In this study, site-directed mutagenesis at a single position (A456V) led to an increase in HDMF glucosylation activity of UGT74AF3b (Fig. 4), which confirmed that amino acids close to the C-terminus affect the enzymatic activity, while mutations in the N-terminus mainly modify the substrate preference^{2,47}.

The comparison of two enzymes with almost identical primary sequences showed that the ability of the enzymes to differentiate between UDP-Glc and UDP-Gal as donor substrates was primarily due to a single residue, Glu303, in the C-terminus, which was not located in the active site⁴⁸. In this study, the A456V mutant significantly increased the activity toward UDP-Glc and UDP-Gal when compared with that of wild-type UGT74AF3b when HDMF, EHMF, and HMF were used as substrates (Fig. 4). The discovery

Fig. 7 Phylogenetic tree of UGT74AF3a, UGT74AF3b and selected plant glycosyltransferases. The phylogenetic tree was generated by the neighbor-joining method within the MEGA 6.0 program using 1,000 bootstrap replications. The following accession numbers were utilized: UGT85K14 (LC021362), UGT85K11 (ABB847092), UGT85K16 (LC312711), UGT76D1 (At2g26480), CsUGT85A53 (ALO19888.1), CsUGT78A14 (ALO1989.1), CsUGT84A22 (ALO19890), FaGT2 (AY663785), CsUGT73A20 (ALO19886.1), UGT94P1 (ABB847093), UGT71K3 (XP_004294260.1), and UGT71C3 (At1g07260). The bar indicates a 0.1 amino acid substitution/site

that residue 456 determines the sugar-donor preference of UGT74AF3 enhances our understanding of site-directed mutagenesis as an efficient method for improving and altering the sugar-donor preference of UGTs.

In planta function assessment of UGT74AF3

For a long time, because the transgenic system of tea plants has not been successfully established, the *in vivo* functional verification of tea plant genes has only been carried out in heterologous plants^{2,27}. In this study, the *in planta* function of UGT74AF3 was first investigated in tobacco, where overexpression of UGT74AF3a resulted in an increased HDMF glucoside content compared to that of the control (Fig. 5), suggesting that UGT74AF3 can glucosylate HDMF in tobacco.

Transient gene suppression using antisense oligonucleotides (AsODNs) is widely used in gene function

analyses, and the use of AsODNs has been extended to various plant species⁴⁹. A recent study showed that the functions of CsLIS/NES-1 and CsLIS/NES-2 in tea plants were successfully verified by AsODN. This method was further validated by treating tea seedlings with AsODN_PDS to suppress carotenoid production, suggesting that the AsODN method is effective in the tea plants⁵⁰. To determine the function of UGT74AF3 in the tea plant, the expression level of *UGT74AF3* was transiently suppressed in *C. sinensis* leaves by gene-specific AsODN suppression according to Zhao et al.³¹. The expression level of *UGT74AF3* in tea leaves treated with AsODN_UGT74AF3 was significantly reduced in accordance with the HDMF glucoside content in comparison with that of the controls (Fig. 6). Finally, *UGT74AF3* gene expression was shown to correlate well with the concentration of HDMF glucosides in leaves of different tea

Fig. 8 The proposed HDMF glucoside biosynthesis pathway in tea plants

plant cultivars (Fig. 6), suggesting that UGT74AF3 can catalyze HDMF glucosylation in tea plants (Fig. 8).

The identification of HDMF glucoside in tea plants and the discovery of a novel-specific UDP-glucose:HDMF glucosyltransferase provide not only the basis for the improvement of tea flavor, but also new insights for understanding the sugar-donor preference of UGTs and the molecular evolution of HDMF glucosyltransferases in plants.

Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (31961133030; 31870678), National Key Research and

Development Program of China (2018YFD1000601), Science Fund for Distinguished Young Scientists of Anhui Province (1908085J12).

Author contributions

C.S., W.S., and Y.C. conceptualized the initial study; Y.C., X.G., T.G., and N.Z. were involved in the experimental layout; Y.C., T.G., X.G., and N.Z. performed the lab experiments; C.S., W.S., X.C., and Y.C. drafted the initial article; all authors discussed the results, reviewed the article, and approved the final article.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-0248-x>).

Received: 28 July 2019 Revised: 28 November 2019 Accepted: 4 January 2020

Published online: 01 March 2020

References

- Arimura, G. I., Matsui, K. & Takabayashi, J. Chemical and molecular ecology of herbivore-induced plant volatiles: proximate factors and their ultimate functions. *Plant Cell Physiol.* **50**, 911–923 (2009).
- Jing, T. et al. Glucosylation of (Z)-3-hexenol informs interspecies interactions in plants: a case study in *Camellia sinensis*. *Plant, Cell Environ.* **42**, 1352–1367 (2019).
- Schuh, C. & Schieberle, P. Characterization of the key aroma compounds in the beverage prepared from Darjeeling black tea: quantitative differences between tea leaves and infusion. *J. Agric Food Chem.* **54.3**, 916–924 (2006).
- Ho, C.-T., Zheng, X. & Li, S. Tea aroma formation. *Food Sci. Hum. Wellness* **4**, 9–27 (2006).
- Larsen, M., Poll, L. & Olsen, C. E. Evaluation of the aroma composition of some strawberry (*Fragaria ananassa* Duch) cultivars by use of odour threshold values. *Z. Leb.* **195**, 536–539 (1992).
- Schieberle, P. & Hofmann, T. Evaluation of the character impact odorants in fresh strawberry juice by quantitative measurements and sensory studies on model mixtures. *J. Agric Food Chem.* **45**, 227–232 (1997).
- Schwab, W. Natural 4-hydroxy-2,5-dimethyl-3(2h)-furanone (Furaneol®). *Molecules* **18**, 6936–6951 (2013).
- Andreas, D., Martin, S. & Matthias, K. Nature's chemical signatures in human olfaction: a foodborne perspective for future biotechnology. *Angew. Chem. Int. Ed.* **53**, 7124–7143 (2014).
- Rodin, J. O., Himel, C. M., Silverstein, R. M., Leeper, R. W. & Gortner, W. A. Volatile flavor and aroma components of pineapple. I. Isolation and tentative identification of 2,5-dimethyl-4-hydroxy-3(2h)-furanone. *J. Food Sci.* **30**, 280–285 (1965).
- Ubeda, C. et al. Glycosidically bound aroma compounds and impact odorants of four strawberry varieties. *J. Agric Food Chem.* **60**, 6095–6102 (2012).
- Sasaki, K. et al. Molecular cloning and characterization of UDP-glucose: furaneol glucosyltransferase gene from grapevine cultivar Muscat Bailey A (*Vitis labrusca* × *V. vinifera*). *J. Exp. Bot.* **66**, 6167–6174 (2015).
- Buttery, R. G. et al. Analysis of furaneol in tomato using dynamic headspace sampling with sodium sulfate. *J. Agric Food Chem.* **49**, 4349–4351 (2001).
- Ito, Y. et al. Identification of potent odorants in Chinese jasmine green tea scented with flowers of *Jasminum sambac*. *J. Agric Food Chem.* **50.17**, 4878–4884 (2002).
- Sasaki, T. et al. Characterisation of odorants in roasted stem tea using gas chromatography-mass spectrometry and gas chromatography-olfactometry analysis. *Food Chem.* **220**, 177–183 (2017).
- Song, C. et al. Attractive but toxic emerging roles of glycosidically bound volatiles and glucosyltransferases involved in their formation. *Mol. Plant* **11**, 1225–1236 (2018).
- Bowles, D. et al. Glycosyltransferases: managers of small molecules. *Curr. Opin. Plant Biol.* **8**, 254–263 (2005).
- Wang, J. & Hou, B. Glycosyltransferases: key players involved in the modification of plant secondary metabolites. *Front Biol. China* **4**, 39–46 (2009).
- Perez, A. G. et al. Furanones in strawberries: evolution during ripening and postharvest shelf life. *J. Agric Food Chem.* **44**, 3620–3624 (1996).
- Wein, M. et al. Isolation, cloning and expression of a multifunctional O-methyltransferase capable of forming 2,5-dimethyl-4-methoxy-3(2H)-furanone, one of the key aroma compounds in strawberry fruits. *Plant J.* **31**, 755–765 (2002).
- Li, X.-X., Fukuhara, K. & Hayata, Y. Concentrations of character impact odorants in 'Toyonoka' strawberries quantified by standard addition method and PQ column extraction with GC-MS analysis. *J. Jpn. Soc. Hortic. Sci.* **78.2**, 200–205 (2009).
- Landmann, C., Fink, B. & Schwab, W. FaGT2: a multifunctional enzyme from strawberry (*Fragaria x ananassa*) fruits involved in the metabolism of natural and xenobiotic compounds. *Planta* **226**, 417–428 (2007).
- Song, C. et al. Glucosylation of 4-hydroxy-2,5-dimethyl-3(2H)-furanone, the key strawberry flavor compound in strawberry fruit. *Plant Physiol.* **171**, 139–151 (2016).
- Yamada, A. et al. A glucosyltransferase specific for 4-hydroxy-2,5-dimethyl-3(2H)-furanone in strawberry. *Biosci. Biotechnol. Biochem.* **53**, 1–8 (2019).
- Xia, E.-H. et al. The tea tree genome provides insights into tea flavor and independent evolution of caffeine biosynthesis. *Mol. Plant* **26**, 1–12 (2017).
- Wei, C. et al. Draft genome sequence of *Camellia sinensis* var. *sinensis* provides insights into the evolution of the tea genome and tea quality. *Proc. Natl Acad. Sci. USA* **115.18**, 622 (2018).
- Cui, L. et al. Identification of UDP-glycosyltransferases involved in the biosynthesis of astringent taste compounds in tea (*Camellia sinensis*). *J. Exp. Bot.* **67**, 2285–2297 (2016).
- Zhao, X. et al. Functional characterization of a new tea (*Camellia sinensis*) flavonoid glucosyltransferase. *J. Agric Food Chem.* **65**, 2074–2083 (2017).
- Ohgami, S. et al. Volatile glucosylation in tea plants: sequential glucosylations for the biosynthesis of aroma β -primeverosides are catalyzed by two *camellia sinensis* glucosyltransferases. *Plant Physiol.* **168**, 464–477 (2015).
- Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* **25**, 402–408 (2001).
- Sheikh, M. O. et al. Rapid screening of sugar-nucleotide donor specificities of putative glucosyltransferases. *Glycobiology* **27**, 206–212 (2017).
- Zhao, M. et al. Sesquiterpene glucosylation mediated by glucosyltransferase UGT91Q2 is involved in the modulation of cold stress tolerance in tea plants. *N. Phytologist*. <https://doi.org/10.1111/nph.16364> (2020).
- Ding, Y. & Lawrence, C. E. A statistical sampling algorithm for RNA secondary structure prediction. *Nucleic Acids Res.* **31.24**, 7280 (2003).
- Caputi, L., Lim, E. K. & Bowles, D. J. Discovery of new biocatalysts for the glucosylation of terpenoid scaffolds. *Chem. Eur. J.* **14**, 6656–6662 (2008).
- Mackenzie, P. I. et al. The UDP glucosyltransferase gene superfamily: recommended nomenclature update based on evolutionary divergence. *Pharmacogenetics* **7**, 255–269 (1997).
- Song, C. et al. Functional characterization and substrate promiscuity of UGT71 glucosyltransferases from strawberry (*Fragaria x ananassa*). *Plant Cell Physiol.* **56**, 2478–2493 (2015).
- Richman, A. et al. Functional genomics uncovers three glucosyltransferases involved in the synthesis of the major sweet glucosides of *Stevia rebaudiana*. *Plant J.* **41**, 56–67 (2005).
- Lim, E. K. et al. Arabidopsis glucosyltransferases as biocatalysts in fermentation for regioselective synthesis of diverse quercetin glucosides. *Biotechnol. Bioeng.* **87**, 623–631 (2004).
- Dean, J. V. & Delaney, S. P. Metabolism of salicylic acid in wild-type, *ugt74f1* and *ugt74f2* glucosyltransferase mutants of *Arabidopsis thaliana*. *Physiol. Plant* **132.4**, 417–425 (2008).
- Jin, S. H. et al. GT74D1 is a novel auxin glucosyltransferase from *Arabidopsis thaliana*. *PLoS ONE* **8**, 1628–1641 (2013).
- Tognetti, V. B. et al. Perturbation of indole-3-butyric acid homeostasis by the UDP-glucosyltransferase UGT74E2 modulates arabidopsis architecture and water stress tolerance. *Plant Cell* **22**, 2660–2679 (2010).
- Hu, Y. & Walker, S. Remarkable structural similarities between diverse glycosyltransferases. *Chem. Biol.* **9**, 1287–1296 (2002).
- Larson, E. T. et al. Structure of A197 from *Sulfolobus turreted* icosahedral virus: a crenarchaeal viral glucosyltransferase exhibiting the GT-A fold. *J. Virol.* **80.15**, 7636–7644 (2006).
- Daniel, B. et al. Positive cooperativity between acceptor and donor sites of the peptidoglycan glucosyltransferase. *Biochem. Pharmacol.* **93.2**, 141–150 (2015).
- Li, Y. et al. Phylogenetic analysis of the UDP-glycosyltransferase multigene family of *Arabidopsis thaliana*. *J. Biol. Chem.* **276**, 4338–4343 (2001).
- Ross, J., Li, Y., Lim, E. & Bowles, D. J. Higher plant glucosyltransferases. *Genome Biol.* **2.2**, 1–6 (2001).
- Paquette, S., Möller, B. L. & Bak, S. On the origin of family 1 plant glucosyltransferases. *Phytochemistry* **62**, 399–413 (2003).
- Bönisch, F. et al. A UDP-glucose:monoterpenol glucosyltransferase adds to the chemical diversity of the grapevine metabolome. *Plant Physiol.* **165**, 561–581 (2014).
- Patenaude, S. I. et al. The structural basis for specificity in human abo(h) blood group biosynthesis. *Nat. Struct. Biol.* **9**, 685–690 (2002).
- Dinc, E. et al. Synthetic antisense oligodeoxynucleotides to transiently suppress different nucleus- and chloroplast-encoded proteins of higher plant chloroplasts. *Plant Physiol.* **157.4**, 1628–1641 (2011).
- Liu, G. et al. Implementation of CSLIS/NES in Linalool Biosynthesis Involves Transcript Splicing Regulation in *Camellia Sinensis*. *Plant Cell Environ.* **41**, 176–186 (2017).