The optimal amount of salt intake

Kazufumi Nakamura¹ · Toru Miyoshi¹

Received: 8 January 2019 / Revised: 16 January 2019 / Accepted: 16 January 2019 / Published online: 14 February 2019
© The Japanese Society of Hypertension 2019

Sodium excretion is associated with increased blood pressure. The INTERSALT study revealed a significant linear relationship between the median 24-h urinary sodium excretion in 52 centers and the slope of systolic blood pressure with age in age- and sex-standardized analysis [1]. A pooled analysis of the results of four large studies, the PURE, EPIDREAM, ONTARGET, and TRANSCEND studies, also showed that the estimated 24-h urinary sodium excretion was associated with increased systolic and diastolic pressure [2].

However, the association between urinary sodium excretion and cardiovascular events remains controversial. The abovementioned pooled analysis showed that salt restriction reduced the risk of cardiovascular events and death in populations with hypertension and that an association of extensively low sodium intake with increased risk of cardiovascular events and death is observed in those with or without hypertension [2]. Meanwhile, the Trials of Hypertension Prevention (TOHP) trial showed an increased risk of mortality for individuals with high sodium intake and a direct relationship with total mortality, even at the lowest levels of sodium intake [3]. Therefore, the optimal level of sodium intake has been unclear.

Reduction of sodium intake is recommended in the guidelines of the European Society of Hypertension, the American Heart Association, the Japanese Society of Hypertension, and the World Health Organization [4–6]. Although the optimal level of sodium intake is controversial, a sodium reduction program is likely to be beneficial for prevention of cardiovascular diseases in communities in which people consume high amounts of salt [7]. The United Kingdom initiated a nationwide salt reduction program in 2003/2004, which was successful and resulted in a 15% reduction in population salt intake by 2011 (from 9.5 g/day in

2003 to 8.1 g/day in 2011). The Health Survey for England conducted from 2003 to 2011 showed decreases of 42 and 40% in the mortality rates of stroke and ischemic heart disease, respectively, along with decreases in systolic/diastolic blood pressure of 3.0/1.4 mm Hg. A recent community-level prospective epidemiological cohort study showed that sodium intake was associated with cardiovascular disease and stroke only in communities in which the mean sodium intake was more than 5 g/day [8]. Data from a prospective cohort study of a Chinese population also showed that high urinary sodium excretion was independently associated with an increased risk of cardiovascular disease [9]. Japan is a country in which people consume a large amount of salt. The National Health and Nutrition Survey in Japan, 2017 showed that the mean dietary salt intake in the period from 2007 to 2017 was 9.9 g/day (https://www.mhlw.go.jp/bunya/kenkou/kenkou_eiyouchousa.html). The ESPRIT study, a prospective observational study, showed that in Japan, high sodium excretion (≥ 4.0 g/day) was associated with composite cardiovascular events including hospitalization due to heart failure, acute coronary syndrome, cerebrovascular events, and cardiovascular deaths [10]. It seems difficult to determine the optimal amount of salt intake, but a strategy of sodium reduction would be appropriate in communities in which people prefer to season their food with salt.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Intersalt Cooperative Research Group. Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 h urinary sodium and potassium excretion. Intersalt Cooperative Research Group. *BMJ*. 1988;297:319–28.

✉ Kazufumi Nakamura
ichibun@cc.okayama-u.ac.jp

¹ Department of Cardiovascular Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, Japan

2. Mente A, O'Donnell M, Rangarajan S, Dagenais G, Lear S, McQueen M, et al. Associations of urinary sodium excretion with cardiovascular events in individuals with and without hypertension: a pooled analysis of data from four studies. *Lancet*. 2016;388:465–75.
3. Cook NR, Appel LJ, Whelton PK. Sodium intake and all-cause mortality over 20 years in the trials of hypertension prevention. *J Am Coll Cardiol*. 2016;68:1609–17.
4. Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M, et al. 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J*. 2013;34:2159–219.
5. Whelton PK, Carey RM, Aronow WS, Casey DE Jr., Collins KJ, Dennison Himmelfarb C, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018;71:1269–324.
6. WHO. Guideline: sodium intake for adults and children. Geneva: World Health Organization; 2012.
7. He FJ, Pombo-Rodrigues S, Macgregor GA. Salt reduction in England from 2003 to 2011: its relationship to blood pressure, stroke and ischaemic heart disease mortality. *BMJ Open*. 2014;4:e004549.
8. Mente A, O'Donnell M, Rangarajan S, McQueen M, Dagenais G, Wielgosz A, et al. Urinary sodium excretion, blood pressure, cardiovascular disease, and mortality: a community-level prospective epidemiological cohort study. *Lancet*. 2018;392:496–506.
9. Liu H, Gao X, Zhou L, Wu Y, Li Y, Mai J, et al. Urinary sodium excretion and risk of cardiovascular disease in the Chinese population: a prospective study. *Hypertens Res*. 2018;41:849–55.
10. Sadanaga T, Hirota S, Enomoto K, Kohsaka S, Tsujita K, Ito M, et al. Evaluation of sodium intake for the prediction of cardiovascular events in Japanese high-risk patients: the ESPRIT Study. *Hypertens Res*. 2019;42:233–40.