

DOI: 10.1038/s41467-018-06710-3

OPEN

Addendum: TGF- β signaling alters H4K20me3 status via miR-29 and contributes to cellular senescence and cardiac aging

Guoliang Lyu¹, Yiting Guan¹, Chao Zhang¹, Le Zong¹, Lei Sun¹, Xiaoke Huang¹, Li Huang¹, Lijun Zhang¹, Xiao-Li Tian², Zhongjun Zhou^{3,4} & Wei Tao¹

Addendum to: *Nature Communications* ; <https://doi.org/10.1038/s41467-018-04994-z>; published online 2 July 2018

Information relating to the in vivo treatments with A-196 and E-616452 was inadvertently omitted from the Methods section of this Article. In Fig. 6, Supplementary Fig. 7, as well as Supplementary Table 1, A-196 and E-616452 were used at a concentration of 25 mg/kg. A-196 and E-616452 were diluted in DMSO to a final concentration of 100 mM. Mice were treated with the inhibitors daily by oral gavage for 1 month.

Published online: 08 October 2018



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018

¹The MOE Key Laboratory of Cell Proliferation and Differentiation, School of Life Sciences, Peking University, Beijing 100871, China. ²Department of Human Population Genetics, Human Aging Research Institute and School of Life Science, Nanchang University, Nanchang 330031, China. ³School of Biomedical Sciences, LKS Faculty of Medicine, The University of Hong Kong, 21 Sassoon Road, Hong Kong, China. ⁴Shenzhen Institute of Innovation and Research, The University of Hong Kong, Nanshan, Shenzhen 518000, China. These authors contributed equally: Guoliang Lyu, Yiting Guan. Correspondence and requests for materials should be addressed to W.T. (email: weitao@pku.edu.cn)