AUTHOR CORRECTION

OPEN

Author Correction: Norgestimate inhibits staphylococcal biofilm formation and resensitizes methicillin-resistant *Staphylococcus aureus* to β -lactam antibiotics

Yutaka Yoshii^{1,2,3}, Ken-ichi Okuda^{1,2}, Satomi Yamada¹, Mari Nagakura¹, Shinya Sugimoto^{1,2}, Tetsuo Nagano⁴, Takayoshi Okabe⁴, Hirotatsu Kojima⁴, Takeo Iwamoto⁵, Kazuyoshi Kuwano³ and Yoshimitsu Mizunoe^{1,2}

npj Biofilms and Microbiomes (2017)3:30; doi:10.1038/s41522-017-0038-x

Correction to: npj Biofilms and Microbiomes (2017); doi:10.1038/s41522-017-0026-1; Published 21 July 2017

At proofing a reference was removed, but references in Fig. 2, Supplementary Information, Supplementary Table 1, and Supplementary Table 9 were not updated to reflect the change. These citations have now been corrected in the revised Figure and Supplementary Information files, in the HTML and PDF versions of this article. A correction has been published and is appended to both the HTML and PDF versions of this paper. The error has been fixed in the paper.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017

Fig. 2 The effects of NGM on the proteome of *S. aureus*. **a–c** Two-dimensional (2-D) electrophoresis in the presence or absence of NGM or 17DN (**a** DMSO control; **b** NGM; **c** 17DN). **d** Identification of proteins, whose expressions were decreased in the presence of NGM on 2-D gel

¹Department of Bacteriology, The Jikei University School of Medicine, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo 105-8461, Japan; ²Jikei Center for Biofilm Science and Technology, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo 105-8461, Japan; ³Division of Respiratory Diseases, Department of Internal Medicine, The Jikei University School of Medicine, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo 105-8461, Japan; ⁴Drug Discovery Initiative, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan and ⁵Division of Molecular Cell Biology, Core Research Facilities for Basic Science, The Jikei University School of Medicine, 3-25-8 Nishi-Shimbashi, Minato-ku, Tokyo 105-8461, Japan
Correspondence: Ken-ichi Okuda (okuda-k@jikei.ac.jp)
Yutaka Yoshii and Ken-ichi Okuda contributed equally to this work.