

PROTOCOL OPEN



Estimating mortality among inpatients with acute exacerbation of chronic obstructive pulmonary disease using registry data

Zhengcun Pei¹, Yixin Sun¹, Shengfeng Wang¹, Yahong Chen², Ting Yang^{3,4}, Kewu Huang^{5,6}, Yan Zhang⁷, Yin Huang⁸, Chen Wang^{4,9} and Siyan Zhan¹✉

The study aimed to investigate the demographic characteristics, clinical features, diagnoses, and treatments of hospitalized exacerbation COPD patients, as well as their disease prognoses and economic costs. The study planned to enroll 7600 hospitalized patients (aged ≥ 18 years with main diagnosis as AECOPD). Study patients were recruited since September 2017, followed up with a 3-year observing period. In the baseline visit, information on demographic characteristics, clinical features, diagnoses, and treatments were collected. In the following visits, treatments and examinations, recurrence of AECOPD, re-admission to hospital, complications, and mortality were recorded. Several validated questionnaires were applied at specific visits. This study included data from 1 September 2017 until 31 December 2022. The data would be used to estimate all-cause mortality during hospital stay, AECOPD recurrence within 1 month after discharge, all-cause and cause-specific mortality, frequency of AECOPD recurrence, lung function, life quality, healthcare costs in the study period, etc.

npj Primary Care Respiratory Medicine (2020)30:28; <https://doi.org/10.1038/s41533-020-0186-y>

INTRODUCTION

Chronic obstructive pulmonary disease (COPD), characterized by persistent airflow limitation, is a common preventable and treatable disease and is the leading cause of mortality and morbidity worldwide¹. More than three million deaths were caused by COPD each year^{2,3}. The prevalence of COPD in adults was estimated to be 4–10% worldwide^{4,5}, and 8.6% in China⁶, accounting for approximately 100 million COPD patients.

Acute exacerbation of COPD (AECOPD), defined by the Global Initiative for Chronic Obstructive Lung Disease, is an acute event characterized by a worsening of the patient's respiratory symptoms that is beyond normal day-to-day variations and leads to a change in medication^{7–9}; the disease burden and its exacerbations have been studied globally^{3,10,11}; exacerbation is a signal of disease progression and a major cause of patient hospitalization.

The severity and frequency of exacerbations strongly correlated with patient prognosis, especially mortality, whereas prevention, early detection, and prompt treatment would predict a better prognosis². However, as the mortality data with regard to AECOPD are limited for China, we speculated that there remains a gap between guidelines and clinical practices in China. This multi-center prospective patient registry study is launched to investigate the clinical features, treatments, and prognoses of AECOPD and meanwhile to build up a COPD management network.

The primary aim of this study was to assess the all-cause and cause-specific mortality among patients admitted to hospital for AECOPD and further to calculate re-admission rate caused again by AECOPD within 1 month after discharge. The secondary aims included analyses with regard to the 3-year follow-up after discharge, specifically all-cause and cause-specific mortality (at 1, 2, 3 years), recurrence of AECOPD (frequency of AECOPD

recurrence, the date of first recurrence after discharge), lung function, and quality of life assessment, as well as healthcare costs.

RESULTS

The primary aim of this study was to assess the in-hospital mortality of AECOPD patients, to describe all-cause and cause-specific mortality, and further to calculate re-admission rate caused again by AECOPD within 1 month after discharge from hospital. The secondary aims included analyses with regard to the 3-year follow-up after discharge from hospital, specifically all-cause and cause-specific mortality (at 1, 2, 3 years), recurrence of AECOPD (frequency of AECOPD recurrence, the date of first recurrence after discharge), lung function, and quality of life assessment, as well as healthcare costs.

The data obtained were planned to be used to describe the all-cause and cause-specific mortality of AECOPD patients, re-admission rates at the time points of 1 month to 3 years after discharge from hospital, as well as the recurrence of AECOPD, lung function decline, and quality of life.

Mortality rate, recurrence rate, and re-admission rate were planned to be reported pooled and stratified by sex, age, area, disease severity, etc. Their potential risk factors and confounders in consideration included COPD severity (lung function, AECOPD symptoms), medical history (COPD history, frequency of hospitalization, tobacco exposure, etc.), comorbidities (respiratory diseases, cardiovascular diseases, metabolic diseases, digestive diseases, etc.), treatment (bronchodilators, glucocorticoids, antibiotics, oxygen therapy, ventilatory support, etc.), questionnaire, and scales (COPD assessment test (CAT), modified British Medical Research Council (mMRC), St. George's respiratory questionnaire (SGRQ), hospital anxiety and depression scale (HADS)). The

¹Department of Epidemiology and Biostatistics, School of Public Health, Peking University, Beijing, China. ²Department of Pulmonary and Critical Care Medicine, Peking University Third Hospital, Beijing, China. ³Department of Pulmonary and Critical Care Medicine, Center of Respiratory Medicine, China-Japan Friendship Hospital, Beijing, China. ⁴National Clinical Research Center for Respiratory Diseases, Beijing, China. ⁵Department of Pulmonary and Critical Care Medicine, Beijing ChaoYang Hospital, Beijing, China. ⁶Department of Respiratory Medicine, Capital Medical University, Beijing, China. ⁷Best-Road Medi-Tech (Beijing) Ltd, Beijing, China. ⁸Beijing Natureself Technology Co. Ltd, Beijing, China. ⁹Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China. ✉email: siyan-zhan@bjmu.edu.cn

Table 1. Schematic diagram of the ACURE study.

Visit	Study period							
	Baseline survey	Follow-up survey						
	V0	V1	V2	V3	V4	V5	V6	V7
Time point	Admission–discharge	D30	M3	M6	M9	M12	M24	M36
Contact type	V	V	P	V	P	V	V	V
Eligibility screen	X							
Informed consent	X							
Demographics	X							
Medical history	X							
Personal history	X							
Treatment	X	X	X	X	X	X	X	X
Change of treatment for respiratory system		X	X	X	X	X	X	X
Physical examination	X							
Lung function test	X	X		X		X	X	X
Laboratory test	X	X		X		X	X	X
ECG	X							
Lung CT	X					X	X	X
UCG	X							
Pulmonary perfusion imaging	X							
Venous ultrasound of lower extremity	X							
PEACE questionnaire	X ^a	X	X	X	X	X	X	X
CAT questionnaire	X	X	X	X	X	X	X	X
mMRC questionnaire		X	X	X	X	X	X	X
SGRQ		X				X	X	X
HADS				X				
Prognosis (including death)	X	X		X		X	X	X
RICU/ICU admission	X							
Re-admission		X	X	X	X	X	X	X
AECOPD assessment		X	X	X	X	X	X	X
Lost to follow-up		X		X		X	X	X
Direct cost	X	X	X	X	X	X	X	X
Inhalation equipment				X				

ACURE estimating mortality among inpatients with Acute exacerbation of Chronic obstructive pulmonary disease using REgistry data, AECOPD acute exacerbation of chronic obstructive pulmonary disease, CAT COPD assessment test, CT computed tomography, D30 30 days after discharge, ECG electrocardiogram, HADS hospital anxiety and depression scale, M3, M6, M9, M12 3/6/9/12 months after discharge, mMRC modified British Medical Research Council, P phone call, PEACE questionnaire from the study by Zheng et al.¹³, RICU respiratory intensive care unit, SGRQ St. George's respiratory questionnaire, UCG ultrasound cardiogram, V visit, X information collected.

^aEveryday during hospitalization.

association between death event and its potential influential factors, the association between recurrence and readmission with their risk factors, the association between treatment and future outcomes, and similar associations of interest were planned to be assessed using multivariate linear/logistic regressions and Cox regression to assess time to death and its relevant factors. A mixed-effects model for longitudinal data analysis would be an optional choice to validate long-term effect of potential risk factors. In addition, comparative effectiveness research on different inpatient therapies (bronchodilators, glucocorticoids, antibiotics, oxygen therapy, ventilatory support) and prognoses (recovery, death, length of hospitalization), stable COPD management (pharmacological treatment, tobacco cessation, pulmonary rehabilitation, oxygen therapy), and disease progression (death, readmission, new comorbidities, lung function decline), as well as the impact of medication adherence on disease progression were planned to be compared after propensity score matching. Statistical significance were defined as the two-sided p value < 0.05 . All the

data were planned to be analyzed using the R software version 3.6, SAS V9.4 statistical package or STATA V15.0.

DISCUSSION

The ACURE (estimating mortality among inpatients with Acute exacerbation of Chronic obstructive pulmonary disease using REgistry data) study aimed to describe the overall clinical features and treatment procedures of AECOPD patients. The data obtained could be used to better understand the long-term outcome and risk factors of AECOPD and lung function declines.

As a pioneering study for proper measuring mortality of AECOPD patients all over the country, this study enabled our clinicians and researchers to address fundamental issues regarding the “real world” situation of AECOPD in China. It will also serve as a harmonized, evidence-based registry and platform for conducting future research, which will ultimately improve the management care provided to AECOPD patients.

Table 2. Steering committee of the ACURE study.

Center	Province/city	Principal investigator/coordinator
Peking University Third University	Beijing	Yahong Chen, MD
Chinese PLA General Hospital	Beijing	Junchang Cui, MD
Peking Union Medical College Hospital	Beijing	Jinglan Wang, MD
Beijing Hospital	Beijing	Chun Pu, MSc
The Second Hospital of Hebei Medical University	Hebei	Xixin Yan, MD
The First Affiliated Hospital of Chongqing Medical University	Chongqing	Shuliang Guo, MD
West China Hospital Sichuan University	Sichuan	He Yu, MSc
Changhai Hospital	Shanghai	Yuchao Dong, MD
The Second Affiliated Hospital of Chongqing Medical University	Chongqing	Daixin Wang, MD
The First Hospital of China Medical University	Liaoning	Hongwen Zhao, MD
General Hospital of East China	Jiangsu	Xiaoyong Xu, MD
China-Japan Friendship Hospital	Beijing	Ting Yang, MD

ACURE estimating mortality among inpatients with Acute exacerbation of Chronic obstructive pulmonary disease using REgistry data, PLA People's Liberation Army.

questionnaires [CAT questionnaire, PEACE questionnaire (consisting of eight questions assessing daily variance of COPD symptoms, i.e., dyspnea, purulent sputum, sputum volume, upper respiratory tract infection, fever, wheeze, cough, breath rate), mMRC questionnaire], medical examinations, laboratory tests, and treatments was recorded. Comorbidities including respiratory diseases (pneumonia, pulmonary embolism, interstitial lung disease, pulmonary arterial hypertension, lung cancer, asthma, respiratory failure), cardiovascular diseases (myocardial infarction, angina pectoris, hypertension, chronic heart failure, ventricular premature beat, right bundle branch block, atrial fibrillation), metabolic diseases (diabetes, osteoporosis), and digestive diseases (gastroesophageal reflux disease, peptic ulcer, cirrhosis), as well as malignancies other than lung cancer, peripheral arterial disease, venous thromboembolism, cerebrovascular disease, anxiety/depression, musculoskeletal dysfunction, chronic kidney disease, etc. were recorded. When a patient was discharged from hospital, information on diagnosis, CAT questionnaire, disease prognosis, intensive care unit stay (if any), and costs was collected. During each follow-up visit, information on the management; progress; recurrence; and prognosis of COPD, pharmacological, and non-pharmacological treatment was collected; meanwhile, scale and questionnaire surveys [CAT, mMRC, SGRQ, HADS] were conducted. An overview of the collected data is listed in Supplementary Table 1.

The primary aim of the study was to estimate the mortality rate during hospital stay; the sample size required is 6080 individuals, which was calculated based on an estimated mortality rate of 5% with an absolute precision of 2%, 95% confidence interval, and high design effect when recruiting patients from 40 hospitals, using the software CSurvey 2.0¹²; and finally 7600 individuals are required with an assumption of 20% drop-outs, as loss to follow-up is inevitable in such longitudinal design.

DATA AVAILABILITY

No datasets were generated or analyzed during the current study.

CODE AVAILABILITY

No codes were generated during the current study.

Received: 3 February 2020; Accepted: 7 May 2020;

Published online: 16 June 2020

REFERENCES

1. GBD 2015 Chronic Respiratory Disease Collaborators. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Respir. Med.* **5**, 691–706 (2017).
2. Fang, X., Wang, X. & Bai, C. COPD in China: the burden and importance of proper management. *Chest* **139**, 920–929 (2011).

3. Lozano, R. et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* **380**, 2095–2128 (2012).
4. Halbert, R. J., Isonaka, S., George, D. & Iqbal, A. Interpreting COPD prevalence estimates: what is the true burden of disease? *Chest* **123**, 1684–1692 (2003).
5. Buist, A. S. et al. International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet* **370**, 741–750 (2007).
6. Wang, C. et al. Prevalence and risk factors of chronic obstructive pulmonary disease in China (the China Pulmonary Health [CPH] study): a national cross-sectional study. *Lancet* **391**, 1706–1717 (2018).
7. Rodriguez-Roisin, R. Toward a consensus definition for COPD exacerbations. *Chest* **117**, 398S–401S (2000).
8. Celli, B. R. & Barnes, P. J. Exacerbations of chronic obstructive pulmonary disease. *Eur. Respir. J.* **29**, 1224–1238 (2007).
9. Burge, S. & Wedzicha, J. A. COPD exacerbations: definitions and classifications. *Eur. Respir. J. Suppl.* **41**, 46s–53s (2003).
10. Vogelmeier, C. F. et al. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease 2017 Report. GOLD Executive Summary. *Am. J. Respir. Crit. Care Med.* **195**, 557–582 (2017).
11. Wedzicha, J. A. & Seemungal, T. A. COPD exacerbations: defining their cause and prevention. *Lancet* **370**, 786–796 (2007).
12. Farid, M. N. & Frerichs, R. R. CSurvey 2.0. <http://www.ph.ucla.edu/epi/cs Survey.html> (2008).
13. Zheng, J.-P. et al. Effect of carbocysteine on acute exacerbation of chronic obstructive pulmonary disease (PEACE Study): a randomised placebo-controlled study. *Lancet* **371**, 2013–2018 (2008).

ACKNOWLEDGEMENTS

This study is funded by the National Key R&D Program of China (Project number: 2016YFC1304300).

AUTHOR CONTRIBUTIONS

The study concept and design was conceived by C.W. and S.Z. Analysis will be performed by Z.P., Y.S., and S.W. Z.P. prepared the first draft of the manuscript. Y.C., T.Y., and K.H. provided expert clinical advice, Y.Z. and Y.H. provided expert advice on research scheme and EDC system. All authors read and approved the final manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information is available for this paper at <https://doi.org/10.1038/s41533-020-0186-y>.

Correspondence and requests for materials should be addressed to S.Z.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give

appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020