ARTICLE OPEN

Differential effects of changes in cardiorespiratory fitness on worst- and best- school subjects

Toru Ishihara¹, Noriteru Morita²✉, Toshihiro Nakajima³, Koji Yamatsu⁴, Koichi Okita⁵, Masato Sagawa² and Keita Kamijo⁶✉

Accumulating evidence shows a beneficial association between physical fitness and school children's academic performance. However, several other studies have failed to demonstrate such an association. We reanalyzed data of a two-year longitudinal study of the association between changes in cardiorespiratory fitness and academic performance of school children by focusing on intra-individual variability in grade points as a possible source of this discrepancy. We analyzed data from 469 junior high school students to examine if improvements in cardiorespiratory fitness had a differential effect on an individual student's worst and best grade points. Results indicated that improvements in physical fitness were associated with an improvement in the worst grade points. On the contrary, we did not observe a similar longitudinal association with the best grade points. These findings suggest that improving cardiorespiratory fitness improves the worst grade points of an individual, selectively. We suggest that intra-individual variability in grade points might moderate the association between physical fitness and academic performance changes.

npj Science of Learning (2021)6:8; <https://doi.org/10.1038/s41539-021-00086-8>

INTRODUCTION

The epidemic of physical inactivity and the corresponding reduction in physical fitness during childhood has become a global public health problem^{1,2}. A large body of cross-sectional studies has consistently found a positive association between physical fitness, including several of its dimensions, such as cardiorespiratory fitness, muscular fitness, and motor fitness, and the academic performance of school children^{3–6}, measured by academic subjects' grade points (GP), or standardized academic test scores. These findings indicate the crucial role of regular physical activity and the associated improvements in physical fitness on the healthy development of the brain. Moreover, several longitudinal studies, including observational studies, randomized controlled trials (RCTs), and cluster RCTs have supported these cross-sectional findings by also demonstrating a positive causal link between improvements in physical fitness associated with regular physical activity and academic performance^{7–11}. However, several recent large-scale cluster RCTs have reported no effect of physical activity interventions on academic performance^{12–14}, casting doubt on the causality in this relationship. Multiple factors might cause the discrepant findings of longitudinal studies, including participant characteristics (e.g., sex) and the characteristics of the physical activity investigated (e.g., type and duration)¹⁵. We focused on intra-individual variability in GP for academic subjects, among other factors, to identify possible moderators of the association between physical fitness and academic performance.

Most cross-sectional studies have demonstrated a positive association between childhood fitness and academic performance. Nevertheless, this association might differ based on the subject. For example, several studies have reported a positive association between cardiorespiratory fitness and mathematics, but not with reading^{16,17}, suggesting that only certain subjects might benefit from the effects of physical fitness. Another study, however, has reported the opposite association, indicating that

cardiorespiratory fitness is significantly associated with reading, but not with mathematics¹⁸. Similarly, longitudinal studies have shown contradictory findings by indicating different changes for different academic subjects^{9,19}. We assumed that this discrepancy might be partly due to the baseline performance of a student. Ishihara, Drollette, Ludyga, Hillman, and Kamijo²⁰ reanalyzed three RCTs to examine whether the effects of physical activity interventions on executive function, which refers to prefrontal dependent goal-directed cognitive processes²¹, are moderated by the baseline performance. A large body of evidence has shown that executive function predicts both GP and standardized academic test scores²², indicating an association between physical fitness and academic performance. Ishihara et al.²⁰ indicated that improvements in executive function resulting from physical activity interventions were greater for individuals with lower baseline cognitive performance. Likewise, subgroup analysis of a large-scale cluster RCT¹³ that found no effects of physical activity interventions on academic performance, however, also indicated that children with the lowest tertile group for math performance at baseline improved their academic performance following the intervention, which was not the case for children in the higher tertile groups. Therefore, a low baseline performance might be associated with greater benefits from physical activity interventions. These findings suggest that inter-individual differences in baseline performance might moderate the effects of physical activity interventions on academic performance.

From the alternative perspective of baseline performance, intra-individual variability in GPs might potentially affect the longitudinal association between improved physical fitness and academic performance. Studies focusing on inter-individual differences in baseline performance^{13,20} suggest that an individual's worst school subjects are more sensitive to the beneficial effects of improved physical fitness than the best subjects. Moreover, the two studies cited above also indicated that physical

¹Graduate School of Human Development and Environment, Kobe University, Kobe, Hyogo, Japan. ²Department of Sports Cultural Studies, Hokkaido University of Education, Iwamizawa, Hokkaido, Japan. ³Department of Teachers Training, Hokkaido University of Education, Sapporo, Hokkaido, Japan. ⁴Faculty of Education, Saga University, Saga, Saga, Japan. ⁵Department of Sport Education, Hokusho University, Ebetsu, Hokkaido, Japan. ⁶University of Tsukuba, Tsukuba, Ibaraki, Japan. ✉email: morita.noriteru@i.hokkyodai.ac.jp; ktakamijo@gmail.com

activity interventions did not cause a deterioration of executive or academic performance even in individuals with high baseline performance, suggesting that increasing the time spent on physical activities does not detract from academic performance, which was also corroborated in a systematic review by Donnelly et al.²³. Therefore, findings at the intra-individual level suggest that improving physical fitness does not reduce the best grades of an individual student.

Recent studies in this field have employed rigorous designs controlling for potential confounders such as socioeconomic status (SES), which might be a possible source of inconsistencies in early studies²³. Nevertheless, most studies have failed to consider out-of-school learning activities, which is a critical factor affecting academic performance. Out-of-school learning activities, such as doing homework, attending cram schools, and private tutoring, are known to influence academic performance directly^{3,24}. Therefore, it is essential to examine whether the expected association between physical fitness and academic performance would be observed after controlling for out-of-school learning time.

The present study reanalyzed data from a two-year longitudinal study¹¹ to investigate the association between changes in physical fitness and academic performance of junior high school children by focusing on intra-individual variability in school grades. We specifically examined if improvements in cardiorespiratory fitness had differential effects on an individual's worst and best GPs. We focused on cardiorespiratory fitness, which several studies have indicated was positively associated with academic performance^{3–5,7,10,11}, although the exact changes in fitness components that most contribute to improving academic performance remain unclear. Moreover, we rigorously controlled for potentially

significant confounding variables, including out-of-school learning time and SES, when examining the association between cardiorespiratory fitness and academic performance. We predicted that physical fitness improvements would be associated with an improvement in the lowest GP, whereas it would have weak or no association with the highest GP.

RESULTS

The results of Pearson's correlation analyses are presented in Table 1. It can be seen that changes in cardiorespiratory fitness were not associated with changes in learning time. However, changes in learning time were positively associated with changes in the lowest and highest GPs, whereas changes in cardiorespiratory fitness were positively associated with only changes in the lowest GP. No association was found between changes in cardiorespiratory fitness and the highest GP. The associations between changes in cardiorespiratory fitness and changes in the lowest and highest GPs are illustrated in Fig. 1. Multiple regression analyses indicated that these associations between changes in cardiorespiratory fitness with changes in the lowest and highest GPs remained unchanged (Cohen's $f^2 = 0.02$, $1-\beta = 0.87$ and Cohen's $f^2 = 0.003$, $1-\beta = 0.22$, respectively) after controlling for the household income, maternal education, sex, change in BMI, and change in learning time (Table 2).

DISCUSSION

This study's main findings corroborated the study's hypothesis by indicating that improvements in physical fitness were associated

Table 1. Results of Pearson's correlation analyses.

	1	2	3	4	5	6	7
1. Change in cardiorespiratory fitness	–						
2. Change in lowest GP	0.15*	–					
3. Change in highest GP	0.09	0.35*	–				
4. Household income	0.12*	0.01	0.04	–			
5. Maternal education	0.12*	0.09	–0.06	0.21*	–		
6. Sex	0.35*	0.04	0.07	0.08	0.13*	–	
7. Change in BMI	–0.27*	0.04	–0.04	–0.01	–0.04	–0.26*	–
8. Change in learning time	0.05	0.17*	0.21*	0.09	0.04	–0.10*	0.05

GP grade point, BMI body mass index.

* $p < 0.05$.

Fig. 1 The associations between changes in cardiorespiratory fitness and changes in the lowest (left panel) and highest (right panel) GPs. Regression lines are shown with 95% confidence bands (shaded areas).

Table 2. Results of multiple regression analyses predicting changes in the lowest and highest grade points.

Variables	Change in lowest GP				Change in highest GP			
	β	SE	95% CI	<i>p</i>	β	SE	95% CI	<i>p</i>
Change in cardiorespiratory fitness	0.15	0.05	0.06 to 0.25	0.002	0.06	0.05	−0.04 to 0.16	0.25
	Adjusted $\Delta R^2 = 0.02$				Adjusted $\Delta R^2 = 0.003$			
Confounders								
Household income	−0.04	0.05	−0.14 to 0.07	0.48	0.01	0.05	−0.09 to 0.12	0.81
Maternal education	0.07	0.05	−0.03 to 0.17	0.17	−0.10	0.05	−0.20 to 0.008	0.07
Sex	0.01	0.05	−0.08 to 0.11	0.78	0.08	0.05	−0.02 to 0.18	0.10
Change in BMI	0.06	0.06	−0.05 to 0.18	0.26	−0.009	0.06	−0.12 to 0.10	0.88
Change in learning time	0.16	0.05	0.07 to 0.25	<0.001	0.22	0.05	0.13 to 0.31	<0.001

BMI body mass index, GP grade point, SE standard error.

with an improvement in the worst GPs, whereas a similar longitudinal association was not observed for the highest GP. These findings are consistent with the results of RCTs that focused on inter-individual differences in baseline performance and reported that lower baseline performance is associated with greater benefits from physical activity interventions^{13,20}. The present study extended this line of research by showing that the longitudinal association between improvements in physical fitness and academic performance is moderated by intra-individual variability in academic grades. Stated differently, these findings suggest that the beneficial association between physical fitness and the lowest GP could be observed even in students with high baseline performance.

Moreover, this study suggests that the null effect of physical activity interventions on academic performance observed in previous RCTs^{12–14} could be partially explained by the difficulty of academic tests. The beneficial effects of physical activity interventions on academic performance might not be detectable when the overall test scores of an individual are relatively high. Also, this study indicated that physical fitness changes were not statistically related to the highest GP, which supports the position of the American College of Sports Medicine that increasing physical activities does not negatively influence academic performance²³.

The present study provides novel findings regarding out-of-school learning time. Not surprisingly, increased out-of-school learning time increased both the lowest and highest GPs. More importantly, the longitudinal association between changes in cardiorespiratory fitness and the lowest GP was independent of out-of-school learning time, suggesting that the observed increase in the lowest GP might have resulted from changes in classroom behaviors. Several RCTs have shown that physical activity interventions leading to increased cardiorespiratory fitness improve prefrontal cortex dependent executive function^{25–28}, demonstrating a bidirectional relationship with academic performance²⁹. Furthermore, executive function is likely to have critical functions in sustained attention³⁰ and academic motivation³¹. Although speculative, improvements in fitness might improve students' classroom behaviors, including the ability to concentrate on learning, and the motivation for learning might collectively increase the lowest GP. In contrast, no association was observed in the highest GP, probably because each participant might have been more motivated for the best subject even at baseline. Several studies have examined the effects of physical activity on on-task classroom behaviors. These studies suggest that on-task classroom behaviors can be improved immediately after a single bout of physical activity (i.e., acute positive effects)³². However, Szabo-Reed et al.³³ also reported that a 3-year physical activity intervention did not change on-task classroom behaviors (i.e., no

long-term effects). To date, studies on the long-term effects of physical activity on classroom behaviors are scarce, and more research evaluating the effects of physical activity on classroom behaviors is required in the future to clarify how fitness improvements selectively influence an individual's worst academic subjects.

Specific limitations of this study should be noted. Firstly, the possible effects of catch-up growth cannot be ruled out because this study used an observational design. From the perspective of inter-individual differences in baseline performance, students with low baseline performance and relatively low physical fitness could experience greater improvements in academic performance and physical fitness at follow-up due to catch-up growth. However, the present study focused on intra-individual variability, which indicated beneficial associations between increased physical fitness and the lowest GP, even in students with a high baseline score. Therefore, we believe that the observed increase in the lowest GP is mainly attributed to improvements in physical fitness rather than catch-up growth. Secondly, this study used cardiorespiratory fitness as a proxy for engagement in regular physical activities. It is unclear whether physical fitness improvements caused by the effects of development, or by physical activity per se, are critical for improving academic performance. Furthermore, investigations assessing both physical fitness and physical activity levels are required to clearly distinguish the effects of regular physical activities from developmental effects. Thirdly, the present study only focused on cardiorespiratory fitness. A recent meta-analysis has suggested that the positive effects of physical activity interventions on executive function, which is pivotal for academic performance^{34,35}, are different for different types of physical activity, with coordinative training being more effective than cardiorespiratory and resistance training¹⁵. It is suggested that future research examine whether the longitudinal relationship between improvements in physical fitness and intra-individual variability of academic subjects differs according to the components of fitness. Fourthly, the GP system varies among countries. The present study only targeted Japanese junior high schools, in which the subject teacher evaluates the GP for each academic subject. The Japanese subject teachers base their assessments on a wide variety of perspectives, including class activities, reflection papers, and paper test scores. Therefore, we believe that the findings of this study can be generalized to other GP assessment systems to a certain extent. Fifthly, none of the participants were receiving special education services for cognitive or attentional disorders. Nevertheless, the lack of assessment and information regarding the participants' mental health, learning disabilities, or neurological disorders is another limitation of the present study. As a result, we could not determine whether neuropsychological variables cofounded the observed

Fig. 2 Flow diagram of participant recruitment and follow-up procedures.

longitudinal associations between changes in fitness and GPs or whether these findings could be generalized to children with psychopathologies. Finally, the possibility of selection bias in the sample cannot be ruled out because approximately one-fourth of students were not included in the study due to missing informed consent.

This two-year longitudinal study demonstrated that a positive association between improvements in physical fitness and academic performance was strongly observed for an individual's worst GPs. Furthermore, the findings of this study regarding the highest GPs support the new perspective that the increase in physical fitness associated with regular physical activity has no adverse effects on academic performance. In conclusion, this study highlights the importance of childhood fitness for healthy brain development and academic success.

METHOD

Participants

Figure 2 shows the flow diagram of the participant recruitment and follow-up processes. We requested 20 public junior high schools in Sapporo, which is the prefectural capital of Hokkaido in northern Japan, and cities near Sapporo, for permission to recruit seventh-grade students (i.e., 12–13 years old). Eleven schools declined to participate, and three schools refused to respond to the SES questionnaire. Two schools were excluded from the analysis due to the lack of follow-up. Only students whose parents or guardians returned the signed informed consent form were included in this study (consent rate = 74%). Finally, the data of participants that completed the two-year study period ($N = 469$, 214 girls and 255 boys) were analyzed. Table 3 shows the characteristics of the participants in this study. The study was approved by the institutional review board of the Hokkaido University of Education and the principals of participating schools.

Procedure and measures

The academic year runs from April to March in Japan. We collected data on the participants' academic performance at the end of the seventh and the ninth grades (i.e., two years apart). Participants' fitness and out-of-school learning time were assessed in October when they were in the seventh and the ninth grade. The participants' SES was assessed in October when they were in the seventh grade.

Academic performance. The GPs of five academic subjects (Japanese, mathematics, social studies, science, and English) in the national standard curriculum was obtained from schools and used to measure academic performance. Students earn 1–5 points for each subject, with high GPs indicating high academic performance. We analyzed an individual's lowest and highest GPs at the end of the seventh and ninth grades.

Table 3. Characteristics of study participants.

Variables	7th grade	9th grade
<i>N</i>	469	
Girls/boys	214/255	
Household income ^a	3 (1)	
Maternal education ^a	3 (1)	
BMI (kg/m ²)	19 (3)	20 (3)
Lowest GP	3 (1)	3 (1)
Highest GP	4 (1)	4 (1)
Cardiorespiratory fitness	5 (2)	6 (2)
Learning time ^{a, b}	5 (1)	5 (2)

Values are presented as *N* or mean (SD).

BMI body mass index, *GP* grade point.

^aThese measures were assessed by using a five-point scale: household income, 1 = <2 million yen, 2 = 2 to 4 million yen, 3 = 4 to 6 million yen, 4 = 6 to 8 million yen, 5 = >8 million yen; maternal education, 1 = junior high school, 2 = high school, 3 = vocational school, 4 = junior college, 5 = undergraduate studies; learning time, 1 = <30 min, 2 = 30 min to 1 h, 3 = 1 to 2 h, 4 = 2 to 3 h, 5 = >3 h.

^bThis score is presented as the sum of scores on weekdays and weekends.

Cardiorespiratory fitness. Participants performed a cardiorespiratory fitness test during a physical education class conducted in each school. This fitness test has been adopted by the Ministry of Education, Culture, Sports, Science, and Technology of Japan and annually conducted nationwide. In the test, each school can select either a 20-m shuttle run or an endurance run of 1500 m for boys and 1000 m for girls as a cardiorespiratory fitness test. The data obtained from the cardiorespiratory fitness test was converted into a score ranging from 1 to 10, based on age- and sex-specific normative data for Japanese people³⁶, such that high scores indicated high cardiorespiratory fitness.

Learning time. Out-of-school learning time was assessed by using a five-point, self-reported questionnaire inquiring about the participants' out-of-school learning time on weekdays and weekends (from "<30 min/day" to ">3 h/day"). The scores for weekdays and weekends were summed and analyzed. We could obtain learning time data of only 451 participants at baseline and 466 participants at follow-up because of missing values.

Body mass index. The data on body weight and height were obtained from schools. We obtained body mass index (BMI) data, calculated as body weight (kg)/height (m²), for 468 and 356 participants at baseline and follow-up, respectively, because the follow-up data of one school were missing.

Socioeconomic status. SES was assessed only at baseline by using a five-point questionnaire administered to participants' parents or guardians, which requested the participants' parents or guardians about their household income (from "<2 million yen" to ">8 million yen"), and maternal educational attainment (ranging from "complete junior high school" to "earned a bachelor's degree"). We obtained the SES data consisting of household income and maternal educational attainment only from 354 and 361 participants, respectively, due to non-respondents.

Statistical analysis

Pearson's correlation analyses were conducted to determine the association between changes in cardiorespiratory fitness and out-of-school learning time with the lowest and the highest GPs. Then, multiple regression analyses with full-information maximum likelihood estimation were conducted for predicting the lowest and highest GPs after controlling for household income, maternal education, sex, change in BMI, and change in learning time. All statistical analyses were conducted with $\alpha = 0.05$ using the R Studio software, version 1.1.463. As mentioned, this study was a reanalysis of our previous study¹¹. Therefore, post-hoc power analyses were conducted to determine statistical power using G*Power 3.1.2³⁷. The effect size was defined as a small, medium, and large when Cohen's $f^2 = 0.02$, 0.15, and 0.35, respectively³⁸.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this article.

DATA AVAILABILITY

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

CODE AVAILABILITY

No custom and original software were used in this study.

Received: 4 September 2020; Accepted: 5 February 2021;

Published online: 01 April 2021

REFERENCES

- Guthold, R., Stevens, G. A., Riley, L. M. & Bull, F. C. Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc. Health* **4**, 23–35 (2020).
- Tomkinson, G. R., Lang, J. J. & Tremblay, M. S. Temporal trends in the cardiorespiratory fitness of children and adolescents representing 19 high-income and upper middle-income countries between 1981 and 2014. *Br. J. Sports Med.* **53**, 478–486 (2019).
- Ishihara, T. et al. Direct and indirect relationships of physical fitness, weight status, and learning duration to academic performance in Japanese schoolchildren. *Eur. J. Sport Sci.* **18**, 286–294 (2018).
- Kao, S.-C., Westfall, D. R., Parks, A. C., Pontifex, M. B. & Hillman, C. H. Muscular and aerobic fitness, working memory, and academic achievement in children. *Med. Sci. Sports Exerc.* **49**, 500–508 (2017).
- Kyan, A., Takakura, M. & Miyagi, M. Mediating effect of aerobic fitness on the association between physical activity and academic achievement among adolescents: a cross-sectional study in Okinawa, Japan. *J. Sports Sci.* <https://doi.org/10.1080/02640414.2018.1554552> (2018).
- Geertsens, S. S. et al. Motor skills and exercise capacity are associated with objective measures of cognitive functions and academic performance in pre-adolescent children. *PLoS ONE* **11**, e0161960 (2016).
- Bartee, R. T., Heelan, K. A. & Dority, B. L. Longitudinal evaluation of aerobic fitness and academic achievement among schoolchildren. *J. Sch. Health* **88**, 644–650 (2018).
- Booth, J. N. et al. Associations between objectively measured physical activity and academic attainment in adolescents from a UK cohort. *Br. J. Sports Med.* **48**, 265–270 (2014).
- Davis, C. L. et al. Exercise improves executive function and achievement and alters brain activation in overweight children: a randomized, controlled trial. *Health Psychol.* **30**, 91–98 (2011).
- Hsieh, S.-S. et al. The relations between 3-year changes in physical fitness and academic performance in nationally representative sample of junior high school students. *Sci. Rep.* **8**, 15978 (2018).
- Ishihara, T. et al. Relationship of participation in specific sports to academic performance in adolescents: a 2-year longitudinal study. *Scand. J. Med. Sci. Sports* **30**, 1471–1482 (2020).
- Donnelly, J. E. et al. Physical activity and academic achievement across the curriculum: Results from a 3-year cluster-randomized trial. *Prev. Med.* **99**, 140–145 (2017).
- Resaland, G. K. et al. Effects of physical activity on schoolchildren's academic performance: The Active Smarter Kids (ASK) cluster-randomized controlled trial. *Prev. Med.* **91**, 322–328 (2016).
- Resaland, G. K. et al. Gender-specific effects of physical activity on children's academic performance: The Active Smarter Kids cluster randomized controlled trial. *Prev. Med.* **106**, 171–176 (2018).
- Ludyga, S., Gerber, M., Pühse, U., Looser, V. N. & Kamijo, K. Systematic review and meta-analysis investigating moderators of long-term effects of exercise on cognition in healthy individuals. *Nat. Human Behav.* <https://doi.org/10.1038/s41562-020-0851-8> (2020).
- de Greeff, J. W. et al. Physical fitness and academic performance in primary school children with and without a social disadvantage. *Health Educ. Res.* **29**, 853–860 (2014).
- Hansen, D. M., Hermann, S. D., Lambourne, K., Lee, J. & Donnelly, J. E. Linear/nonlinear relations of activity and fitness with children's academic achievement. *Med. Sci. Sports Exerc.* **46**, 2279–2285 (2014).
- Scudder, M. R. et al. The association between aerobic fitness and language processing in children: Implications for academic achievement. *Brain Cognition* **87**, 140–152 (2014).
- Reed, J. A. et al. Examining the impact of integrating physical activity on fluid intelligence and academic performance in an elementary school setting: a preliminary investigation. *J. Phys. Act. Health* **7**, 343–351 (2010).
- Ishihara, T., Drollette, E. S., Ludyga, S., Hillman, C. H. & Kamijo, K. Baseline cognitive performance moderates the effects of physical activity on executive functions in children. *J. Clin. Med.* **9**, 2071 (2020).
- Miyake, A. & Friedman, N. P. The nature and organization of individual differences in executive functions: four general conclusions. *Curr. Dir. Psychol. Sci.* **21**, 8–14 (2012).
- Duckworth, A. L., Taxer, J. L., Eskreis-Winkler, L., Galla, B. M. & Gross, J. J. Self-control and academic achievement. *Annu. Rev. Psychol.* **70**, 373–399 (2019).
- Donnelly, J. E. et al. Physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Med. Sci. Sports Exerc.* **48**, 1197–1222 (2016).
- Ishihara, T. et al. Modeling relationships of achievement motivation and physical fitness with academic performance in Japanese schoolchildren: moderation by gender. *Physiol. Behav.* **194**, 66–72 (2018).
- Hillman, C. H. et al. Effects of the FITKids randomized controlled trial on executive control and brain function. *Pediatrics* **134**, e1063–1071 (2014).
- Kamijo, K. et al. The effects of an afterschool physical activity program on working memory in preadolescent children: fitness and working memory in children. *Dev. Sci.* **14**, 1046–1058 (2011).
- Drollette, E. S. et al. Effects of the FITKids physical activity randomized controlled trial on conflict monitoring in youth. *Psychophysiology* **55**, e13017 (2018).
- Ludyga, S., Gerber, M., Herrmann, C., Brand, S. & Pühse, U. Chronic effects of exercise implemented during school-break time on neurophysiological indices of inhibitory control in adolescents. *Trends Neurosci. Educ.* **10**, 1–7 (2018).
- Peng, P. & Kievit, R. A. The development of academic achievement and cognitive abilities: a bidirectional perspective. *Child Dev. Perspect.* **14**, 15–20 (2020).
- Thomson, D. R., Besner, D. & Smilek, D. A resource-control account of sustained attention: evidence from mind-wandering and vigilance paradigms. *Perspect. Psychol. Sci.* <https://doi.org/10.1177/1745691614556681> (2015).
- Eisenberg, N., Duckworth, A. L., Spinrad, T. L. & Valiente, C. Conscientiousness: origins in childhood? *Dev. Psychol.* **50**, 1331–1349 (2014).
- Mahar, M. T. et al. Effects of a classroom-based program on physical activity and on-task behavior. *Med. Sci. Sports Exerc.* **38**, 2086–2094 (2006).
- Szabo-Reed, A. N. et al. Impact of three years of classroom physical activity bouts on time-on-task behavior. *Med. Sci. Sports Exerc.* **49**, 2343–2350 (2017).
- Best, J. R., Miller, P. H. & Naglieri, J. A. Relations between executive function and academic achievement from ages 5 to 17 in a large, representative national sample. *Learn. Individ. Differences* **21**, 327–336 (2011).
- Duckworth, A. & Gross, J. J. Self-control and grit: related but separable determinants of success. *Curr. Dir. Psychol. Sci.* **23**, 319–325 (2014).
- Ministry of Education, Culture, Sports, Science and Technology of Japan. Standard operating procedure for physical fitness and exercise performance tests. https://www.mext.go.jp/a_menu/sports/stamina/03040901.htm.

37. Faul, F., Erdfelder, E., Buchner, A. & Lang, A.-G. Statistical power analyses using G*Power 3.1: tests for correlation and regression analyses. *Behav. Res. Methods* **41**, 1149–1160 (2009).
38. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences*. (Academic Press, 2013).

ACKNOWLEDGEMENTS

This study was supported by the Japan Society for the Promotion of Science (JSPS) KAKENHI Grant Numbers JP24650379, JP23500737, JP26282188, JP15K12723, and JP17H06479 from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

AUTHOR CONTRIBUTIONS

T.I. and K.K. developed the study concept. All authors contributed to the study design. T.I. performed the data analysis and interpretation under the supervision of K.K. T.I. and K.K. drafted the manuscript, and T.N., K.Y., K.O., M.S., and N.M. provided critical revisions and important intellectual content. All authors approved the final version of the manuscript for submission.

COMPETING INTERESTS

The authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41539-021-00086-8>.

Correspondence and requests for materials should be addressed to Noriteru Morita or Keita Kamijo.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021, corrected publication 2021