

REVIEW ARTICLE OPEN



Progress towards a glycoconjugate vaccine against Group A Streptococcus

Keira Burns^{1,2}, Helge C. Dorfmüller³, Brendan W. Wren^{1,2}, Fatme Mawas¹ and Helen A. Shaw¹✉

The Group A Carbohydrate (GAC) is a defining feature of Group A Streptococcus (Strep A) or *Streptococcus pyogenes*. It is a conserved and simple polysaccharide, comprising a rhamnose backbone and GlcNAc side chains, further decorated with glycerol phosphate on approximately 40% GlcNAc residues. Its conservation, surface exposure and antigenicity have made it an interesting focus on Strep A vaccine design. Glycoconjugates containing this conserved carbohydrate should be a key approach towards the successful mission to build a universal Strep A vaccine candidate. In this review, a brief introduction to GAC, the main carbohydrate component of Strep A bacteria, and a variety of published carrier proteins and conjugation technologies are discussed. Components and technologies should be chosen carefully for building affordable Strep A vaccine candidates, particularly for low- and middle-income countries (LMICs). Towards this, novel technologies are discussed, such as the prospective use of bioconjugation with PglB for rhamnose polymer conjugation and generalised modules for membrane antigens (GMMA), particularly as low-cost solutions to vaccine production. Rational design of “double-hit” conjugates encompassing species specific glycan and protein components would be beneficial and production of a conserved vaccine to target Strep A colonisation without invoking an autoimmune response would be ideal.

npj Vaccines (2023)8:48; <https://doi.org/10.1038/s41541-023-00639-5>

INTRODUCTION

Vaccination is considered one of the most successful health interventions known to man¹. Antibiotics and alternative therapeutics such as intravenous immunoglobulins are not suitable to control *Streptococcus pyogenes* or Group A Streptococcus (Strep A) infections at the population level or stop transmission within communities². The emergence of antimicrobial resistance (AMR), along with the persistence of penicillin sensitivity, has led the WHO and CDC to both highlight vaccines as urgent safeguards against AMR^{3,4}. Safe and efficacious Strep A specific vaccines are required for better control of Strep A related morbidity and mortality. Strep A is a major global pathogen, with disease manifestations ranging from Strep throat pharyngitis and impetigo to scarlet fever and invasive diseases such as toxic shock syndrome and necrotising fasciitis. These infections are often fast progressing and highly contagious, further emphasising the need for a vaccine. Invasive infections have high mortality and morbidity, and secondary diseases from autoimmune sequelae such as rheumatic heart disease (RHD) result in significant disease adjusted life years (DALYs), particularly in Low- and Middle-Income Countries (LMICs)⁵. In recent years there has further been a re-emergence of scarlet fever particularly noted in the UK and Europe, with particular concern in the current season with rising rates of sepsis associated with these infections^{6,7}. There is therefore an urgent need globally to develop a vaccine against Strep A. With humans as the sole natural host there is also potential to eradicate the pathogen if transmission can be blocked.

Vaccine development against any pathogen is a complex and lengthy process, however Strep A vaccine development has had arguably a more complicated history with many challenges and hurdles. Strep A vaccine development has an official impeded status recognised by the WHO, due to many factors loosely

relating to bacterial and host amongst others. First and foremost, Strep A is a complex pathogen with high antigen genomic heterogeneity due to recombination events, possessing different virulence factor expression profiles between strains, as well as complicated diverse global epidemiology of circulating Strep A serotypes. Strep A serotypes are based on variations in the major virulence factor, the M protein, with numerous variant serotypes globally and no consistent serotype observed in geographic regions or correlation with disease manifestations. At the genomic level there are even greater variations in the M-type, or *emm* type, with >200 variants. A recent genomic study showed that no single protein investigated during vaccine development has been 100% conserved between all the analysed Strep A isolates⁸. This has led to difficulty targeting a particular protein to cover all serotypes causing infections globally as a universal vaccine candidate. Gene exchange and single nucleotide polymorphisms (SNPs) causing protein sequence variation within vaccine candidates, such as within the most progressed NTD (N-terminal domain) M protein-based vaccines leads to the requirement of multicomponent inclusion to obtain acceptable levels of cross protection. Additionally broad spectrum of disease makes finding an effective long-term vaccine strategy challenging. Disproportionate Strep A disease burden and diverse serotype prevalence within LMICs, related to vaccine efficacy and protective coverage predication complicates testing when deciding on vaccine clinical endpoints generally and within these settings.

Strep A can be distinguished from other beta-haemolytic streptococci species by Lancefield serotyping. This technique is based on the identification of type-specific surface exposed carbohydrates that bind to specific antibodies⁹, with *S. pyogenes* being named Strep A due to the presence of the conserved group A carbohydrate (GAC) on its surface.

¹Vaccine Division, Scientific Research & Innovation Group, MHRA, Potters Bar, UK. ²Department of Infection Biology, London School of Hygiene and Tropical Medicine, London, UK. ³Division of Molecular Microbiology, School of Life Sciences, Dow Street, Dundee, UK. ✉email: alexandra.shaw@nibsc.org

Table 1. Summary of studies progressing vaccines containing Group A Carbohydrate (GAC) as well as derivative GAC^{PR} (polyrhymose) and synthetic polymers as stated.

Saccharide	Carrier protein/ nanotechnology	PS activation conjugation resulting structure	Animal model	Immune response	Protection	Autoimmune	Refs.
WT GAC	TT	Periodate oxidation Chemical coupling Selective sunlike	Mouse	IgG response	60–85% Passive (Rb sera) protection in mouse IP challenge 70–80% survival in mouse IP challenge Reduced nasal colonisation	Indirect immunofluorescence – negative brain, heart, kidney	31
Synthetic WT GAC (6mer and 12mer)	CRM ₁₉₇	Disuccinimidyl adipate activation (6/12mer)/periodate oxidation (GAC)	Mouse	IgG response (12mer and WT GAC high)	40% OPA killing (Rb sera) 30–60% survival in mouse IP challenge	NT	14
WT GAC	CRM ₁₉₇	Chemical coupling Selective sunlike		Antibody binding to GAS			
Synthetic GAC (6mer)	TT	Cysteamine activation Chemical coupling Selective sunlike	Mouse	IgG and IgM response	NT	NT	63
Synthetic GAC (3mer, 6mer, 9mer)	C5a peptidase	Hydrogenation and acylation/ DSG activated esters Chemical coupling Selective sunlike	Mouse	IgG response Antibody binding to GAS	30–60% OPA killing (6/9mer) 50–80% survival in mouse IP challenge	NT	65,66
GAC ^{PR}	Recombinant pneumococcal protein SP_0435	Chemical biotinylation Affinity interaction Selective	Rabbit Passive mouse	Antibody binding to GAS	Whole blood and OPA killing Survival from passive immunisation (Rb) in mouse IP challenge	Reduced ARF GicNac Mab reactivity Negative ELISA to human cardiac tissue lysate	10
GAC ^{PR}	Arginine Deiminase Protein	CDAP Chemical coupling Random	Mouse	IgG response Antibody binding to GAS	60% killing indirect bactericidal assay Clearance of GAS in mouse skin challenge No protection in mouse SC challenge	NT	56
GAC	CRM ₁₉₇ SpyCEP SpyAD SLO	Periodate oxidation Chemical coupling Selective sunlike / random mesh	Mouse	IgG response (GAC) Significant reduction in anti-protein IgG Antibody binding to GAS No neutralisation of specific protein activity	NT	NT	57
GAC ^{PR}	SpyAD Non-carrier SLO and C5a peptidase	CDAP and dibenzocyclooctyne derivatization Site-direct CLICK chemistry Selective sunlike	Rabbit (sera) Mouse (challenge)	Protein IgG titres Antibody binding to GAS SLO neutralisation	OPA killing ~50% survival from passive immunisation in mouse IP challenge 100% survival from multicomponent immunisation in mouse IP challenge	Negative for antibody cross- reactivity with human heart lysates by western blot	35
GAC ^{PR}	SLO	CDAP and dibenzocyclooctyne derivatization Site-direct CLICK chemistry Selective sunlike	Mouse	IgG response	90% survival from SLO-GAC ^{PR} immunisation in mouse IP challenge	NT	62

Table 1 continued

Saccharide	Carrier protein/ nanotechnology	PS activation conjugation resulting structure	Animal model	Immune response	Protection	Autoimmune	Refs.
Synthetic GAC ^{PR}	Gold nanoparticles	Thiolation activation Chemical coupling Selective sunlike (star/sphere)	N/A	Competitive ELISA with anti-Rha polyclonal IgG	NT	NT	102
Synthetic tri- rhamnose	Ac-PADRE-lipid core	-	Mouse	IgG response	50–100% OPA killing	NT	36
WT GAC	GMMMA	Adipic acid dihydrazide activation Chemical conjugation <i>Salmonella</i> Typhimurium GMMMA	Mouse	IgG response	NT	NT	103
Recombinant GAC ^{PR}	OMV	Biological anchoring to KDO Recombinant expression <i>E. coli</i> OMV	Mouse	IgG response Antibody binding to GAS	NT	NT	96

The group A carbohydrate (GAC)

Group A carbohydrate (GAC) is conserved across all Strep A strain cell surfaces^{8,10} and shown to be a key survival determinant¹⁰. GAC is abundant making up 40–60% of the total cell wall mass¹¹, functioning to provide structural support as an environmental barrier, maintain cell morphology and enable cellular division¹². In addition, GAC is also a major virulence determinant, providing resistance to zinc toxicity and resistance to neutrophil mediated killing^{10,13,14}.

GAC polymers have an average molecular mass of 8.9 ± 1.0 kDa, corresponding to 18 repeating units¹⁴, though different purification methods result in varying average sizes¹⁵. GAC is made up of a linear polyrhamnose backbone with alternating GlcNAc side-chains, with a trisaccharide repeating unit of $[3\alpha\text{-L-Rhap}(1\rightarrow2)[\beta\text{-D-GlcpNAc}(1\rightarrow3)]\alpha\text{-L-Rhap}(1\rightarrow3)]_n$ (Fig. 1)^{14,16}. GlcNAc is attached to the 2-hydroxyl linked rhamnose residue¹⁷ extending out to the periphery from the rhamnose helix¹⁸. Approximately 25–30% of the GlcNAc residues on GAC polymers contain glycerol phosphate (GroP) modifications, specifically on the C6-hydroxyl group. This polysaccharide modification has remained elusive for some time and the specific function has not been fully characterised. It may, however, contribute to the immune evasion activity through resistance to zinc toxicity¹⁹. A Strep A mutant devoid of GlcNAc, and therefore the GroP modification, was more susceptible to killing by human whole blood, as well as in the presence of purified human neutrophils, shown mechanistically to be due to greater binding of cationic human defence peptides, cathelicidin, LL-37 to the mutant polysaccharide. This loss of GlcNAc also seems to reduce the binding of cationic bactericidal enzyme human Group IIA secreted phospholipase A2 to mutant Strep A cell surfaces^{19,20}.

Branched polysaccharides composed of rhamnose and GlcNAc sidechains have been shown to be immunogenic, evidenced by rabbit and human antiserum^{21,22}, computer simulations^{18,23,24} and by NMR²⁵ techniques to demonstrate mAb and GAC interaction. The GlcNAc sidechain is predicted to play a role in human specific pathogenicity or immune evasion strategies. However, a consideration with using GAC as a vaccine component is the potential generation of autoimmune antibodies due to the presence of GlcNAc in many human glycan structures and similarities between GlcNAc and components of the extracellular matrix (ECM) in humans^{26,27}. Clinically this has been observed with acute rheumatic fever (ARF) patients having 2- to 3- fold higher anti-GAC antibody titres at the point of infection compared to patients with pharyngitis²⁸, as well as longer lived anti-GAC antibody populations in patients which have rheumatic heart disease (RHD)²⁹. Isolated monoclonal antibodies from ARF patients appear to only recognise wildtype GAC, not cross reacting with mutated GAC which does not contain GlcNAc sidechains¹⁰, demonstrating the antibody specificity to GlcNAc epitopes. The role of GlcNAc is, however, controversial with the majority of autoimmune responses attributed instead to the M-protein in the manifestation of ARF and RHD³⁰.

Antibody binding experiments have shown that GAC polymers are mainly localised to the outer surface of the cell wall¹¹. This surface localisation, conservation between strains and protective properties makes GAC and derivatives of interest as a polysaccharide component of a glycoconjugate vaccine.

GAC immunogenicity and considerations for carrier proteins

Immunological studies have revealed that GAC is accessible to antibody binding^{14,31,32}, with affinity-purified anti-GAC antibodies showing opsonic properties specifically against M3, M6, M14 and M28 serotypes³². Host infection with Strep A induces circulating GAC specific antibodies, which may be slow to generate initially, but are thought to gradually increase with age, peaking in adolescents³³. Some studies suggest the GlcNAc sidechain portion




Fig. 1 Structures of the GAC trisaccharide repeating unit. GAC is composed of a polyrhamnose backbone and GlcNAc side chains, decorated with glycerol phosphate. The wildtype repeating unit is $[3]\alpha\text{-L-Rhap}(1 \rightarrow 2)[\beta\text{-D-GlcpNAc}(1 \rightarrow 3)]\alpha\text{-L-Rhap}$ for 70% of the repeating units (i), and 30% modified on GlcNAc side chains with glycerol phosphate (ii) $[3]\text{-}\alpha\text{-L-Rhap}\text{-}(1 \rightarrow 2)[\beta\text{-D-GlcpNAc6P(S)Gro}\text{-}(1 \rightarrow 3)]\text{-}\alpha\text{-L-Rhap}\text{-}(1 \rightarrow 3)$. $\Delta gacI$ mutants and certain GAS strains passaged in mice have the repeating structure $[3]\text{-}\alpha\text{-L-Rhap}\text{-}(1 \rightarrow 2)\alpha\text{-L-Rhap}\text{-}(1 \rightarrow 3)\text{-}\alpha$ deficient in GlcNAc sidechains (iii).

of the polymer³², particularly the branch points within the trisaccharide repeat structure, to be important in recognition and generation of opsonophagocytic IgG antibodies^{22,32,34}. However, more recent data has shown antibodies also recognise rhamnose epitopes within the polymer backbone^{10,35}. The entire trisaccharide of Rha-Rha-GlcNAc has been demonstrated to bind to monoclonal antibodies by several techniques including 1D-TOCSY NMR²⁵. Further, that the branch structure itself is important, but GlcNAc can be substituted with rhamnose for the same response³⁶. Titres of anti-GAC antibodies are thought to correlate with reduced Strep A infection incidence in adolescents compared to young children, suggesting that anti-GAC antibodies may be important in long lasting immunity to Strep A infection. This justifies the argument that carbohydrate-based antibodies are a suitable approach towards protection from disease. This concept was highlighted in a study of Mexican children which showed high titres of GAC specific IgG antibodies correlated with protection against throat carriage³¹.

Polysaccharide antigens alone are T cell independent antigens, containing a repetitive structure which can be recognised and cross-linked by B cell receptors (BCR) on B cells, but provide no T cell epitopes. This leads to B cell differentiation into plasma cells which secrete antibodies directed against the polysaccharide epitopes but with reduced immunological memory particularly in infants^{37–40}. Conjugation of polysaccharide to a carrier protein enables the protein moiety to provide such T cell epitopes through presentation on MHC Class II complexes to generate CD4⁺ T cell help⁴¹. This strengthens and improves immune response longevity by increasing polysaccharide specific antibody levels, affinity maturation and proliferation of polysaccharide-specific B cells from memory pools, resulting in IgM to IgG isotype switching leading to higher antibody avidity⁴².

Protein carriers selected for inclusion in glycoconjugate vaccines are ideally themselves immunogenic. Suitable protein carriers must therefore enable induction of effective anti-polysaccharide immune responses and be compatible with conjugation techniques, being safe and produced at high yields and low costs^{43–45}. Traditionally, carrier proteins included in currently licenced glycoconjugate vaccines were heterologous to the organism against which you wish to vaccinate^{44,46}. There are currently five carrier proteins in licenced glycoconjugate vaccines; Tetanus Toxoid (TT), Diphtheria Toxoid, CRM₁₉₇ (a non-toxic mutant of diphtheria toxin), recombinant *E. coli* produced *Haemophilus influenzae* protein D and outer membrane protein complex of serogroup B meningococcus^{44,46}. These traditional carriers have been successfully included in licenced vaccines due to their compatibility with well-characterised conjugation chemistries, and their ability to induce effective long lasting anti-

polysaccharide immune responses⁴⁵. Recently, however there has been a drive for new carriers to be investigated⁴⁴, as data suggests that repeated immunisation with the same classical carrier for different glycoconjugate vaccines in some cases dampens immunological potency and efficacy^{47–50}. This is mediated by inhibition of polysaccharide antibody responses by carrier-specific B cells and suppressor T cells due to pre-existing carrier protein immunity referred to as carrier-induced epitopic suppression⁴⁹.

Currently, the potential benefit from protective antibodies directed against the protein component has not been fully exploited. In addition to avoiding carrier-specific epitopic suppression, vaccines containing both pathogen specific polysaccharide and protein antigens provides a “double-hit” approach^{45,51,52}, which may achieve broader immunity⁴⁴. There have been several studies on conserved Strep A protein antigens as vaccine candidates, often included in multicomponent vaccines and these have potential to be utilised as glycoconjugate carrier proteins^{35,53–58}. Rational protein antigen design to improve pathogen specific immune responses, in addition to providing effective T helper cell function to improve polysaccharide responses, is a key consideration for a Strep A glycoconjugate vaccine.

Chemical conjugation for glycoconjugates

Glycoconjugate vaccines can be synthesised through several conjugation methodologies, and the method choice is also an important consideration. This has traditionally been through chemical conjugation, which requires extraction and purification of polysaccharide directly from the organism’s cell wall or capsule against which you wish to protect. It also requires purification of recombinantly expressed protein carriers before covalent attachment of the two components can occur (Fig. 2a–c). Conjugation chemistries can utilise naturally present reactive groups or alternatively add in either cross-linking reagents to artificially introduce compatible reactive groups or chemically modify components through activation for attachment. Both these modifying reactions can improve conjugation efficiency but can also damage conformational epitopes in the process. Chemical modification can be through reducing end selective activation, or random multiple activation along polysaccharide chains⁵⁹ (Fig. 2d). The choice of chemical conjugation approach is often governed by structure, size, and composition of vaccine components⁶⁰. For example, larger polysaccharides are usually randomly activated, and smaller polysaccharides activated at the reducing end to preserve protective epitopes⁵⁹. Some selectivity can be achieved by modulating component stoichiometry⁶¹, targeting or introducing specific non-natural amino acids on the protein facilitating site directed attachment⁶². Such selective or milder




Fig. 2 Simplified schematic representation of chemical conjugation. Polysaccharide (PS) extracted from the native organism, e.g., GAS from Strep A cells (a), and recombinant protein carriers expressed and purified from *E. coli* cells (b). Extracted PS and recombinant protein carriers containing compatible reactive groups are conjugated together using a compatible cross-linking chemical reagent (c) yielding heterogeneous glycoconjugate vaccine candidates depending on the selected approach (d). Schematic shows different approaches yielding different glycoconjugate species, with reducing end chemistries leading to terminal single ended glycoconjugate products (sun-like structures), and random activation chemistries yielding cross-linked mesh-like structures of higher molecular weights with several attached protein-polysaccharide molecules.

approaches can be important for carrier proteins with a dual purpose to maintain protective B cell epitopes.

Progress towards a GAC glycoconjugate vaccine

There have been several studies investigating GAC or components of GAC as vaccine candidates. In this section we briefly summarise key studies and advancements from selected studies, while a list of all published studies on this topic is provided (Table 1). Early studies focussed on GAC immunogenicity when conjugated to classical carrier proteins to determine the anti-polysaccharide response and the immunological epitopes for protection. These included synthetic polymers, native GAC and GAC mutated to contain no GlcNAc sidechains (GAC^{PR}). The move towards modified GAC aimed to mitigate the potential risks of generating an autoimmune response^{26,27}. There has been a renewed interest in GAC as a vaccine component, with studies in the last five years building on incorporation of GAC into multicomponent formulations or “double-hit” glycoconjugates. To date, these studies have all made use of chemical methods for conjugation (Table 1).

An early study by Sabharwal et al., utilised native GAC conjugated to TT and showed protection of mice from intranasal colonisation and intraperitoneal lethal challenge with two different M Strep A serotypes³¹. This study was built on by

Kabanova et al. where synthetic polymers of varying lengths were conjugated to CRM₁₉₇ to investigate size dependent immunogenicity¹⁴. These were also able to generate protective antibodies and demonstrated that a hexasaccharide (composed of four rhamnose sugars, two of which were decorated with a GlcNAc sidechain) was the minimal epitope able to invoke a robust immune response. This reiterated earlier observations that a hexasaccharide was key to the natural immune response to GAC²² and was further validated by a study that demonstrated equivalent immunogenicity from a hexasaccharide hapten as native GAC when conjugated to TT⁶³.

To address potential vaccine safety concerns, van Sorge et al., used modified GAC devoid of GlcNAc (GAC^{PR}) extracted from a genetically modified Strep A strain and conjugated to *S. pneumoniae* protein SP0435¹⁰. They showed that antibodies generated from the vaccine promoted phagocytic killing of multiple Strep A serotypes and protected mice against systemic infection following passive immunisation with rabbit anti-GAC antisera¹⁰. The conjugation methodology was unique using a Multiple Antigen Presenting System (MAPS)⁶⁴, incorporating chemical methods for biotinylation of GAC to allow for a non-covalent bond between GAC and SP0435 containing a biotin-binding domain. This genetically mutated GAC was also able to show disease reduction with partial protection from bacteraemia and skin infections in further studies⁵⁶, without any cross-reactivity with human heart or brain tissue lysates³⁵. Further justification for modified GAC has been provided by a recent study using self-adjuvanting lipopeptides³⁶. Using synthetic repeating unit epitopes this study demonstrated that the GlcNAc residue was not essential for a robust immune response to GAC. Instead, replacement with a third rhamnose could substitute for this structurally related immunogenicity, with a higher anti-GAC IgG response and equivalent or higher killing activity on Strep A strains than the wildtype repeating unit. These studies laid the foundation for interest in GAC as a viable broadly protective vaccine antigen.

The first study with a “double-hit” approach to a GAC glycoconjugate demonstrated that conjugation of modified GAC to Strep A protein ADI (arginine deiminase protein) gave a robust immune response to both components, without a loss of immunogenicity to the protein component compared with a protein alone response⁵⁶. Following this, conjugation of synthetic polysaccharides (tri-, hexa- and nonasaccharides) to C5a peptidase (ScpA) demonstrated superior functional immunity of antibody binding to Strep A and opsonophagocytic killing over conjugation to CRM₁₉₇ or TT with hexasaccharide haptens, and equivalent immunogenicity between carriers with a nonasaccharide^{65,66}.

Recent studies have highlighted the need to preserve protein epitopes when using species specific protein carriers. This was demonstrated by di Benedetto et al., where the benefits and disadvantages of different conjugation chemistries was investigated with native GAC conjugated to CRM₁₉₇. This revealed that random conjugation demonstrated an equivalent anti-GAC IgG response to selective conjugation but affected the anti-protein response⁵⁷. Despite high anti-GAC IgG titres from random conjugation of GAC to three Strep A proteins SLO, SpyAD and SpyCEP, the anti-protein responses were significantly impacted and resulted in a loss of functional immunity, such as neutralisation of SpyCEP IL-8 cleavage activity. Random conjugation produces a mesh-like structure, which has been demonstrated to be effective for anti-polysaccharide responses, but likely will mask immunologically relevant protein epitopes. Selective conjugation, meanwhile, is likely to be the most effective method for “double-hit” glycoconjugates, as demonstrated with a site-specific click-chemistry approach to produce a SpyAD-GAC^{PR} conjugate³⁵. In this study, four lysine residues were replaced with a non-native amino acid, p-azidomethyl phenylalanine (pAMF) in a cell-free expression system, as target sites for dibenzocyclooctyne (DMCO)




Fig. 3 Glycoconjugate production using *C. jejuni* PglB bioconjugation inside *E. coli* host cells. **a** Plasmids containing genes encoding polysaccharide (PS) biosynthesis (green), the protein acceptor (blue) containing a glycotag (yellow star), and the oligosaccharyltransferase (OST) PglB (red) are co-transformed into *E. coli* host cells. **b** Bioconjugates are produced as follows; (1) The PS biosynthesis locus is expressed and built onto undecaprenol-pyrophosphate (Und-PP) lipid linkers within the inner membrane. (2) The PS is flipped from the cytoplasm to the periplasm by a specific flippase enzyme. (3) Synthesised carrier proteins are exported to the periplasm through the Sec secretion system. (4) In the periplasm both the PS, and the carrier protein containing a specific glycotag can be recognised by the PglB OST enzyme. PglB transfers the PS from Und-PP onto the asparagine residue within the glycotag D/E-X-N-X-S/T motif on the fully folded carrier protein, resulting in protein glycosylation. (5) An inexhaustible supply of glycoproteins can be subsequently purified from *E. coli* cells.

derivatised modified GAC (conjugated through 1-Cyano-4-dimethyl aminopyridinium tetrafluoroborate, CDAP, chemical conjugation). IgG titres against SpyAD were high for both conjugate and protein alone, and improved shifts in antibody binding by flow cytometry for SpyAD-GAC^{PR} compared with SpyAD alone. Combination of SpyAD-GAC^{PR} with SLO and C5a peptidase resulted in improved protection in both passive and active immunisation murine challenge models compared with components alone, with no cross-reactivity observed to human heart lysates. This site-directed click chemistry approach was further investigated with SLO as a carrier protein for GAC^{PR62}. Conjugates showed high antibody titres for GAC in all conjugates produced, and similar antibody titres for SLO in all conjugates produced compared with SLO variants alone. A specific SLO neutralising assay was not conducted to demonstrate retained functional neutralising antibodies, but an *in vivo* model with intraperitoneal challenge showed significant protection with the SLO-GAC^{PR} conjugate compared with immunisation with SLO and CRM₁₉₇-GAC^{PR62}. This was a successful demonstration that site-specific conjugation of GAC to a Strep A carrier protein provides effective antibody responses to both components and superior protection *in vivo*.

Though promising, the conjugates here relied on costly chemical conjugation, which has limitations such as technical challenges, low product yields, and batch to batch variation⁶⁷. Alternative conjugation methods and vaccine structures are therefore of benefit to Strep A vaccinology to reduce costs and improve manufacturing consistency.

Protein Glycan Coupling Technology

In recent years the development of Protein Glycan Coupling Technology (PGCT) or bioconjugation has provided an alternative and in some cases superior method to chemical conjugation^{68,69}. PGCT has the potential to simplify glycoconjugate production, and several vaccines using this technology are in clinical trials^{70–73}.

PGCT relies on the innate capability of certain prokaryotic cells to synthesise polysaccharides and attach them to proteins as a post-translational modification. Using PGCT the polysaccharide attachment on the carrier protein can be an asparagine (*N*-linked)

or serine/threonine (*O*-linked) amino acid glycosylation^{74,75}. *N*-linked glycosylation, specifically using oligosaccharyltransferase (OST) PglB from *Campylobacter jejuni* (CjPglB), is the most applied bioconjugation approach⁷⁵. The key experiment for glycoengineering and vaccine development was the demonstration that the *pglB* operon could be cloned and fully expressed in *E. coli* cells for functional recombinant glycosylation⁷⁶.

Polysaccharides for exploitation of this system, are first synthesised onto an undecaprenol-pyrophosphate (Und-PP) lipid linker attached to the cytoplasmic leaflet of the inner membrane. A flippase enzyme is then responsible for the translocation of the fully synthesised polysaccharide across to the periplasm, where it can subsequently be recognised by PglB or other OST and attached onto a recognition sequon on a given acceptor protein (Fig. 3)^{77,78}. The protein recognition sequon for *N*-linked glycosylation is D/E-X-N-X-S/T, where X represents any amino acid except proline, and positive D/E and S/T amino acids are at the ± 2 positions, pivotal in locating asparagine (N) as the acceptor amino acid⁷⁹. Unlike eukaryotic glycosylation, bacterial *N*-linked glycosylation occurs after protein folding, therefore to be accessible to PglB engineered Glycotags of the D/E-X-N-X-S/T sequon can be added to the N- and C- termini of carrier proteins allowing enhanced glyco-modification in vaccine design^{80,81}.

PGCT is a feasible alternative to chemical conjugation⁷⁸, with benefits such as *E. coli* systems producing inexhaustible fully synthesised recombinant polysaccharide resources, and readily purified glycoproteins at reduced costs and improved yields⁸². This great promise is currently limited by PglB specificity, such that only polysaccharides with a reducing end containing an acetamido group at the C2 position are permissive for wildtype enzyme transfer^{79,83}. However, this limitation can be mitigated by modification of the PglB enzyme by directed evolutionary mutagenesis to improve transfer compatibility and efficiency^{84–86}.

Alternative OST enzymes are also widely used for *N*-linked glycosylation^{74,77,87,88}, and *O*-linked OST enzymes, such as PglI and PglS for glycans with galactose^{89,90} and glucose end groups^{91,92}. As the discovery and understanding of bacterial OSTs increases there will be further opportunities to develop PGCT for custom designed glycoconjugate vaccines such that theoretically almost any glycan could be coupled to any protein. In addition,




Fig. 4 The *gac* operon in *Strep A* produces GAC containing a polyrhamnose backbone decorated with alternate GlcNAc side chains. **a** Schematic representation of the *gac* operon (*gacA-L*) in *Strep A*. Horizontal arrows represent each gene designation with colour denoting predicted gene function. Green, polyrhamnose biosynthesis; blue, GlcNAc biosynthesis. **b** Schematic diagram of GAC biosynthesis. GAC biosynthesis is initiated on lipid linked GlcNAc attached to the inner leaflet of the periplasmic membrane for polyrhamnose synthesis catalysed by rhamnosyltransferase enzymes (GacBCFG). After polymerisation, the polyrhamnose backbone is flipped to the outer leaflet by an ABC transporter (GacDE complex) before GlcNAc (GacL) and glycerol phosphates (GacH) are transferred to the polyrhamnose backbone as a sidechain modification. A LytR-CpsS-Psr (LCP) phosphotransferase protein is hypothesised to attach GAC to peptidoglycan via a phosphodiester bond¹⁷.

PGCT provides the opportunity to couple multiple glycans at precise positions on a given carrier protein, which has been demonstrated to increase glycoconjugate vaccine efficacy⁹³.

Furthermore, PGCT offers benefits such as minimal alteration to the protein carrier and target recombinant polysaccharide, dissimilar to chemical conjugation approaches which are often harsher in their attachment method.

GAC biosynthesis and exploitation of recombinant polyrhamnose structures

GAC polymers are encoded by a conserved 12-gene cluster termed *gacA-L*^{8,10} (Fig. 4a). The operon is highly conserved with one study finding that 2017 of 2083 tested *Strep A* genomes had >70% DNA sequence similarity for the entire 12-gene cluster⁸ supporting observations from a smaller dataset¹⁰. Within the operon the first seven genes (*gacA-G*) encode biosynthesis of the polyrhamnose backbone, conserved across other streptococci groups, specifically A, B, C and G⁹⁴, whilst 3 genes (*gacI-K*) are known to be implicated in GlcNAc sidechain attachment and modification¹⁰. To date not all genes have been fully characterised, and some are believed not to be essential to survival due to frameshift mutations in some *Strep A* genomes^{8,13}. However, the rhamnose encoding genes¹⁰ and the availability of L-rhamnose substrates¹² have been shown to be essential to *Strep A* survival.

There is a strong case for the exploitation of GAC with PGCT. A rhamnose backbone structure related to GAC in *Streptococcus mutans* has previously been expressed and exported to the surface of *E. coli* cells, altering the LPS profiles, with similarity shown between native rhamnose biosynthesis pathways in *S.*

mutans and the reconstituted pathways expressed and synthesised in *E. coli* host cells⁹⁵. Recently, Castro et al followed a similar recombinant approach, where *E. coli* cells encoded the genes for GAC rhamnose backbone biosynthesis and produced Outer Membrane Vesicles (OMVs) loaded with the carbohydrate⁹⁶. Their work has shown that the GAC rhamnose backbone can be built successfully in *E. coli* with the correct chemical structure and produces antibodies in mice and rabbits that can target *Strep A* serotypes. This system can now serve as a foundation for PGCT with *Strep A* antigens.

GAC has a GlcNAc at the reducing end, suggesting it would be suitably recognised by CjPglB, however the following sugar in the chain, rhamnose, is attached to the GlcNAc by a β -1,4 linkage by GacB⁹⁴. This particular linkage may not be recognised or transferred by CjPglB efficiently, as demonstrated by studies assessing CjPglB transfer capability of a GlcNAc β -1,4 GlcNAc disaccharide attached to either a non-native eukaryotic isoprene lipid linker⁹⁷, or a native Und-PP lipid linker^{98,99}. Therefore, alternative strategies may be required to transfer *Strep A* rhamnose polymers by PGCT.

Concluding remarks

The global imperative to develop a *Strep A* vaccine has resulted in a burgeoning field of GAS targeted glycoconjugate vaccines. Following initial studies demonstrating a robust immune response to GAC, focus has shifted to “double-hit” glycoconjugates incorporating species specific carrier proteins along with GAC, either native or modified to remove GlcNAc side chains. Studies have suggested that milder selective approaches are more

appropriate for “double-hit” glycoconjugates to prevent shielding of protein immune epitopes, as they are less likely to disrupt the protein structure and conformation, ensuring its role as a T cell epitope carrier, as well as a protective antigen itself. However, such approaches may reduce overall polysaccharide attachment due to limiting the region of the chain which is activated and available for attachment¹⁰⁰. This in some respects is similar to bioconjugation where protein carriers are engineered with glycotags at the N- and C- termini, which is more likely to preserve the protein’s stability and structure, and therefore B and T cell epitopes⁸¹. Investigation of the use of bioconjugation for generation of a Strep A glycoconjugate would help lead the way to a cost-effective Strep A vaccine.

Studies with Generalised Modules for Membrane Antigens (GMMA) showed that different mechanisms of presentation, specifically the structure and chain length of the polysaccharide antigen, dictates the dominance of T cell responses. Based on a *S. typhi* capsule glycoconjugate, shorter polysaccharide antigens appear to require more T cell help for robust immune stimulation compared to glycoconjugates containing longer polysaccharide antigens¹⁰¹. Therefore, carrier proteins which can strongly stimulate T cell help responses may be necessary for future vaccines with GAC. Such T cell activity is important in the magnitude as well as response duration, as without T cell activity apoptosis of polysaccharide-specific B cells in the spleen and depletion of polysaccharide-specific B cells in the bone marrow can occur leading to hypo-responsiveness¹⁰¹. Such impact of polysaccharide size on B and T cell interactions is known to be antigen-specific^{41,100}, requiring future studies to focus on deciphering the best protein carrier to induce T cell activity and enable GAC directed memory responses.

There has also been increasing interest in novel non-protein carriers for GAC. Gold nanoparticles have been shown to successfully present GAC for recognition by immune sera, though its own intrinsic immunogenicity was not assessed¹⁰². Self-adjuvating lipopeptides have shown great promise with GAS peptides and have also shown a robust immune response against GAC³⁶, with recent studies taking this further showing strong immune responses to conjugated GMMA¹⁰³ and recombinant OMVs⁹⁶. Such technologies will help to build up an understanding of a protective immune response to GAC and derivatives, and push for a low-cost solution to the global need for a vaccine. Recombinant technology, particularly PGCT, may answer this need with its low-cost, inexhaustible and renewable supply along with its ability for rational design⁸². Carrier proteins and glycans can easily be manipulated such that glycosylation sites can be precise, glycan number increased as needed and chain lengths potentially controlled.

In summary, GAC is a conserved antigen, found in all isolated Strep A strains, a property which recommends it for a universal global vaccine against Strep A infection. Investigation into cost-effective solutions for glycoconjugate production is imperative for a global vaccine, particularly to target low- and middle-income countries. Newer technologies such as PGCT and GMMA/OMVs show great promise towards this aim and examination with GAC is warranted. Impressive progress has been made towards a Strep A glycoconjugate vaccine, and further investigation into production of “double-hit” conjugates and their protection from infection would be a sweet reward.

Received: 5 October 2022; Accepted: 27 February 2023;

Published online: 28 March 2023

REFERENCES

- Bonanni, P. Demographic impact of vaccination. *Vaccine* **17**, 120–125 (1999).
- Vekemans, J. et al. The path to Group A Streptococcus vaccines: WHO research and development technology roadmap and preferred product characteristics. *Clin. Infect. Dis.* **16**, 877–883 (2019).
- Fay, K. et al. Patterns of antibiotic nonsusceptibility among invasive group A Streptococcus Infections—United States, 2006–2017. *Clin. Infect. Dis.* **73**, 1957–1964 (2021).
- WHO. *The Current Evidence for the Burden of Group A Streptococcal Diseases*. <https://apps.who.int/iris/handle/10665/69063> (2005).
- Watkins, D. A. et al. Global, regional, and national burden of rheumatic heart disease 1990–2015. *N. Engl. J. Med.* **377**, 713–722 (2017).
- Herdman, M. T. et al. Clinical management and impact of scarlet fever in the modern era: findings from a cross-sectional study of cases in London, 2018–2019. *BMJ Open* **11**, e057772 (2021).
- Rümke, L. W. et al. Dominance of M1UK clade among Dutch M1 Streptococcus pyogenes. *Lancet Infect. Dis.* **20**, 539–540 (2020).
- Davies, M. R. et al. Atlas of Group A Streptococcal vaccine candidates compiled using large-scale comparative genomics. *Nat. Genet.* **51**, 1035–1043 (2019).
- Lancefield, R. A serological differentiation of human and other groups of hemolytic Streptococci. *J. Exp. Med.* **57**, 571–595 (1933).
- van Sorge, N. M. et al. The classical lancefield antigen of group a streptococcus is a virulence determinant with implications for vaccine design. *Cell Host Microbe* **15**, 729–740 (2014).
- Mccarty, M. The lysis of Group A hemolytic Streptococci by extracellular enzymes of Streptomyces albus. Nature of the cellular substrate attacked by the lytic enzymes. *J. Exp. Med.* **96**, 569–580 (1952).
- van der Beek, S. L. et al. GacA is essential for Group A Streptococcus and defines a new class of monomeric dTDP-4-dehydrorhamnose reductases (RmlD). *Mol. Microbiol.* **98**, 946–962 (2015).
- Henningham, A. et al. Virulence role of the GlcNAc side chain of the Lancefield cell wall carbohydrate antigen in non-M1-serotype Group A Streptococcus. *MBio* **9**, 1–12 (2018).
- Kabanova, A. et al. Evaluation of a group A Streptococcus synthetic oligosaccharide as vaccine candidate. *Vaccine* **29**, 104–114 (2010).
- Hamada, S. et al. Isolation and immunochemical characterization of carbohydrate antigens prepared from Group A Streptococcus pyogenes. *Zentralbl. Bakteriologie. Mikrobiol. Hyg. A* **256**, 37–48 (1983).
- Rush, J. S. et al. The molecular mechanism of N-acetylglucosamine side-chain attachment to the Lancefield Group A carbohydrate in Streptococcus pyogenes. *J. Biol. Chem.* **292**, 19441–19457 (2017).
- Rush, J. S. et al. PplD is a de-N-acetylase of the cell wall linkage unit of Streptococcal rhamnopolysaccharides. *Nat. Commun.* **13**, 590 (2022).
- Kreis, U. C., Varma, V. & Pinto, B. M. Application of two-dimensional NMR spectroscopy and molecular dynamics simulations to the conformational analysis of oligosaccharides corresponding to the cell-wall polysaccharide of Streptococcus Group A. *Int. J. Biol. Macromol.* **17**, 117–130 (1995).
- Edgar, R. J. et al. Discovery of glycerol phosphate modification on Streptococcal rhamnose polysaccharides. *Nat. Chem. Biol.* **15**, 463–471 (2019).
- van Hensbergen, V. P. et al. Streptococcal Lancefield polysaccharides are critical cell wall determinants for human Group IIA secreted phospholipase A2 to exert its bactericidal effects. *PLoS Pathog.* **14**, e1007348 (2018).
- Reimer, K. B., Gidney, M. A. J., Bundle, D. R. & Mario Pinto, B. Immunochemical characterization of polyclonal and monoclonal Streptococcus Group A antibodies by chemically defined glycoconjugates and synthetic oligosaccharides. *Carbohydr. Res.* **232**, 131–142 (1992).
- Michon, F. et al. Doubly branched hexasaccharide epitope on the cell wall polysaccharide of Group A Streptococci recognized by human and rabbit antisera. *Infect. Immun.* **73**, 6383–6389 (2005).
- Weimar, T., Harris, S. L., Pitner, J. B., Bock, K. & Pinto, B. M. Transferred nuclear overhauser enhancement experiments show that the monoclonal antibody strep 9 selects a local minimum conformation of a Streptococcus Group A trisaccharide-hapten. *Biochemistry* **34**, 13672–13681 (1995).
- Stuik-Prill, R. & Pinto, B. M. Conformational analysis of oligosaccharides corresponding to the cell-wall polysaccharide of the Streptococcus Group A by Metropolis Monte Carlo simulations. *Carbohydr. Res.* **279**, 59–73 (1995).
- Johnson, M. A. & Pinto, B. M. Saturation transfer difference 1D-TOCSY experiments to map the topography of oligosaccharides recognized by a monoclonal antibody directed against the cell-wall polysaccharide of Group A Streptococcus. *J. Am. Chem. Soc.* **124**, 15368–15374 (2002).
- Adderson, E. E., Shikhman, A. R., Ward, K. E. & Cunningham, M. W. Molecular analysis of polyreactive monoclonal antibodies from rheumatic carditis: human anti-N-acetylglucosamine / anti-myosin antibody V region genes. *J. Immunol.* **161**, 2020–2031 (1998).
- Shikhman, A. R., Cunningham, M. W., Hildebrand, W., Fischetti, V. A. & Cunningham, M. W. Immunological mimicry between N-acetyl-beta-D-glucosamine

- and cytokeratin peptides. Evidence for a microbially driven anti-keratin antibody response. *J. Immunol.* **152**, 4375–4387 (1994).
28. Martins, T. B. et al. Comprehensive analysis of antibody responses to Streptococcal and tissue antigens in patients with acute rheumatic fever. *Int. Immunol.* **20**, 445–452 (2008).
 29. Dudding, B. A. & Ayoub, E. M. Persistence of Streptococcal Group A antibody in patients with rheumatic valvular disease. *J. Exp. Med.* **128**, 1081–1098 (1968).
 30. Faé, K. C. et al. How an autoimmune reaction triggered by molecular mimicry between Streptococcal M protein and cardiac tissue proteins leads to heart lesions in rheumatic heart disease. *J. Autoimmun.* **24**, 101–109 (2005).
 31. Sabharwal, H. et al. Group A *Streptococcus* (GAS) carbohydrate as an immunogen for protection against GAS infection. *J. Infect.* **193**, 129–135 (2006).
 32. Salvadori, L. G., Blake, M. S. & Mccarty, M. Group A *Streptococcus*-liposome ELISA antibody titers to group a polysaccharide and opsonophagocytic capabilities of the antibodies. *J. Immunol.* **171**, 593–600 (1995).
 33. Zimmerman, R. A., Auernheimer, A. H. & Taranta, A. Precipitating antibody to Group A Streptococcal polysaccharide in humans. *J. Immunol.* **107**, 832–83241 (1971).
 34. Zabriskie, J. B., Poon-King, T., Blake, M. S., Michon, F. & Yoshinaga, M. Phagocytic, serological, and protective properties of Streptococcal Group A carbohydrate antibodies. *Adv. Exp. Med. Biol.* **418**, 917–919 (1997).
 35. Gao, N. J. et al. Site-specific conjugation of cell wall polyrhannose to protein SpyAD envisioning a safe universal group a streptococcal vaccine. *Infect. Microbes Dis.* **3**, 87–100 (2021).
 36. Khatun, F. et al. Immunogenicity assessment of cell wall carbohydrates of group A *Streptococcus* via self-adjuvanted glyco-lipopeptides. *ACS Infect. Dis.* **7**, 390–405 (2021).
 37. Barrett, D. J. Human immune responses to polysaccharide antigens: an analysis of bacterial polysaccharide vaccines in infants. *Adv. Pediatr.* **32**, 139–158 (1985).
 38. Weintraub, A. Immunology of bacterial polysaccharide antigens. *Carbohydr. Res.* **338**, 2539–2547 (2003).
 39. Kelly, D. F., Pollard, A. J. & Moxon, E. R. Immunological memory: the role of B cells in long-term protection against invasive bacterial pathogens. *JAMA* **294**, 3019–3023 (2005).
 40. Pollard, A. J., Perrett, K. P. & Beverley, P. C. Maintaining protection against invasive bacteria with protein-polysaccharide conjugate vaccines. *Nat. Rev. Immunol.* **9**, 213–220 (2009).
 41. Costantino, P., Rappuoli, R. & Berti, F. The design of semi-synthetic and synthetic glycoconjugate vaccines. *Expert Opin. Drug Discov.* **6**, 1045–1066 (2011).
 42. Avci, F. Y., Li, X., Tsuji, M. & Kasper, D. L. A novel mechanism for glycoconjugate vaccine activation of the adaptive immune system. *Nat. Med.* **17**, 1602–1609 (2012).
 43. Dagan, R., Poolman, J. & Siegrist, C. A. Glycoconjugate vaccines and immune interference: A review. *Vaccine* **28**, 5513–5523 (2010).
 44. Micoli, F., Adamo, R. & Costantino, P. Protein carriers for glycoconjugate vaccines: History, selection criteria, characterization and new trends. *Molecules* **23**, 1–18 (2018).
 45. Bröker, M., Berti, F., Schneider, J. & Vojtek, I. Polysaccharide conjugate vaccine protein carriers as a “neglected valency” – Potential and limitations. *Vaccine* **35**, 3286–3294 (2017).
 46. Pichichero, M. E. Characteristics, development, and clinical trials. *Hum. Vaccin. Immunother.* **9**, 2505–2523 (2013).
 47. Schutze, M. P., Leclerc, C., Jolivet, M., Audibert, F. & Chedid, L. Carrier-induced epitopic suppression, a major issue for future synthetic vaccines. *J. Immunol.* **135**, 2319–2322 (1985).
 48. Schutze, M.-P., Leclerc, C., Vogel, F. R. & Chedid, L. Epitopic suppression in synthetic vaccine models: Analysis of the effector mechanisms. *Cell. Immunol.* **104**, 79–90 (1987).
 49. Etlinger, H. M. et al. Use of prior vaccinations for the development of new vaccines. *Science* **249**, 423–425 (1990).
 50. Falugi, F. et al. Rationally designed strings of promiscuous CD4+ T cell epitopes provide help to *Haemophilus influenzae* type b oligosaccharide: a model for new conjugate vaccines. *Eur. J. Immunol.* **31**, 3816–3824 (2001).
 51. Reglinski, M. et al. A recombinant conjugated Pneumococcal vaccine that protects against murine infections with a similar efficacy to Prevnar-13. *npj Vaccines* **3**, 1–11 (2018).
 52. Pozzi, C. et al. Opsonic and protective properties of antibodies raised to conjugate vaccines targeting six *Staphylococcus aureus* antigens. *PLoS ONE* **7**, e46648 (2012).
 53. Bensi, G. et al. Multi high-throughput approach for highly selective identification of vaccine candidates: the group A *Streptococcus* Case. *Mol. Cell. Proteom.* **11**, M111.015693 (2012).
 54. Rivera-Hernandez, T. et al. An experimental group A *Streptococcus* vaccine that reduces pharyngitis and tonsillitis in a nonhuman primate model. *MBio* **10**, 1–10 (2019).
 55. Rivera-Hernandez, T. et al. Vaccine-induced Th1-type response protects against invasive Group A *Streptococcus* infection in the absence of opsonizing antibodies. *MBio* **11**, 1–11 (2020).
 56. Rivera-Hernandez, T. et al. Differing efficacies of lead group a streptococcal vaccine candidates and full-length M protein in cutaneous and invasive disease models. *MBio* **7**, 1–9 (2016).
 57. Di Benedetto, R. et al. Rational design of a glycoconjugate vaccine against Group A *Streptococcus*. *Int. J. Mol. Sci.* **21**, 1–21 (2020).
 58. Castro, S. A. & Dorfmüller, H. C. A brief review on Group A *Streptococcus* pathogenesis and vaccine development. *R. Soc. Open Sci.* **8**, 1–17 (2021).
 59. Kuberan, B. & Linhardt, R. J. Carbohydrate Based Vaccines. *Curr. Org. Chem.* **4**, 653–677 (2000).
 60. Zarei, A. E., Almelhdar, H. A. & Redwan, E. M. Hib vaccines: past, present, and future perspectives. *J. Immunol. Res.* **2016**, 1–18 (2016).
 61. Crotti, S. et al. Defined conjugation of glycans to the lysines of CRM197 guided by their reactivity mapping. *ChemBioChem* **15**, 836–843 (2014).
 62. Kapoor, N. et al. Non-native amino acid click chemistry-based technology for site-specific polysaccharide conjugation to a bacterial protein serving as both carrier and vaccine antigen. *ACS Omega* **7**, 24111–24120 (2022).
 63. Auzanneau, F. I., Borrelli, S. & Pinto, B. M. Synthesis and immunological activity of an oligosaccharide-conjugate as a vaccine candidate against Group A *Streptococcus*. *Bioorg. Med. Chem. Lett.* **23**, 6038–6042 (2013).
 64. Zhang, F., Lu, Y.-J. & Malley, R. Multiple antigen-presenting system (MAPS) to induce comprehensive B- and T-cell immunity. *Proc. Natl Acad. Sci. USA* **110**, 13564–13569 (2013).
 65. Zhao, Y. et al. Synthesis and immunological studies of Group A *Streptococcus* cell-wall oligosaccharide–Streptococcal C5a peptidase conjugates as bivalent vaccines. *Org. Chem. Front.* **6**, 3589–3596 (2019).
 66. Wang, S. et al. Group A *Streptococcus* cell wall oligosaccharide-streptococcal C5a peptidase conjugates as effective antibacterial vaccines. *ACS Infect. Dis.* **6**, 281–290 (2020).
 67. Frasch, C. E. Preparation of bacterial polysaccharide-protein conjugates: analytical and manufacturing challenges. *Vaccine* **27**, 6468–6470 (2009).
 68. Langdon, R. H., Cuccui, J. & Wren, B. W. N-linked glycosylation in bacteria: an unexpected application. *Future Microbiol.* **4**, 401–412 (2009).
 69. Cuccui, J. & Wren, B. Hijacking bacterial glycosylation for the production of glycoconjugates, from vaccines to humanised glycoproteins. *J. Pharm. Pharmacol.* **67**, 338–350 (2015).
 70. Riddle, M. S. et al. Safety and immunogenicity of a candidate bioconjugate vaccine against *Shigella flexneri* 2a administered to healthy adults: a single-blind, randomized phase I study. *Clin. Vaccine Immunol.* **23**, 908–917 (2016).
 71. Huttner, A. et al. Safety, immunogenicity, and preliminary clinical efficacy of a vaccine against extraintestinal pathogenic *Escherichia coli* in women with a history of recurrent urinary tract infection: a randomised, single-blind, placebo-controlled phase 1b trial. *Lancet Infect. Dis.* **17**, 528–537 (2017).
 72. Hatz, C. F. R. et al. Safety and immunogenicity of a candidate bioconjugate vaccine against *Shigella dysenteriae* type 1 administered to healthy adults: a single blind, partially randomized Phase I study. *Vaccine* **33**, 4594–4601 (2015).
 73. Micoli, F. et al. Glycoconjugate vaccines: current approaches towards faster vaccine design. *Expert Rev. Vaccines* **18**, 881–895 (2019).
 74. Nothaft, H. & Szymanski, C. M. Protein glycosylation in bacteria: Sweeter than ever. *Nat. Rev. Microbiol.* **8**, 765–778 (2010).
 75. Iwashkiw, J. A., Voza, N. F., Kinsella, R. L. & Feldman, M. F. Pour some sugar on it: the expanding world of bacterial protein O-linked glycosylation. *Mol. Microbiol.* **89**, 14–28 (2013).
 76. Young, N. M. et al. Structure of the N-linked glycan present on multiple glycoproteins in the Gram-negative bacterium, *Campylobacter jejuni*. *J. Biol. Chem.* **277**, 42530–42539 (2002).
 77. Kowarik, M. et al. Definition of the bacterial N-glycosylation site consensus sequence. *EMBO J.* **25**, 1957–1966 (2006).
 78. Feldman, M. F. et al. Engineering N-linked protein glycosylation with diverse O antigen lipopolysaccharide structures in *Escherichia coli*. *Proc. Natl Acad. Sci. USA* **102**, 3016–3021 (2005).
 79. Lizak, C., Gerber, S., Numao, S., Aebi, M. & Locher, K. P. X-ray structure of a bacterial oligosaccharyltransferase. *Nature* **474**, 350–356 (2011).
 80. Kowarik, M. et al. N-linked glycosylation of folded proteins by the bacterial oligosaccharyltransferase. *Science* **314**, 1148–1150 (2006).
 81. Fisher, A. C. et al. Production of secretory and extracellular N-linked glycoproteins in *Escherichia coli*. *Appl. Environ. Microbiol.* **77**, 871–881 (2011).
 82. Dow, J. M., Mauri, M., Scott, T. A. & Wren, B. W. Improving protein glycan coupling technology (PGCT) for glycoconjugate vaccine production. *Expert Rev. Vaccines* **19**, 507–527 (2020).
 83. Wacker, M. et al. Substrate specificity of bacterial oligosaccharyltransferase suggests a common transfer mechanism for the bacterial and eukaryotic systems. *Proc. Natl Acad. Sci. USA* **103**, 7088–7093 (2006).

84. Ihssen, J. et al. Increased efficiency of *Campylobacter jejuni* N-oligosaccharyltransferase PglB by structure-guided engineering. *Open Biol.* **5**, 140227 (2015).
85. Ollis, A. A. et al. Substitute sweeteners: diverse bacterial oligosaccharyltransferases with unique N-glycosylation site preferences. *Sci. Rep.* **5**, 1–13 (2015).
86. Ihnken, L. A. et al. *Mutated PglB Oligosaccharyltransferase Enzymes*. 1–107 (2020).
87. Jervis, A. J. et al. Characterization of N-linked protein glycosylation in *Helicobacter pullorum*. *J. Bacteriol.* **192**, 5228–5236 (2010).
88. Mills, D. C. et al. Functional analysis of N-linking oligosaccharyl transferase enzymes encoded by deep-sea vent proteobacteria. *Glycobiology* **26**, 398–409 (2016).
89. Pan, C. et al. Biosynthesis of conjugate vaccines using an O-linked glycosylation system. *MBio* **7**, 1–11 (2016).
90. Sun, P. et al. Design and production of conjugate vaccines against *S. paratyphi A* using an O-linked glycosylation system in vivo. *npj Vaccines* **3**, 1–9 (2018).
91. Harding, C. M. et al. A platform for glycoengineering a polyvalent Pneumococcal bioconjugate vaccine using *E. coli* as a host. *Nat. Commun.* **10**, 1–11 (2019).
92. Feldman, M. F. et al. A promising bioconjugate vaccine against hypervirulent *Klebsiella pneumoniae*. *Proc. Natl Acad. Sci. USA* **116**, 18655–18663 (2019).
93. Marshall, L. E. et al. An O-antigen glycoconjugate vaccine produced using protein glycan coupling technology is protective in an inhalational rat model of tularemia. *J. Immunol. Res.* **2018**, 1–12 (2018).
94. Zorzoli, A. et al. Group A, B, C, and G *Streptococcus* Lancefield antigen biosynthesis is initiated by a conserved α -D-GlcNAc- β -1,4-L-rhamnosyltransferase. *J. Biol. Chem.* **294**, 15237–15256 (2019).
95. Shibata, Y., Yamashita, Y., Ozaki, K., Nakano, Y. & Koga, T. Expression and characterization of Streptococcal *rgp* genes required for rhamnan synthesis in *Escherichia coli*. *Infect. Immun.* **70**, 2891–2898 (2002).
96. Castro, S. A. et al. Recombinant Group A Carbohydrate backbone embedded into Outer Membrane Vesicles is a potent vaccine candidate targeting Group A *Streptococcus* from *Streptococcus pyogenes* and *Streptococcus dysgalactiae* subsp. *bioRxiv*. p. 1–21 Preprint at <https://www.biorxiv.org/content/10.1101/2021.11.12.468441v1> (2021).
97. Chen, M. M., Glover, K. J. & Imperiali, B. From peptide to protein: Comparative analysis of the substrate specificity of N-linked glycosylation in *C. jejuni*. *Biochemistry* **46**, 5579–5585 (2007).
98. Valderrama-Rincon, J. D. et al. An engineered eukaryotic protein glycosylation pathway in *Escherichia coli*. *Nat. Chem. Biol.* **8**, 434–436 (2012).
99. Glasscock, C. J. et al. A flow cytometric approach to engineering *Escherichia coli* for improved eukaryotic protein glycosylation. *Metab. Eng.* **47**, 488–495 (2018).
100. Jones, C. Vaccines based on the cell surface carbohydrates of pathogenic bacteria. *An. Acad. Bras. Cienc.* **77**, 293–324 (2005).
101. Micoli, F. et al. Short Vi-polysaccharide abrogates T-independent immune response and hyporesponsiveness elicited by long Vi-CRM197 conjugate vaccine. *Proc. Natl Acad. Sci. USA* **117**, 24443–24449 (2020).
102. Pitirollo, O. et al. Gold nanoparticles morphology does not affect the multivalent presentation and antibody recognition of Group A *Streptococcus* synthetic oligorhamnans. *Bioorg. Chem.* **99**, 1–8 (2020).
103. Micoli, F. et al. Generalized modules for membrane antigens as carrier for polysaccharides: impact of sugar length, density, and attachment site on the immune response elicited in animal models. *Front. Immunol.* **12**, 1–12 (2021).

ACKNOWLEDGEMENTS

All images created with BioRender.com.

AUTHOR CONTRIBUTIONS

K.B. and H.A.S. wrote the manuscript, H.D., B.W. and F.M. edited and provided feedback.

COMPETING INTERESTS

H.C.D. holds a patent on the rhamnose polysaccharide platform technology (WO2020249737A1). B.W.W. holds a patent for an *E. coli* bioconjugation strain (US20150344928A1).

ADDITIONAL INFORMATION

Correspondence and requests for materials should be addressed to Helen A. Shaw.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© Crown 2023