Credit: iStock/Getty Images Plus

Acute stroke produces a massive release of pro-inflammatory alarmin proteins

plaque load after stroke compared with control. Similarly, the use of RAGE-specific small interfering RNA or HMGB1-specific monoclonal antibodies reduced vascular inflammation after stroke.

The acute stroke response also involved sympathetic activation, which drove egress of monocytic cells from the bone marrow. “Enhanced vascular inflammation after stroke depended on a synergistic effect of sympathetic recruitment of monocytes from the bone marrow niche and subsequent activation of innate immune and endothelial cells by circulating alarmins,” summarize the researchers. Liesz and colleagues now plan to investigate whether “the exacerbation of vascular inflammation after stroke leads to plaque rupture with arterio-arterial embolism, which would explain the increased risk of recurrent stroke events”.

Gregory B. Lim

ORIGINAL ARTICLE Roth, S. et al. Brain-released alarmins and stress response synergize in accelerating atherosclerosis progression after stroke. *Sci. Transl. Med.* **10**, eaao1313 (2018)

This finding provides the first genetic evidence linking NLRP3 inflammasome activation to neutrophil accumulation and NETosis

and NETosis in plaques, which was reversed by *Nlrp3* or *Casp1* and *Casp11* deficiency. This finding “provides the first genetic evidence linking NLRP3 inflammasome activation to neutrophil accumulation and NETosis in atherosclerotic plaques”. However, “the mechanisms linking cholesterol accumulation in myeloid cells to inflammasome activation need to be further explored”, says Westertorp.

Patients with Tangier disease carry mutations in *ABCA1* and have higher plasma levels of IL-1 β compared with age-and-gender-matched individuals, suggesting conservation of cholesterol-dependent inflammasome activation in humans. Low levels of myeloid *ABCA1* and *ABCG1* are seen in diabetes mellitus and ageing. “Our findings point to the importance of inflammasome-targeted, anti-inflammatory therapy, such as IL-1 β antibodies,” concludes Westertorp.

Alexandra Le Bras

ORIGINAL ARTICLE Westertorp, M. et al. Cholesterol efflux pathways suppress inflammasome activation, NETosis and atherogenesis. *Circulation* <https://doi.org/10.1161/CIRCULATIONAHA.117.032636> (2018)

Credit: V. Summersby/Macmillan Publishers Limited

Redefining leukocytes in atherosclerosis

Distinct leukocyte populations have been associated with atherosclerosis development and progression or with atheroprotection. However, most of these subsets were defined using preselected markers, and the actual subset heterogeneity in atherosclerotic lesions and the exact contribution of each leukocyte population is not well defined. Two studies now take advantage of novel high-dimensional techniques to provide a more detailed insight into leukocyte phenotypic and transcriptional heterogeneity in atherosclerosis.

Combining mass cytometry (CyTOF) and single-cell RNA sequencing (single-cell RNA-seq) with bioinformatic tools, Dennis Wolf and colleagues found high leukocyte diversity in mouse atherosclerotic aortas, with 11 leukocyte populations with distinct phenotypic and spatial characteristics. By contrast, mouse healthy aortas had a small number of leukocyte subsets. The phenotypic diversity of the clusters was confirmed with a 35-panel CyTOF and conventional flow cytometry, and the subsets showed a high correlation with previously defined transcriptomes. Genetic deconvolution analysis of microdissected mouse aorta and of 126 human plaques showed that monocytes, macrophages, and T cells were predominantly present in plaques and B cells in the adventitia. Finally, absence of one T-cell cluster was associated with a higher cardiovascular risk in patients. “Our data suggest that the traditional classification of leukocytes based on principal haematopoietic lineages and known lineages is too coarse,” says Wolf. “We hope that our cell atlas will open new avenues in atherosclerosis research and inspire others to more precisely look at leukocyte diversity.” The researchers plan to study further some of the newly identified subsets to understand the biological meaning of their findings.

The team led by Alma Zernecke, Clément Cochain, and Antoine-Emmanuel Saliba, used single-cell RNA-seq to analyse the transcriptional profile of CD45+ cells from mouse healthy and atherosclerotic aortas. The analysis identified 13 cell clusters. A population of resident-like macrophages was found in both healthy and atherosclerotic aortas, whereas monocytes, monocyte-derived dendritic cells, and two macrophage populations were almost only detectable in atherosclerotic aortas. The two macrophage populations associated with atherosclerotic lesions were inflammatory macrophages enriched in *Il1b* and *TREM2*^{high} macrophages, characterized by a gene expression signature reminiscent of osteoclasts (which suggests a role in lesion calcification) and putative functions in lipid metabolism and catabolism, as revealed by differential gene expression and gene ontology enrichment analyses. *TREM2*^{high} macrophages were also detected in advanced atherosclerosis, and *TREM2* expression was found in human atherosclerotic lesions. “Our work provides a global overview of the transcriptional landscape of macrophages and other immune cells in atherosclerotic vessels with unprecedented resolution, and may prove a valuable resource for other researchers,” explain the investigators. The research team plans to assess whether similar macrophage populations are present in human atherosclerotic lesions and to investigate some of the newly identified targets that might control macrophage-mediated inflammation in atherosclerosis.

Irene Fernández-Ruiz

ORIGINAL ARTICLES Winkels, H. et al. Atlas of the immune cell repertoire in mouse atherosclerosis defined by single-cell RNA-sequencing and mass cytometry. *Circ. Res.* <https://doi.org/10.1161/CIRCRESAHA.117.312513> (2018) | Cochain, C. A. et al. Single-cell RNA-seq reveals the transcriptional landscape and heterogeneity of aortic macrophages in murine atherosclerosis. *Circ. Res.* <https://doi.org/10.1161/CIRCRESAHA.117.312509> (2018)