Credit: bymuratdeniz/Getty

GDF15 is an inflammation-induced hormone that facilitates survival during acute inflammation, via metabolic adaptations

patients in intensive care units increased survival in the GDF15-blocked mouse model of bacterial inflammation. In this model, the triglyceride treatment rescued defects in cardiac and thermal function.

In summary, the findings suggest that GDF15 is an inflammation-induced hormone that facilitates survival during acute inflammation, via metabolic adaptations that decrease tissue damage.

Shimona Starling

ORIGINAL ARTICLE Luan, H. H. et al. GDF15 is an inflammation-induced central mediator of tissue tolerance. *Cell* **178**, 1231–1244 (2019)

BONE

Osteocalcin linked to stress response

Credit: Science Photo Library — STEVE GSCHMEISSNER/Getty

Bone is an endocrine organ that is integral to a number of processes, ranging from energy metabolism to cognition. Now, new research from Gerard Karsenty and colleagues has furthered our understanding of bone by implicating it in the acute stress response.

Karsenty and colleagues previously posited that evolutionarily bone equipped vertebrates with the means to escape threats. For example, in addition to its mechanical properties, bone facilitates hearing, which ensures physical threats can be heard. Furthermore, osteocalcin, a bone derived hormone, is key to physiological processes that ensure vertebrates can predict and avoid danger (via actions on memory formation) and employ fight or flight reactions (via actions on exercise capacity).

In addition to the aforementioned processes, the acute stress response is inextricably linked to an animal's ability to avoid or escape danger as it helps regulate and maintain homeostasis during a fight or flight response. While the sympathetic nervous system is considered the key regulator of the acute stress response, the response is accompanied by increases in circulating glucocorticoid hormones, which are synthesized in the adrenal cortex. "We thought, however, that the increase in glucocorticoid hormones could not account for an endocrine mediation of the stress response as steroids such as glucocorticoids need hours to act, whereas the stress response is quasi immediate," explains Karsenty. "This realisation brought in the notion that an endocrine mediation of the acute stress response would be best mediated by a peptide hormone."

For this reason, and considering the known roles bone has in enabling vertebrates to escape threats, Karsenty and colleagues investigated whether bone-derived hormones contribute to the acute stress response. Through the use of animal models, including loss of function mutants, as well as human participants, the team found that circulating osteocalcin levels quadruple very quickly (within 2 minutes in mice) when rodents or humans are exposed to stressors.

Karsenty and colleagues also found that bone, via osteocalcin, is required for the development of an acute stress response. "Osteocalcin signals in post-ganglionic parasympathetic neurons to inhibit their activity, thereby removing a break on the fight or flight response," adds Karsenty. The authors findings also help explain, in part, why in the absence of adrenal glands, rodents and humans develop a normal stress response. Mammals without adrenal glands still have circulating osteocalcin, which arises because glucocorticoids repress osteocalcin expression.

"We now need to unravel the road map linking the brain to glutamatergic neurons in bone," concludes Karsenty.

Alan Morris

ORIGINAL ARTICLE Berger, J. M. et al. Mediation of the acute stress response by the skeleton. *Cell Metab.* <https://doi.org/10.1016/j.cmet.2019.08.012> (2019)

Credit: Daniel Grizelj/Getty

for the leptin resistance induced by GIP. "Our study shows that GIP is a circulating signal that drives neural leptin resistance and that GIP acts in the brain to mediate the regulation of energy balance," explains Fukuda.

"Beyond leptin resistance, it will be interesting to determine the exact roles of GIP in mediating other features of obesity in the central nervous system, such as hypothalamic inflammation," concludes Fukuda. "We plan to further prove our model by using mouse genetics with Cre-LoxP technology."

Claire Greenhill

ORIGINAL ARTICLE Kaneko, K. et al. Gut-derived GIP activates central Rap1 to impair neural leptin sensitivity during overnutrition. *J. Clin. Invest.* <https://doi.org/10.1172/JCI126107> (2019)

GIP is a circulating signal that drives neural leptin resistance