Obesity increases the risk of developing type 2 diabetes mellitus (T2DM). However, it is incompletely understood why some individuals with obesity develop the disease and others do not.

To answer this question, Mototsugu Nagao, Lena Eliasson and colleagues studied pancreatic islets from people with obesity who had developed T2DM and compared them with islets from people with obesity who had not developed T2DM.

“When Mototsugu Nagao came to my laboratory, he had published a paper describing that pancreatic islets from mice prone to diet-induced diabetes mellitus had high expression of the fatty acid transporter CD36 (*Cd36*) and, at the same time, lower expression of some exocytotic genes,” says Eliasson, co-corresponding author of the study. “We decided to investigate together if this was also the case in human islets, and if it might have a pathogenic role in T2DM in relation to obesity.”

The researchers observed that islets from donors with obesity-associated T2DM had reduced insulin secretion and β -cell exocytosis compared with islets from donors with obesity but not T2DM. Furthermore, dysfunctional insulin secretion and impaired exocytosis were associated with increased expression of CD36.

Using a rat-derived β -cell line the team investigated the mechanisms underlying the human findings. Their findings suggest that increased levels of CD36 modulate β -cell insulin signalling and suppress insulin granule exocytosis.

To test the therapeutic potential of their work, the researchers treated a human β -cell line with CD36-neutralizing antibodies. They observed that the treatment increased the expression of exocytosis-associated proteins and enhanced insulin secretion. “Taken together, our findings further highlight the pathogenic role of β -cell CD36 in the development of T2DM, especially in relation to obesity,” explains Eliasson.

“We believe that the blockage of CD36 function could be an attractive approach to improve insulin secretion, especially in obesity-related T2DM,” summarizes Eliasson. “A further study to test the CD36-neutralizing antibody on human islets of donors with obesity and T2DM is warranted as the next step.”

Anna Kriebs, Associate Editor,
Nature Communications

ORIGINAL ARTICLE Nagao, M. et al. Potential protection against type 2 diabetes in obesity through lower CD36 expression and improved exocytosis in β -cells. *Diabetes* <https://doi.org/10.2337/db19-0944> (2020)

OBESITY

Maternal obesity disrupts hypothalamic development

In the USA, ~50% of pregnant women have overweight or obesity. Now, new data in mice show that maternal obesity can disrupt the development of the hypothalamic circuits that are involved in metabolic regulation and feeding. Furthermore, a high-fat, high-sucrose (HFHS) diet during pregnancy and lactation was associated with lifelong metabolic alterations in the offspring, including increased body weight, fat mass and insulin resistance.

“Previous studies demonstrated that endoplasmic reticulum (ER) stress has an important role in obesity-induced insulin resistance and type 2 diabetes mellitus (T2DM) during adult life,” explains corresponding author Sebastien Bouret. “However, the role of ER stress during early life and whether it could contribute to maternal obesity-induced metabolic dysfunction in the offspring have

not yet been established.” Therefore, lead author Soyoung Park and colleagues investigated the link between maternal diet-induced obesity and ER stress in offspring and whether ER stress contributes to hypothalamic development.

The authors examined the consequences of maternal obesity on the offspring’s metabolism and hypothalamic development and whether it also induced ER stress in female mice that were fed a HFHS diet during pregnancy and lactation. The team conducted all of their analyses in metabolically relevant tissues, including the arcuate nucleus, paraventricular nucleus, pancreas, adipose tissue and liver.

“We found that maternal obesity causes abnormal development of hypothalamic circuits involved in feeding and metabolic regulation,” explains Bouret. “Moreover, we report that maternal HFHS feeding

BONE

Gut hormones involved in bone remodelling

Gut hormones seem to regulate bone metabolism, but the mechanisms are unclear. New research suggests that glucose-dependent insulinotropic polypeptide (GIP) and xenin (hormones secreted by enteroendocrine K cells) are key factors in regulating several aspects of bone metabolism.

Studies have shown that genetic deletion of the GIP receptor (GIPR) slightly increases trabecular bone mass. By contrast, mice given GIP analogues have increased trabecular bone mass. “It appears clear to us that a better understanding of the physiology of the GIP–GIPR pathway would lead to the development of second-generation medicines for the prevention and treatment of bone fragility,” explains author Guillaume Mabileau.

The researchers found that ablation of the cells that produce GIP in mice (GIP–DT mice) did not alter bone strength or bone physiology. However, in mice

that only lacked GIP (but retained expression of the other hormones produced by K cells; GIP–GFP–KI mice) trabecular bone mass was reduced, which was linked to increased bone fragility. These apparently conflicting results led the researchers to hypothesize that antagonistic actions of other gut hormones produced by K cells were involved in a feedback loop. Additional experiments revealed that xenin was involved in bone metabolism as part of a feedback loop that counteracts the action of GIP. “We also showed that other K cell-produced peptides, namely cholecystokinin, somatostatin and secretin, had very little or no effect on bone metabolism,” says Mabileau.

The researchers suggest that their results support the development of GIP analogues for the treatment of bone disorders featuring low bone mass and increased bone fragility. “We are actively working on the development