Reply to 'Can we predict microbial keystones?'

Samiran Banerjee, Klaus Schlaeppi^{1,2} and Marcel G. A. van der Heijden^{1,3}

In response to our recent Opinion article (Keystone taxa as drivers of microbiome structure and functioning. *Nat. Rev. Microbiol.* **16**, 567–576 (2018))¹, Röttgers and Faust (Can we predict keystones? *Nat. Rev. Microbiol.* <https://doi.org/10.1038/s41579-018-0132-y> (2018))² highlight an important discussion in microbial ecology: how accurately can keystones be predicted from co-occurrence networks? We agree with their claim that the identification of highly connected OTUs (hubs) in microbial networks does not necessarily reveal their role as keystones. In our Review article¹, we have extensively discussed the idea that the identification of keystones by network-based scores must be complemented with empirical data to uncover their true importance. Therefore, we summarized keystones into two categories: those identified through computational inference ('keystone OTUs') and those with explicit experimental evidence for their role in the microbiome ('keystone taxa'). While Röttgers and Faust only consider empirically established keystone taxa, our aim was to generate a candidate list for further validation but also to identify keystone OTUs, for which it might be a daunting task to collect enough empirical evidence to classify them as keystone taxa. Some microorganisms are uncultivable, or current methods are not sensitive enough to culture them. Moreover, without an initial statistical screening for potential keystones, it may be nearly impossible to identify them in view of the enormous diversity and complexity in microbial communities (for example, almost

any substrate on Earth is typically colonized by thousands of microbial taxa).

Röttgers and Faust also underline that the 'conditionality' of co-occurrence networks often confounds the accurate identification of keystones. Indeed, network-based scores can be biased by habitat filtering and other environmental factors^{3,4}, and this was also extensively discussed in our article. However, we posit that keystone OTUs may also be important in the community in case they have been consistently identified across ecosystems. For example, we speculated in our article that mycorrhizae, an important and widespread group of beneficial fungi that contribute substantially to plant nutrition, might also be keystones. Our recent study found that the majority of keystone OTUs in the wheat-root microbiome were arbuscular mycorrhizal fungi⁵. Moreover, our study found that some of the keystone OTUs were members of the *Dioszegia*, a genus that was identified as keystone taxa by Agler et al. (2016)⁶. The consistent identification of *Dioszegia* as keystone OTUs across studies not only highlights their importance in the plant microbiome but also indicates the usefulness of network scores for an initial identification of keystones, which can then be tested empirically. We would like to stress that owing to the ubiquitous nature and enormous diversity of microorganisms, we expect that there are many more keystone OTUs than the 200 listed in supplementary table 1 (REF.¹).

In summary, we wish to thank Röttgers and Faust for reinforcing the importance of empirical evidence in confirming the role of keystone OTUs. We explicitly stated that

network-mediated identification requires further proof and, thus, we proposed several approaches, including metagenomics and metatranscriptomics. Recent culturomics⁷ and microbiome-on-a-chip⁸ approaches may also be useful. In science, experimental evidence often comes years after theoretical proposition. A classic example of this is 'comammox', or complete nitrifiers, for which experimental validation⁹ came a decade after it was originally postulated¹⁰. As such, the identification of keystone OTUs will help to target key members of microbial communities.

Samiran Banerjee¹*, Klaus Schlaeppi^{1,2} and Marcel G. A. van der Heijden^{1,3}*

¹Plant-Soil Interactions, Department of Agroecology and Environment, Agroscope, Reckenholz, Zürich, Switzerland.

²Institute of Plant Sciences, University of Bern, Bern, Switzerland.

³Department of Plant and Microbial Biology, University of Zürich, Zürich, Switzerland.

*e-mail: samiran.banerjee@agroscope.admin.ch; marcel.vanderheijden@agroscope.admin.ch

<https://doi.org/10.1038/s41579-018-0133-x>

1. Banerjee, S., Schlaeppi, K. & van der Heijden, M. G. A. Keystone taxa as drivers of microbiome structure and functioning. *Nat. Rev. Microbiol.* **16**, 567–576 (2018).
2. Röttgers, L. & Faust, K. Can we predict keystones? *Nat. Rev. Microbiol.* <https://doi.org/10.1038/s41579-018-0132-y> (2018).
3. Faust, K. & Raes, J. Microbial interactions: from networks to models. *Nat. Rev. Microbiol.* **10**, 538–550 (2012).
4. Berry, D. & Widder, S. Deciphering microbial interactions and detecting keystone species with co-occurrence networks. *Front. Microbiol.* **5**, 219 (2014).
5. Banerjee, S. et al. Agricultural intensification reduces microbial network complexity and the abundance of keystone taxa in roots. Preprint at *bioRxiv* <https://doi.org/10.1101/416271> (2018).
6. Agler, M. T. et al. Microbial hub taxa link host and abiotic factors to plant microbiome variation. *PLOS Biol.* **14**, e1002352 (2016).
7. Lagier, J.-C. et al. Culture of previously uncultured members of the human gut microbiota by culturomics. *Nat. Microbiol.* **1**, 16203 (2016).
8. Stanley, C. E. & van der Heijden, M. G. A. Microbiome-on-a-chip: new frontiers in plant-microbiota research. *Trends Microbiol.* **25**, 610–613 (2017).
9. Daims, H. et al. Complete nitrification by *Nitrospira* bacteria. *Nature* **528**, 504–509 (2015).
10. Costa, E., Pérez, J. & Kreft, J. U. Why is metabolic labour divided in nitrification? *Trends Microbiol.* **14**, 213–219 (2006).

Competing interests

The authors declare no competing interests.