GENOME WATCH

Yeasts and how they came to be

Sara Calhoun, Stephen J. Mondo and Igor V. Grigoriev

This month's Genome Watch highlights a large-scale sequencing project that enriches our understanding of yeast evolution and diversity.

In contrast to most eukaryotes, fungi have relatively small and simple genomes, making them an attractive resource for large-scale genomic studies. Even more attractive, the budding yeast subphylum (Saccharomycotina) has some of the most straightforward genomes to assemble and annotate, as they are generally small (10–20 Mb) and intron-poor. The Saccharomycotina also harbour some of the most important fungi for bioindustry, being essential for converting sugars into ethanol and other valuable chemicals.

The genome of *Saccharomyces cerevisiae* was the first sequenced eukaryotic genome, which led to many breakthroughs in understanding eukaryotic biology and enabling genetic engineering¹. Now, as part of the Y1000+ project, with the ambitious goal to sequence over 1000 yeast genomes, Shen et al.² sequenced and annotated 220 of these fungi. Despite their small genomes, budding yeasts exhibit an extraordinary range across their functional repertoire. Combining these data with 100 published yeast genomes and phenotypic information on metabolic capabilities for all species in their study, the authors were able to make key connections between gene pathways and particular metabolic traits. For example, they inferred that the common ancestral budding yeast could assimilate

nitrate, xylose and galactose, but lacked the ability to ferment glucose.

This study also explored how distribution of these traits across the subphylum came to be. Through phylogeny reconstruction paired with metabolic analysis, the authors were able to trace the loss and gain of new traits through subphylum history. By inferring the traits of the budding yeast common ancestor, they implicate gene loss as a main driver of diversification. Since this work has been published, several follow-up studies have been conducted that use subsets of the genomic data from this project to explore budding yeast diversity and evolution.

To follow up on gene loss as the evolutionary driver of diversity, Steenwyk et al.³ explored a fast-evolving lineage in the genus *Hanseniaspora*, a cosmopolitan lineage that is often found in high abundance on grapes and in wine must. They found that this lineage lost many cell cycle and DNA repair genes and represents an unusual example of long-term survival of a hypermutator lineage. Hypermutator species have lost many genes in addition to these, harbouring the lowest number of genes compared to any other budding yeast. Furthermore, the observed gene losses were consistent with their known metabolic traits (for example, the inability to grow on various carbon sources).

Although gene losses are important for driving diversification in budding yeasts, horizontal gene transfer (HGT) is another interesting contributor to their evolution.

Kominek et al.⁴ identified the first example of an HGT event involving transfer of a full operon from bacteria into yeasts. This operon contained siderophore biosynthesis pathway genes, was functional, and enabled these yeasts to acquire a

new metabolic trait for iron uptake. By exploring conservation of iron uptake and storage systems across 175 fungi including 17 closely related genomes from the clade encompassing *Wickerhamiella* and *Starmerella* yeasts, the authors found that this biosynthetic capability came from a single HGT event at the root of *Wickerhamiella* and *Starmerella*. From 186 budding yeast genomes, over 800 genes were predicted to be acquired from bacteria by HGT, and most of these genes are associated with metabolism².

Thanks to the sequencing efforts of Shen et al. and others, we now have a rich sampling of an ancient (~400 million years old) eukaryotic subphylum. This provides us with the first opportunity to explore eukaryotic evolution at this magnitude and has already provided various important insights into the basic biology of eukaryotes. In addition to those discussed above, these data have been used to characterize variation and alterations of codon usage⁵ and reinvention of mating-type switching⁶. We expect that the completion of the Y1000+ project will provide an even fuller picture of the subphylum and has the potential to reveal many new and important insights into the biology of yeasts and eukaryotes at large.

Sara Calhoun, Stephen J. Mondo and Igor V. Grigoriev*
DOE Joint Genome Institute Walnut Creek,
California, USA

*e-mail: JGI-Microbe@lbl.gov

<https://doi.org/10.1038/s41579-019-0274-6>

1. Goffeau, A. et al. Life with 6000 genes. *Science* **274**, 546–567 (1996).
2. Shen, X. et al. Tempo and mode of genome evolution in the budding yeast subphylum. *Cell* **175**, 1533–1545 (2018).
3. Steenwyk, J. L. et al. Extensive loss of cell-cycle and DNA repair genes in an ancient lineage of bipolar budding yeasts. *PLOS Biology* **17**, e3000255 (2019).
4. Kominek, J. et al. Eukaryotic acquisition of a bacterial operon. *Cell* **176**, 1356–1366 (2019).
5. Labella, A. L. et al. Variation and selection on codon usage bias across an entire subphylum. *PLOS Genetics* **15**, e1008304 (2019).
6. Krassowski, T. et al. Multiple reinventions of mating-type switching during budding yeast evolution. *Current Biology* **29**, 2555–2562 (2019).

Competing interests

The authors declare no competing interests.

Credit: Philip Patenall/Springer Nature Limited