

Dynamical models to evaluate structure–function relationships in network neuroscience

Andrea I. Luppi¹, Joana Cabral¹, Rodrigo Cofre¹, Alain Destexhe¹, Gustavo Deco¹ and Morten L. Kringelbach¹

In their timely Review, Váša and Mišić provide an insightful review of the range of null models available for hypothesis testing in network neuroscience (Váša, F. & Mišić, B. Null models in network neuroscience. *Nat. Rev. Neurosci.* **23**, 493–504 (2022))¹. A central part of their Review is dedicated to generative models, understood as models for the processes that lead (brain) networks to exhibit the organization that they do. Here, we wish to draw attention to a distinct but

complementary kind of generative model that can be used to embody and then test null hypotheses in network neuroscience: namely, dynamical models of mechanisms. In these models, brain network structure (and, increasingly, additional aspects such as regional heterogeneity) gives rise to the ebb and flow of brain activity and functional connectivity that unfold over time^{2–4} (FIG. 1).

In a notable recent example⁵, the topology of the structural connectome underlying a

Kuramoto model was systematically varied, showing that a modular but not a random topology produces ‘events’ in the simulated time-resolved functional connectivity, analogous to those observed in empirical functional MRI signals. From the perspective of null models, this result showed that such events are attributable neither to motion and physiological artefacts nor solely to cognitive operations, as the model incorporates neither. Rather, the topology of the structural network is sufficient to determine whether such events will be observed in the functional dynamics. This work exemplifies how dynamical models can be used to arbitrate between plausible causal mechanisms for a phenomenon of interest, by explicitly building such mechanisms into the model and assessing whether the phenomenon in question emerges.

Another notable study enriched connectome-based mean-field models with PET-derived maps for the 5-HT_{2A} receptor,




Fig. 1 | Use of whole-brain dynamical models to test null hypotheses about the relationship among brain network structure, regional heterogeneity and brain activity. Dynamical whole-brain models simulate the emergence of brain activity and functional connectivity *in silico*, combining a model of local dynamics (red), with varying degrees of biological realism, and global connectivity (yellow), typically obtained from diffusion MRI tractography in humans or tract-tracing in animals. The role of regionally heterogeneous features of neurobiology in shaping brain function can be evaluated by modulating local dynamics according to empirical maps,

such as T1w:T2w ratio, receptor density from PET or transcriptomics, the principal component of gene expression (PC1), or excitation:inhibition ratio (among others). The heterogeneous model's goodness-of-fit to empirical functional data is then compared against a distribution of models enriched with null maps. The contribution of specific features of structural network topology to brain function can be evaluated by comparing the model's goodness-of-fit to empirical data against a distribution of null models, rewired to remove the network feature, or features, of interest. Figure adapted with permission from REF.⁷, Elife; REF.⁸, Elsevier; and REF.⁹, AAAS.

recapitulating blood oxygen level-dependent dynamics under the effects of the serotonergic psychedelic lysergic acid diethylamide (LSD)⁶. Null models enriched with alternative 5-HT receptors, or with uniform or scrambled 5-HT_{2A} maps, allowed the authors to embody and then reject alternative hypotheses whereby the spatial distribution of the 5-HT_{2A} receptor does not contribute to the dynamics of LSD. A subsequent transcriptomics-enriched model evaluated non-5-HT receptors and the role of spatial autocorrelation⁷, highlighting the synergy between different facets of null hypothesis testing in network neuroscience. Increasingly, dynamical models are being enriched with other aspects of neurobiology, such as T1w:T2w ratio⁸, excitatory:inhibitory ratio⁹, the principal component of gene expression⁹, or feedback versus feedforward connectivity¹⁰ (to name just a few in a fast-growing literature), representing additional avenues with which to embody and test hypotheses about annotated brain networks.

Overall, generative models of brain dynamics represent a powerful recent addition to the hypothesis-testing toolset of network neuroscientists. They provide an avenue to embody and assess mechanistic hypotheses, not only of how the organization of structural brain networks came to be, but also of how in turn it interacts with regional dynamics to shape the temporal unfolding of brain activity. We hope that explicit recognition of this flexible framework for testing mechanistic hypotheses will facilitate the emerging research on the interrelationships of brain network structure, dynamics and neuromodulation, bringing additional aspects of neurobiology into consideration for the study of both structural and functional brain networks.

Andrea I. Luppi^{1,2,3}, Joana Cabral^{4,5,6},
Rodrigo Cofre^{7,8}, Alain Destexhe^{7,9},
Gustavo Deco^{10,11,12} and
Morten L. Kringelbach^{4,5,6,13}

¹Division of Anaesthesia and Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK.

²Leverhulme Centre for the Future of Intelligence, University of Cambridge, Cambridge, UK.

³The Alan Turing Institute, London, UK.

⁴Life and Health Sciences Research Institute (ICVS), School of Medicine, University of Minho, Braga, Portugal.

⁵Centre for Eudaimonia and Human Flourishing, Linacre College, University of Oxford, Oxford, UK.

⁶Center for Music in the Brain, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark.

⁷Institute of Neuroscience (NeuroPSI-CNRS), Paris-Saclay University, Gif sur Yvette, France.

⁸CIMFAV-Ingemat, Facultad de Ingeniería, Universidad de Valparaíso, Valparaíso, Chile.

⁹European Institute for Theoretical Neuroscience, Gif sur Yvette, France.

¹⁰Center for Brain and Cognition, Computational Neuroscience Group, Universitat Pompeu Fabra, Barcelona, Spain.

¹¹Institució Catalana de la Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

¹²Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany.

¹³Department of Psychiatry, University of Oxford, Oxford, UK.

✉e-mail: al857@cam.ac.uk

<https://doi.org/10.1038/s41583-022-00646-w>

- Váša, F. & Mišić, B. Null models in network neuroscience. *Nat. Rev. Neurosci.* **23**, 493–504 (2022).
- Breakspear, M. Dynamic models of large-scale brain activity. *Nat. Neurosci.* **20**, 340–352 (2017).
- Shine, J. M. et al. Computational models link cellular mechanisms of neuromodulation to large-scale neural dynamics. *Nat. Neurosci.* **24**, 765–776 (2021).
- Suárez, L. E., Markello, R. D., Betzel, R. F. & Misić, B. Linking structure and function in macroscale brain networks. *Trends Cogn. Sci.* **24**, 302–315 (2020).
- Pope, M., Fukushima, M., Betzel, R. F. & Sporns, O. Modular origins of high-amplitude co-fluctuations

in fine-scale functional connectivity dynamics. *Proc. Natl Acad. Sci. USA* **118**, e2109380118 (2021).

- Deco, G. et al. Whole-brain multimodal neuroimaging model using serotonin receptor maps explains non-linear functional effects of LSD. *Curr. Biol.* **28**, 3065–3074.e6 (2018).
- Burt, J. B. et al. Transcriptomics-informed large-scale cortical model captures topography of pharmacological neuroimaging effects of LSD. *eLife* **10**, e69320 (2021).
- Demirtaş, M. et al. Hierarchical heterogeneity across human cortex shapes large-scale neural dynamics. *Neuron* **101**, 1181–1194.e13 (2019).
- Deco, G. et al. Dynamical consequences of regional heterogeneity in the brain's transcriptional landscape. *Sci. Adv.* **7**, eabf4752 (2021).
- Joglekar, M. R., Mejias, J. F., Yang, G. R. & Wang, X. J. Inter-areal balanced amplification enhances signal propagation in a large-scale circuit model of the primate cortex. *Neuron* **98**, 222–234.e8 (2018).

Acknowledgements

A.I.L. was supported by a Gates Cambridge Scholarship (OPP 1144), the E. G. Fearnside's Travel Fund and a Travel Grant from the Boehringer Ingelheim Fonds. J.C. is funded by the Portuguese Foundation for Science and Technology grants UIDB/50026/2020, UIDP/50026/2020 and CEECIND/03325/2017, Portugal. R.C. and A.D. are supported by the European Union's Horizon 2020 Framework Programme for Research and Innovation under the Specific Grant Agreement No. 945539 (Human Brain Project SGA3). G.D. is supported by HBP SGA3 Human Brain Project Specific Grant Agreement 3 (945539), funded by the EU H2020 FET Flagship program; the Spanish national research project (PID2019-105772GB-I00 MCIU AEI), funded by the Spanish Ministry of Science, Innovation and Universities (MCIU), State Research Agency (AEI); SGR Research Support Group (reference 2017 SGR 1545), funded by the Catalan Agency for Management of University and Research Grants (AGAUR); Neurotwin Digital twins for model-driven non-invasive electrical brain stimulation (grant agreement 101017716), funded by the EU H2020 FET Proactive program; euSNN (grant agreement 860563), funded by the EU H2020 MSCA-ITN Innovative Training Networks; The Emerging Human Brain Cluster (CECH) (001-P-001682) within the framework of the European Research Development Fund Operational Program of Catalonia 2014–2020; Brain-Connects: Brain Connectivity during Stroke Recovery and Rehabilitation (201725.33), funded by the Fundació La Marató TV3; and Corticity, FLAG-ERA JTC 2017 (reference PCI2018-092891), funded by the MCIU, AEI. M.L.K. is supported by the Center for Music in the Brain, funded by the Danish National Research Foundation (DNR117), and the Centre for Eudaimonia and Human Flourishing, funded by the Pettit Foundation and Carlsberg Foundation.

Competing interests

The authors declare no competing interests.