Author Correction: TP53 loss creates therapeutic vulnerability in colorectal cancer

<https://doi.org/10.1038/s41586-021-03664-3>

Published online: 25 August 2021

Correction to: *Nature* <https://doi.org/10.1038/nature14418>

Published online 22 April 2015

Check for updates

Yunhua Liu, Xinna Zhang, Cecil Han, Guohui Wan, Xingxu Huang, Cristina Ivan, Dahai Jiang, Cristian Rodriguez-Aguayo, Gabriel Lopez-Berestein, Pulivarthi H. Rao, Dipen M. Maru, Andreas Pahl, Xiaoming He, Anil K. Sood, Lee M. Ellis, Jan Anderl & Xiongbin Lu

In Extended Data Fig. 9a of this Article, the casp-3 immunohistochemical staining images in the HEA125 and Ama-3 treatment groups of POLR2A^{loss} tumours were inadvertently taken from serial tissue sections of a tumour in the Ama-3 treatment group of POLR2A^{neutral} tumours. In Fig. 1 of this Amendment, we have replaced these two images with corrected ones (highlighted by red boxes). All of the original casp-3 staining images from the POLR2A^{neutral} and POLR2A^{loss} tumours with indicated treatments are shown in Supplementary Information to this Amendment. The immunohistochemistry staining images were not used for quantitative analysis in the study, and this mistake does not affect the conclusions of the original Article. The original Article has not been corrected online.

Supplementary information is available in the online version of this Amendment.

© The Author(s), under exclusive licence to Springer Nature Limited 2021

Original Extended Data Figure 9a

Corrected Extended Data Figure 9a

Fig. 1 | This is the original and corrected Extended Data Fig. 9a of the original Article. The affected panels are highlighted by red boxes.