








Publisher Correction: BRD4 orchestrates genome folding to promote neural crest differentiation

Ricardo Linares-Saldana, Wonho Kim, Nikhita A. Bolar, Haoyue Zhang, Bailey A. Koch-Bojalad, Sora Yoon, Parisha P. Shah, Ashley Karnay, Daniel S. Park , Jennifer M. Luppino , Son C. Nguyen , Arun Padmanabhan, Cheryl L. Smith, Andrey Poleshko , Qiaohong Wang, Li Li, Deepak Srivastava , Golnaz Vahedi, Gwang Hyeon Eom, Gerd A. Blobel , Eric F. Joyce and Rajan Jain 

Correction to: *Nature Genetics* <https://doi.org/10.1038/s41588-021-00934-8>, published online 5 October 2021.

In the version of this Article initially published, there were errors in Fig. 8. In the schematic for BRD4 depletion at right, the downward blue arrow representing stabilization of NIPBL on chromatin should have been comparatively thinner than the upward arrow corresponding to NIPBL dissociation from chromatin. Similarly, the green arrow at bottom right indicating differentiation should have been comparatively thinner to indicate less differentiation to smooth muscle cells as compared to wild-type neural crest progenitors at bottom left.

The errors have been corrected in the online version of the article.

Published online: 1 November 2021

<https://doi.org/10.1038/s41588-021-00980-2>

© The Author(s), under exclusive licence to Springer Nature America, Inc. 2021