after vaccination will be assessed. This trial will reveal whether chemotherapy, immunotherapy, or chemo-immunotherapy influences how patients respond to vaccination and could serve as a model for translational studies of other vulnerable populations or comparable cohorts vaccinated with different vaccines against COVID-19.

Understanding whether this group of patients can mount a sufficient immune response to a vaccine against COVID-19 will provide information for supporting and counseling them during this pandemic and will give them a voice. Is the antibody titer high enough to be protective against COVID-19? What is the durability of antibody titers? Do these patients mount a T cell response sufficient to support the formation of memory B cells? Are two vaccinations sufficient, or are additional doses required? Are alternative measures required?

To assure rapid dissemination of knowledge, we aim to make the results publicly available as soon as possible. Moreover, data sharing will allow comparison between VOICE results and those from other studies for rapid amplification of the knowledge gained. □

Astrid A. M. van der Veldt^{1,15},
Sjoukje F. Oosting^{2,15},
Anne-Marie C. Dingemans³,
Rudolf S. N. Fehrmann²,
Corine GeurtsvanKessel⁴,
Mathilde Jalving², Guus F. Rimmelzwaan⁵,
Pia Kvistborg⁶, Christian U. Blank⁷,
Egbert F. Smit⁸, Valery E. E. P. Lemmens^{9,10},
T. Jeroen N. Hiltermann¹¹,
Marion P. G. Koopmans^{12,4},
Anke L. W. Huckriede¹², Nynke Y. Rots¹³,
Cecile A. C. M. van Els^{13,14}, Debbie van Baarle^{12,13},
John B. A. G. Haanen^{12,16} and
Elisabeth G. E. de Vries^{12,16} ✉

¹Department of Medical Oncology and Department of Radiology & Nuclear Medicine, Erasmus Medical Center-Cancer Institute, Rotterdam, the Netherlands. ²Department of Medical Oncology,

University Medical Center Groningen, University of Groningen, Groningen, the Netherlands.

³Department of Respiratory Medicine, Erasmus Medical Center-Cancer Institute, Erasmus Medical Center, Rotterdam, the Netherlands. ⁴Department of Viroscience, Erasmus Medical Center, Rotterdam, the Netherlands. ⁵Research Center for Emerging Infections and Zoonoses, University of Veterinary Medicine Hannover, Hannover, Germany. ⁶Department of Molecular Oncology and Immunology, Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital, Amsterdam, the Netherlands. ⁷Department of Medical Oncology, Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital, Amsterdam, the Netherlands. ⁸Department of Thoracic Oncology, Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital, Amsterdam, the Netherlands. ⁹Comprehensive Cancer Organization the Netherlands/Netherlands Cancer Registry, Utrecht, the Netherlands. ¹⁰Department of Public Health, Erasmus Medical Center, Rotterdam, the Netherlands. ¹¹Department of Pulmonary Diseases, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands. ¹²Department of Medical Microbiology and Infection Prevention, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands. ¹³Center for Infectious Disease Control, National Institute for Public Health and the Environment, Bilthoven, the Netherlands. ¹⁴Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine, Utrecht University, Utrecht, the Netherlands.

¹⁵These authors contributed equally: Astrid A. M. van der Veldt, Sjoukje F. Oosting. ¹⁶These authors jointly supervised this work: John B.A.G. Haanen, Elisabeth G.E. de Vries.

✉e-mail: e.g.e.de.vries@umcg.nl

Published online: 15 February 2021
<https://doi.org/10.1038/s41591-021-01240-w>

References

- Ribas, A. et al. *Cancer Discov.* **11**, 233–236 (2021).
- Au, L. et al. *Cell* **183**, 4–10 (2020).
- de Joode, K. et al. *Eur. J. Cancer* **136**, 132–139 (2020).
- Nederlandse Federatie van Kankerpatiëntenorganisaties. https://nfk.nl/media/1/Downloads/201210-DJE-corona-II_rapportage_finaal.pdf (2020).
- I&O Research. <https://www.io-research.nl/actueel/lagere-bereidheid-tot-vaccinatie> (2020).

- US Food and Drug Administration. <https://www.fda.gov/emergency-preparedness-and-response/coronavirus-disease-2019-covid-19/covid-19-vaccines> (accessed January 2021).
- Garassino, M. C. et al. *European Society for Medical Oncology* <https://www.esmo.org/covid-19-and-cancer/covid-19-vaccination?hit=ehp> (2020).
- American Society of Clinical Oncology. <https://www.asco.org/asco-coronavirus-resources/covid-19-patient-care-information/covid-19-vaccine-patients-cancer> (accessed January 2021).
- Society for Immunotherapy of Cancer. <https://www.sitcancer.org/aboutsitc/press-releases/2020/sitc-statement-sars-cov-2-vaccination-cancer-immunotherapy> (2020).
- Fendler, A. et al. Preprint at <https://www.medrxiv.org/content/10.1101/2020.12.21.20248608v1> (2020).

Competing interests

A.A.M.v.d.V. reports consultancy fees (paid to Erasmus Medical Center) from BMS, MSD, Merck, Sanofi, Eisai, Pfizer, Ipsen, Roche, Pierre Fabre and Novartis. S.F.O. reports research grants from Novartis and Celldex Therapeutics (paid to University Medical Center Groningen). A.-M.C.D. reports consultancy fees (paid to Erasmus Medical Center) from Roche, Eli Lilly, Boehringer Ingelheim, BMS, Amgen, Novartis, Pfizer, Takeda, Chiesi, Pharmamar, Bayer and Sanofi, and research support (paid to Erasmus Medical Center) from BMS, AbbVie and Amgen. Mathilde Jalving reports consultancy fees (paid to University Medical Center Groningen) from BMS, MSD, Merck, Pfizer, AstraZeneca, Pierre Fabre and Novartis. C.U.B. reports an advisory role at BMS, MSD, Roche, Novartis, GSK, AZ, Pfizer, Lilly, GenMab, Pierre Fabre and Third Rock Ventures, research funding from BMS, Novartis and NanoString, and stock ownership in Uniti Cars, and is a co-founder of Immagine BV. E.F.S. reports consultancy fees (all Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital) from AstraZeneca, Bayer, Bristol Myers Squibb, Daiichi Sankyo, Eli Lilly, Merck, MSD, Novartis, Pfizer, Roche Diagnostics, Roche Genentech and Takeda. T.J.N.H. reports consultancy fees (paid to University Medical Center Groningen) from BMS, MSD, Merck, Boehringer, AstraZeneca and Roche. J.B.A.G.H. reports consultancy roles for Achilles Therapeutics, BioNTech, BMS, GSK, Immunocore, Molecular Partners, MSD, Merck Serono, Neogene Therapeutics, Novartis, Pfizer, Roche/Genentech, Sanofi and Third Rock Ventures, and research grants from Amgen, BMS, BioNTech, MSD, Novartis, Neogene Therapeutics (paid to the Netherlands Cancer Institute–Antoni van Leeuwenhoek Hospital). Elisabeth G.E. de Vries reports an advisory role at Daiichi Sankyo, NSABP and Sanofi (paid to University Medical Center Groningen), and research funding from Amgen, AstraZeneca, Bayer, Chugai Pharma, CytomX Therapeutics, G1 Therapeutics, Genentech, Nordic Nanovector, Radius Health, Regeneron, Roche, Servier and Synthon (paid to University Medical Center Groningen).

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41591-021-01240-w>.

Placebo use and unblinding in COVID-19 vaccine trials: recommendations of a WHO Expert Working Group

To the Editor—The grave public-health threat posed by coronavirus disease 2019 (COVID-19) has spurred an unprecedented

accelerated approach to vaccine research and deployment. Currently, hundreds of thousands of participants globally

are enrolled in COVID-19 vaccine trial research, with more trials imminent or proposed¹. As of 1 March 2021, multiple

regulatory agencies^{3–4} and the World Health Organization (WHO)^{5,6} had granted emergency use designation (EUD) to qualifying candidate vaccines against COVID-19, on the basis of early evidence of efficacy and safety. Consequently, qualifying candidate vaccines are being deployed before conclusion of their trials and/or the collection of longer-term data on safety and efficacy.

From a scientific standpoint, it is important for the clinical trials of vaccines granted an EUD to continue to their completion. But the provision of a candidate vaccine under an EUD to millions of people raises urgent questions about the continuation of the control-group arm of these and other trials, and whether trial blinding is still warranted. Current research-ethics guidance documents^{7–9} were not drafted with emergency-use deployment in mind. Given such dilemmas and guidance gaps, the WHO ACT-Accelerator Ethics & Governance Working Group (ACT: Access to COVID-19 Tools; called 'Working Group' here) has developed a policy brief to guide ethical decision-making in these circumstances: 'Emergency Use Designation of COVID-19 candidate vaccines: Ethical considerations for current and future COVID-19 placebo-controlled vaccine trials and trial unblinding'¹⁰.

The Working Group has concluded that although there is a scientific imperative to continue trials of vaccines against COVID-19 after a candidate vaccine is granted an EUD, there is also an ethical imperative to ensure that trial participants who are at substantial risk of infection with the coronavirus SARS-CoV-2, and severe COVID-19 morbidity or mortality—such as healthcare workers at high to very high risk of acquiring and transmitting the disease, and people above 65 years of age—are in a position to access an EUD vaccine as soon as practically possible, should they wish to do so. Candidate vaccines granted an EUD will probably be deployed in a phased manner to ensure the prioritization of those deemed to be at considerable risk. In settings in which candidate vaccines are introduced under an EUD, investigators should explain the scientific benefit of continued trial participation and the implications of unblinding to trial participants deemed to be at substantial risk of infection, severe morbidity or mortality. Participants should then be offered the opportunity to be unblinded, so that they can make an informed decision about whether to withdraw from the trial and access an EUD vaccine

programmatically as soon as practically possible, should they wish to do so. Trial participants who are not deemed to be at substantial risk of SARS-CoV-2 infection and COVID-19 morbidity or mortality and who do not meet prevailing eligibility criteria to access a candidate vaccine granted an EUD should be informed of the scientific benefits of continuing with the trial and should be encouraged to remain enrolled, with full acknowledgment of their right to withdraw from a trial at any point, without penalty. The continued enrollment of as many participants as possible, for as long as possible, will have considerable scientific and public health utility, as doing so will yield invaluable longer-term data on the safety and efficacy of candidate vaccines. Continued enrollment could also highlight the potential risk of emerging variants of SARS-CoV-2 of concern in relation to those previously unexposed to the virus, those previously infected and recent vaccinees. Such information will be crucial for regulatory decision-making about product registration and/or licensure.

The severe threat to public health posed by the COVID-19 pandemic requires sensitive balancing of the interests of participants in trials of vaccines against COVID-19 versus the need to conduct valuable and urgently needed research into vaccines against COVID-19, in the interest of public health. The policy brief developed by Working Group attempts to strike this delicate balance. □

Jerome Amir Singh^{1,2}✉, Sonali Kochhar^{3,4}, Jonathan Wolff⁵ and The WHO ACT-Accelerator Ethics & Governance Working Group*

¹Howard College School of Law, University of KwaZulu-Natal, Durban, South Africa. ²Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada. ³Department of Global Health, University of Washington, Seattle, WA, USA. ⁴Global Healthcare Consulting, Delhi, India. ⁵Blavatnik School of Government, University of Oxford, Oxford, UK. *A list of authors and their affiliations appears at the end of the paper. ✉e-mail: singhj9@ukzn.ac.za

Published online: 16 March 2021
<https://doi.org/10.1038/s41591-021-01299-5>

References

- World Health Organization. <https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines> (accessed 2 March 2021).
- Medicines and Healthcare products Regulatory Agency. <https://www.gov.uk/government/news/uk-medicines-regulator-gives-approval-for-first-uk-covid-19-vaccine> (2 December 2020).

- U.S. Food & Drug Administration. <https://www.fda.gov/news-events/press-announcements/fda-takes-key-action-fight-against-covid-19-issuing-emergency-use-authorization-first-covid-19> (11 December 2020).
- European Medicines Agency. <https://www.ema.europa.eu/en/news/ema-recommends-first-covid-19-vaccine-authorisation-eu> (21 December 2020).
- World Health Organization. <https://www.who.int/news/item/31-12-2020-who-issues-its-first-emergency-use-validation-for-a-covid-19-vaccine-and-emphasizes-need-for-equitable-global-access> (31 December 2020).
- World Health Organization. <https://www.who.int/news/item/15-02-2021-who-lists-two-additional-covid-19-vaccines-for-emergency-use-and-covax-roll-out#:~:text=Today%20WHO%20listed%20two%20versions,the%20Serum%20Institute%20of%20India> (15 February 2021).
- World Medical Association. <https://www.wma.net/policies-post/wma-declaration-of-helsinki-ethical-principles-for-medical-research-involving-human-subjects/> (2013).
- Council for International Organizations of Medical Sciences. <https://cioms.ch/publications/product/international-ethical-guidelines-for-health-related-research-involving-humans/> (2016).
- Rid, A., Saxena, A., Baqui, A. H. & Bhan, A. et al. *Vaccine* **32**, 4708–4712 (2014).
- World Health Organization. https://apps.who.int/iris/bitstream/handle/10665/337940/WHO-2019-nCoV-Policy_Brief-EUD_placebo-controlled_vaccine_trials-2020.1-eng.pdf?sequence=1&isAllowed=y (18 December 2020).

Acknowledgements

The policy brief reported in this work was supported by the grant WHO COVID-19 SPRP R&D, funded by the Ministry of Health of Germany. K. Littler, A. Reis (WHO Health Ethics & Governance Unit), J. Hombach and A. Wilder-Smith (WHO Strategic Advisory Group of Experts on Immunization Secretariat) provided support for the WHO Secretariat. Contributions from G. Enwere and C. Rodriguez (WHO Prequalification/Vaccines & Immunization Devices Assessment Unit), M.-C. Bouésseau (WHO Department of Integrated Health Services) and P. Hummel (WHO Consultant) are acknowledged.

Competing interests

The authors declare no competing interests.

The WHO ACT-Accelerator Ethics & Governance Working Group

Caesar Atuire⁶, Anant Bhan⁷, Ezekiel Emanuel⁸, Ruth Faden⁹, Prakash Ghimire¹⁰, Dirceu Greco¹¹, Calvin Ho¹², Suerie Moon¹³, Ehsan Shamsi-Gooshki¹⁴, Aissatou Touré¹⁵, Beatriz Thomé¹⁶, Maxwell J. Smith¹⁷ and Ross E. G. Upshur¹⁸

⁶University of Ghana, Accra, Ghana. ⁷Yenepoya University, Mangalore, India. ⁸University of Pennsylvania, Philadelphia, PA, USA.

⁹Johns Hopkins University, Baltimore, MD, USA. ¹⁰Tribhuvan University, Kirtipur, Nepal.

¹¹Federal University of Minas Gerais, Belo Horizonte, Brazil. ¹²University of Hong Kong, Hong Kong, China. ¹³Graduate Institute, Geneva, Switzerland. ¹⁴Tehran University of Medical Sciences, Tehran, Iran. ¹⁵Institut Pasteur, Dakar, Senegal.

¹⁶Federal University of Sao Paulo, Sao Paulo, Brazil.

¹⁷Western University, London, Ontario, Canada.

¹⁸University of Toronto, Toronto, Ontario, Canada.