the 'prevention, containment or treatment' of COVID-19, when India and South Africa proposed that the World Trade Organization temporarily suspend Trade Related Aspects of Intellectual Property Rights rules⁷.

Already in 2019, a Lancet Commission recommended increased utilization of legal mechanisms to advance global health through justice systems⁸ and to harness the power of law to promote global health and sustainable development, calling upon governments, the WHO and United Nations organizations to make necessary legal reforms to translate vision into action.

The current crisis is a salient opportunity for strengthening the global response to upcoming pandemics. With the belief that vaccines against COVID-19 are a global common good and that allocation should be based on need, we make the following recommendations.

First, a global legal framework is needed to regulate the commercial determinants of health and to secure universal fair access to essential technologies and vaccine production. This may be financed through existing alliances (such as COVAX, or Gavi, the Vaccine Alliance) or by new G7 or G20 standing funds for vaccine development and allocation⁹. In times of emergency, the World Trade Organization should also grant temporary waivers for Trade Related Aspects of Intellectual Property Rights rules for governments to produce their own vaccines or, if necessary, to exercise compulsory licensing powers for a third party to produce a patented technology. Ideally, perceived legal risk could facilitate the negotiation of purchase transparency, voluntary licensing arrangements and patent sharing while committing the industry to global justice.

Second, to strengthen crisis preparedness⁴ and make the response less blunt and more evidence-based, coordinated and effective, the existing gaps in infrastructure, technology and information must be reduced. The WHO warns that lockdown-induced disruption (e.g., of childhood immunization and essential care) could cause greater death and suffering than COVID-19 itself¹⁰. As LMICs embark on an unprecedented vaccination scale-up, substantial investment is needed to prevent redistribution of scarce health personnel that further undermines routine immunization. Future plans must have a clear health-systems dimension and must include social mobilization and strategies for fighting vaccine misinformation⁷.

Third, as shown by COVID-19, updated global governance mechanisms are needed that better reflect the contemporary geopolitical order and truly encourage international collaboration across sectors, through political and legal solutions rooted in commitments to justice and shared global responsibility^{5,8}. We call for a 21st-century Bretton Woods-type conference, similar to that of 1944, when states met to improve global economic interaction, secure peace and provide assistance to countries devastated by World War II, which resulted in new global governance structures such as the International Monetary Fund and the World Bank.

Using the dynamics of one acute crisis to build resilience against the next could also serve a greater good in line with the growing notion of the United Nations Security Council that health, poverty mitigation, security and peace are interlinked.

Anna Mia Ekström¹✉, Camilla Berggren², Göran Tomson^{2,3}, Lawrence O. Gostin⁴, Peter Friberg^{2,5} and Ole Petter Ottersen³

¹Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden. ²Swedish Institute for Global Health Transformation, Royal Swedish Academy of Sciences, Stockholm, Sweden. ³President's Office, Karolinska Institutet, Stockholm, Sweden.

⁴O'Neill Institute for National and Global Health Law, Georgetown University, Washington, DC, USA.

⁵Department of Public Health and Community medicine, Sahlgrenska Academy, Gothenburg University, Gothenburg, Sweden.

✉e-mail: anna.mia.ekstrom@ki.se

Published online: 11 March 2021

<https://doi.org/10.1038/s41591-021-01288-8>

References

1. *Nat. Med.* **27**, 1 (2021).
2. Launch & Scale Speedometer. <https://launchandscalefaster.org/COVID-19> (2021).
3. Emanuel, E. J. et al. *Science* **369**, 1309–1312 (2020).
4. Ottersen, O. P. & Engebretsen, E. *Nat. Med.* **26**, 1672–1673 (2020).
5. Ottersen, O. P. et al. *Lancet* **383**, 630–667 (2014).
6. COVAX. Gavi, the Vaccine Alliance https://www.gavi.org/sites/default/files/covid/covax/COVAX_Principles-COVID-19-Vaccine-Doses-COVAX.pdf (2020).
7. Deutsche Welle. <https://www.dw.com/en/rich-countries-block-india-south-africa-bid-to-ban-covid-vaccine-patents/a-56460175> (2021).
8. Gostin, L. O. et al. *Lancet* **393**, 1857–1910 (2019).
9. Gostin, L. O., Friedman, E. A. & Moon, S. *Foreign Affairs* <https://www.foreignaffairs.com/articles/world/2021-01-19/wealthy-countries-should-share-vaccine-doses-it-too-late> (2021).
10. World Health Organization. <https://www.who.int/news/item/15-07-2020-who-and-unicef-warn-of-a-decline-in-vaccinations-during-covid-19> (2020).

Author contributions

All authors contributed to the writing of this manuscript; A.M.E. drafted the first version and edited the final version on the basis of feedback from all co-authors; and all authors have seen and approved the submitted version.

Competing interests

The authors declare no competing interests.

Comprehensive large-scale nucleic acid-testing strategies support China's sustained containment of COVID-19

To the Editor—Since identification and initial characterization of COVID-19 in late December 2019 and early January 2020, China's containment goal has been complete interruption of local transmission of the coronavirus SARS-CoV-2¹. The nationwide epidemic wave centered on Wuhan was contained by April 2020 with non-pharmaceutical interventions alone but

was associated with economic loss: China's first-quarter gross domestic product fell 6.8% compared with that in the first quarter of 2019 (ref. ²). Since containment, there have been numerous outbreaks in China caused by imported viruses with local spread, varying in size from several cases to about 1,000 cases, all stopped with non-pharmaceutical interventions alone³. Sustained containment

has high socioeconomic value: economic activities recovered rapidly, with an increase in gross domestic product of 3.2% during April through June 2020, and of 4.9% during July through September 2020 (ref. ²); the government's response to the pandemic was perceived positively by the Chinese public⁴. In this Correspondence, we describe two large-scale PCR-based testing strategies that

Fig. 1 | Local and imported COVID-19 cases from 1 April 2020 to 30 November 2020. Light blue wide bars indicate outbreaks (key) and adjacent labels indicate mass PCR testing strategies.

have played critically important roles in sustaining containment since April 2020 in mainland China: routine, active, PCR-based screening, and mass PCR testing during the outbreak response.

With almost all of China's population being susceptible to SARS-CoV-2 before the induction of any vaccine-induced population immunity⁵, imported viruses posed an enormous threat to containment. All incoming international travelers, workers handling imported goods, medical-center outpatients with fever or respiratory symptoms, and medical staff caring for patients with fever or infectious diseases are actively screened with PCR to determine if they are infected. Incoming travelers have been required to quarantine at designated hotels for 14 days after entry into China, during which time they are tested at least twice. As of 30 November 2020, 3,866 SARS-CoV-2-infected travelers had been diagnosed, which prevented seeding of the virus in susceptible communities (Fig. 1).

Outbreak investigations in Beijing and Dalian found that food contamination could potentially transmit SARS-CoV-2^{6,7}. Isolation of viable virus in the outer packages of imported seafood in Qingdao strengthened the evidence for such transmission. Workers handling imported goods are now required to be tested weekly for SARS-CoV-2, which has prevented several importation-related outbreaks—for example, four dockworkers were found to have become infected in December 2020 in Dalian, and an outbreak investigation in Kashgar, Xinjiang, found importation of SARS-CoV-2 via a contaminated cross-border truck.

Fever clinics have long been established in secondary and tertiary hospitals. Anyone with fever seeking healthcare is screened by PCR. Medical facilities admit anyone with suspected COVID-19, unexplained pneumonia, or severe, acute respiratory-tract infection, and they test the patients and accompanying people by

PCR. This screening strategy has identified index cases of several outbreaks: the Xinfadi market in Beijing; Dalian; Pudong airport⁸; Qingdao⁹; and Manchuria.

Routine testing requires many PCR tests. Laboratories combine five or ten specimens and test the combination for SARS-CoV-2 RNA as an efficiency and cost-saving measure. If the combination tests have a positive result, people whose samples were combined are tested separately to identify the infected person (or people). With a 10:1 pooled sampling approach, the average cost per person tested is approximately 9 RMB yuan (US\$ 1.50) in China.

Routine PCR testing is augmented with testing of all close contacts of an infected person. Since September 2020, all close contacts of infected people have been required to be tested three times during a 14-day centralized quarantine period: at day 1 or 2; 1 day later; and at day 14.

Mass, community-wide PCR testing during or after an outbreak response has

proven to be a useful strategy for identifying infected people, including those with no symptoms, and for determining whether a community is free of SARS-CoV-2. Once a local outbreak is identified, geographic jurisdictions are classified by epidemiologists into regions at low, moderate or high risk of transmitting the virus. People in high-risk settings are tested individually. In moderate-risk areas, a 5:1 or 10:1 pooled sampling approach is used to reduce the number of tests.

The first city-wide mass screening was performed between 14 May and 1 June 2020 in Wuhan city; nearly 10 million people were tested by PCR, and 300 asymptotically infected people were identified for quarantine. The testing results were used to support full reopening of local socioeconomic activities¹⁰. Large screening programs have been implemented following local case reports that indicated possible community transmission. For instance, during the outbreak in Beijing's Xinfadi market, 11.9 million people were tested between 11 June and 14 July 2020; in Dalian, 4.5 million people were tested between 26 July and 31 July 2020; and in Qingdao, 10.9 million people were tested between 11 October and 16 October 2020 (ref. ⁹). These risk-based, large-scale, population-wide screenings contributed to case finding and rapid control of epidemics and provided evidence for the government to reopen economic activities.

The timeliness of PCR test results varies by testing purpose and epidemiological situation. For members of the public who request PCR testing but have no symptoms or exposure history, results are to be reported within 24 hours. Results

for outpatients, inpatients, caregivers, close contacts of confirmed cases and people residing in high-risk areas in a local outbreak setting are to be reported within 12 hours. Results for patients in fever clinics and emergency departments are to be reported within 6 hours.

In China's 'new-normal' situation, comprehensive, active and innovative PCR testing strategies are performed for targeted groups and in outbreak settings; this ensures timeliness of early case detection and interruption of local outbreaks—essential ingredients in the prevention and control strategy. Maintenance of containment has greatly reduced the impact of COVID-19 in terms of suffering, lives lost and socioeconomic progress.

The introduction of vaccines against COVID-19 globally and in China will undoubtedly improve epidemiological situations. PCR testing strategies will be adjusted to fit the changing epidemiological situation in China—probably one in which few or fewer non-pharmaceutical interventions will be needed for effective epidemic control. Inexpensive, readily available and rapid PCR testing will continue to be essential for sensitive surveillance of SARS-CoV-2 well into the future. Throughout 2020, PCR testing served the public well, helping to make and keep China nearly free of SARS-CoV-2, and providing socioeconomic space and time for vaccine development and long-term prevention and control of COVID-19. □

Zhongjie Li^{1,5}, Fengfeng Liu^{1,5}, Jinzhao Cui¹, Zhibin Peng¹, Zhaorui Chang¹, Shengjie Lai^{1,2}, Qiulan Chen¹, Liping Wang¹, George F. Gao^{3,4,6} and Zijian Feng^{3,6}✉

¹Division of Infectious disease, Key Laboratory

of Surveillance and Early Warning on Infectious Disease, Chinese Center for Disease Control and Prevention, Beijing, China. ²WorldPop, School of Geography and Environmental Science, University of Southampton, Southampton, UK. ³Chinese Center for Disease Control and Prevention, Beijing, China. ⁴CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China. ⁵These authors contributed equally: Zhongjie Li, Fengfeng Liu. ⁶These authors jointly supervised this work: George F. Gao and Zijian Feng. ✉e-mail: fengzj@chinacdc.cn

Published online: 15 April 2021
<https://doi.org/10.1038/s41591-021-01308-7>

References

- Li, Z. et al. *Lancet* **396**, 63–70 (2020).
- National Bureau of Statistics of China. http://www.stats.gov.cn/tjsj/zxfb/202010/t20201020_1794939.html (accessed 16 February 2021).
- Zheng, C. et al. *China CDC Weekly* **2**, 902–905 (2020).
- Lazarus, J. V. et al. *PLoS One* **15**, e0240011 (2020).
- Li, Z. et al. *Lancet Reg. Health West. Pac.* **8**, 100094 (2021).
- Pang, X. et al. *Natl. Sci. Rev.* **7**, 1861–1864 (2020).
- Zhao, X. et al. *China CDC Weekly* **2**, 658–660 (2020).
- Fang, F., Song, Y., Hao, L., Nie, K. & Sun, X. *China CDC Weekly* **47**, 910–911 (2020).
- Xing, Y., Wong, G. W. K., Ni, W. & Hu, X. *Engl. J. Med.* **383**, e129–e129 (2020).
- Cao, S. et al. *Nat. Commun.* **11**, 5917–5917 (2020).

Acknowledgements

We thank L. Rodewald for contributions to the content and improvement to the text of this Correspondence.

Author contributions

Z.L., Z.F. and G.F.G. designed and supervised the study; Z.L. and F.L. wrote the manuscript; J.C. designed and drew the figure; Z.P., Z.C., Q.C. and L.W. participated in literature review and data collection; and S.L. and G.F.G. commented on and revised drafts of the manuscript.

Competing interests

The authors declare no competing interests.

Open letter to international funders of science and development in Africa

To the Editor—Recently there was an announcement¹ of a US\$30 million grant awarded to the nonprofit health organization PATH by the US government's President's Malaria Initiative (PMI). The grant funded a consortium of seven institutions in the USA, the UK and Australia to support African countries in the improved use of data

for decision-making in malaria control and elimination.

Not one African institution was named in the press release. The past year has been full of calls from staff and collaborators of various public-health entities for equality and inclusion, so one might imagine that such a partnership to support Africa should

be led from Africa by African scientists, partnering with Western institutions where appropriate, especially where capacity has been demonstrated.

We write this letter to the major international funders of science and development in Africa as African scientists, policy analysts, public-health practitioners