

SCIENTIFIC REPORTS



OPEN

Author Correction:

Periodontal pathogenic bacteria, *Aggregatibacter actinomycetemcomitans* affect non-alcoholic fatty liver disease by altering gut microbiota and glucose metabolism

Rina Komazaki¹, Sayaka Katagiri¹, Hirokazu Takahashi², Shogo Maekawa¹, Takahiko Shiba¹, Yasuo Takeuchi¹, Yoichiro Kitajima^{2,3}, Anri Ohtsu¹, Sayuri Udagawa¹, Naoki Sasaki¹, Kazuki Watanabe¹, Noriko Sato⁴, Naoyuki Miyasaka⁵, Yuichiro Eguchi⁶, Keizo Anzai² & Yuichi Izumi¹

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-017-14260-9>, published online 24 October 2017

This Article contains errors in Table 1, where incorrect values for ‘Fasting plasma glucose (mg/dL)’ and ‘L/S ratio’ are given. The correct Table 1 appears below.

In addition, this Article contains a typographical error in the Methods section, under the subheading ‘Statistical analysis’ where,

“Bonferroni connection”

should read:

“Bonferroni correction”

¹Department of Periodontology, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University (TMDU), Tokyo, Japan. ²Division of Metabolism and Endocrinology, Faculty of Medicine, Saga University, Saga, Japan. ³Eguchi Hospital, Ogi, Saga, Japan. ⁴Department of Molecular Epidemiology, Medical Research Institute, Tokyo Medical and Dental University (TMDU), Tokyo, Japan. ⁵Department of Comprehensive Reproductive Medicine, Tokyo Medical and Dental University (TMDU), Tokyo, Japan. ⁶Liver Center, Saga University Hospital, Saga, Japan. Correspondence and requests for materials should be addressed to S.K. (email: katagiri.peri@tmd.ac.jp) or H.T. (email: takahas2@cc.saga-u.ac.jp)

Variables	All patients	Males	Females
	(n = 52)	(n = 27)	(n = 25)
Age (years)	55 ± 13.8	48 ± 14.6	63 ± 7.5
Weight (kg)	68 ± 12.0	75 ± 11.0	59 ± 6.2
BMI (kg/m ²)	26 ± 2.8	27 ± 3.2	26 ± 2.2
Blood platelet count (× 10 ⁹ /μL)	22 ± 5.5	22 ± 5.3	22 ± 4.7
AST (IU/L)	26 ± 9.1	28 ± 9.5	23 ± 8.1
ALT (IU/L)	30 ± 18.2	37 ± 20.0	22 ± 12.2
ALP (IU/L)	243 ± 69.8	206 ± 63.2	284 ± 52.1
γ-GTP (IU/L)	57 ± 74.4	80 ± 95.8	31 ± 18.6
Fasting plasma glucose (mg/dL)	105 ± 14	106 ± 16	104 ± 11
HOMA-IR	2.1 ± 1.5	1.8 ± 1.1	2.4 ± 1.9
Total cholesterol (mg/dL)	219 ± 48	227 ± 46	218 ± 25
High density lipoprotein (mg/dL)	57 ± 14	54 ± 14	62 ± 12
Triglyceride (mg/dL)	145 ± 100	165 ± 128	124 ± 58
L/S ratio	1.12 ± 0.21	1.07 ± 0.26	1.17 ± 0.15

Table 1. Characteristics of the patients with NAFLD. Data are shown as mean ± SD.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018