

SCIENTIFIC REPORTS



OPEN

Lobohedleolide suppresses hepatitis C virus replication via JNK/c-Jun-C/EBP-mediated down-regulation of cyclooxygenase-2 expression

Chun-Kuang Lin^{1,2}, Chin-Kai Tseng^{3,4}, Chih-Chuang Liaw^{1,5}, Chiung-Yao Huang⁵, Chih-Ku Wei⁶, Jyh-Horng Sheu^{1,5,7} & Jin-Ching Lee^{6,8,9,10}

Hepatitis C virus (HCV) chronically infects 2–3% people of the global population, which leads to liver cirrhosis and hepatocellular carcinoma. Drug resistance remains a serious problem that limits the effectiveness of US Food and Drug Administration (FDA)-approved direct-acting antiviral (DAA) drugs against HCV proteins. The objective of our study was to discover new antivirals from natural products to supplement current therapeutics. We demonstrated that lobohedleolide, isolated from the Formosan soft coral *Lobophytum crassum*, significantly reduced HCV replication in replicon cells and JFH-1 infection system, with EC_{50} values of 10 ± 0.56 and $22 \pm 0.75 \mu\text{M}$, respectively, at non-toxic concentrations. We further observed that the inhibitory effect of lobohedleolide on HCV replication is due to suppression of HCV-induced cyclooxygenase-2 (COX-2) expression. Based on deletion-mutant analysis of the COX-2 promoter, we identified CCAAT/enhancer-binding protein (C/EBP) as a key transcription factor for the down-regulation of COX-2 by lobohedleolide, through which lobohedleolide decreased the phosphorylation of c-Jun NH2-terminal protein kinase and c-Jun to suppress HCV-induced C/EBP expression. The combination treatment of lobohedleolide with clinically used HCV drugs synergistically reduced HCV RNA replication, indicating that lobohedleolide exhibited a high biomedical potential to be used as a supplementary therapeutic agent to control HCV infection.

Hepatitis C virus (HCV) is a pathogen of high risk causing chronic liver diseases, including hepatic fibrosis, liver cirrhosis and hepatocellular carcinoma (HCC)¹. The current clinical standard therapy for chronic hepatitis C (CHC) infection is a combination treatment with pegylated interferon- α (peg-IFN- α) and ribavirin (RBV), but the sustained virologic response (SVR) with only about 40% for HCV genotype 1 patients. As of 2011, two US Food and Drug Administration (FDA)-approved direct-acting antiviral (DAA) agents, boceprevir and telaprevir, against the viral protease activity were shown to significantly improve the antiviral response rate in the treatment of CHC infection². However, the lack of pangenotypic activity and the rapid selection of drug-resistant viral mutants have limited the antiviral response in patients infected with HCV genotypes 2–6³. More recently, the

¹Doctoral Degree Program in Marine Biotechnology, College of Marine Sciences, National Sun Yat-Sen University, Kaohsiung, Taiwan. ²Doctoral Degree Program in Marine Biotechnology, Academia Sinica, Taipei, Taiwan. ³Institute of Basic Medical Sciences, College of Medicine, National Cheng Kung University, Tainan, Taiwan. ⁴Center of Infectious Disease and Signaling Research, College of Medicine, National Cheng Kung University, Tainan, Taiwan. ⁵Department of Marine Biotechnology and Resources, College of Marine Sciences, National Sun Yat-Sen University, Kaohsiung, Taiwan. ⁶Department of Biotechnology, College of Life Science, Kaohsiung Medical University, Kaohsiung, Taiwan. ⁷Department of Medical Research, China Medical University Hospital, China Medical University, Taichung, Taiwan. ⁸Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan. ⁹Research Center for Natural Products and Drug Development, Kaohsiung Medical University, Kaohsiung, Taiwan. ¹⁰Department of Medical Research, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan. Correspondence and requests for materials should be addressed to J.-H.S. (email: sheu@mail.nsysu.edu.tw) or J.-C.L. (email: jcllee@kmu.edu.tw)

second-generation drugs, such as simeprevir and sofosbuvir, were shown to exhibit an improved resistance profile and resulted in a higher cure rate and fewer side effects in patients treated for CHC infection^{4,5}. However, due to the high cost of these drugs, some countries might face a potential difficulty in accepting the therapy⁶. Therefore, there still remains a need to discover new anti-HCV agents to improve the current therapy.

HCV is an enveloped RNA virus and belongs to the *Flaviviridae* family⁷. The RNA encodes a single poly-protein consisting of approximately 3,000 amino acids which is cleaved into 10 viral proteins by viral and host proteases⁸. In HCV-related liver diseases, cyclooxygenase-2 (COX-2) and its metabolite prostaglandin E₂ (PGE₂), are highly expressed and considered as two of the major cellular effectors causing tissue injury, fibrogenesis, chronic hepatitis, liver cirrhosis, and HCC^{9,10}. Several reports and our previous studies have demonstrated that inactivation of COX-2 by COX-2 shRNA or a specific inhibitor can impair HCV replication^{11,12}. Therefore, the interruption of HCV-induced COX-2 signaling pathway has been considered as a promising strategy to decrease HCV infection and HCV-induced inflammatory pathogenesis^{12,13}.

COX-2 is a rate-limiting enzyme that converts arachidonic acid into prostanoids¹⁴. The expression of COX-2 is tightly controlled in most of the tissues and is induced at the sites of inflammation by several extracellular and intracellular stimuli, including reactive oxygen species, chemicals, and viral infections. Both nuclear factor- κ B (NF- κ B) and mitogen-activated protein kinase (MAPK) signaling pathways have been well characterized to activate COX-2 transcription¹⁵. The central transcriptional region of COX-2 contains several binding sites of transcriptional factors for its expression, including NF- κ B, CCAAT/enhancer-binding protein (C/EBP), and activator factor-1 (AP-1)/cyclic adenosine monophosphate (cAMP)-response element (CRE)^{16,17}.

Soft corals of the genus *Lobophytum* have been demonstrated to consist of a rich harvest of cembranoids and steroids, which exhibit anti-HIV and anti-inflammatory bioactive properties^{18,19}. Previous studies have shown that lobohedleolide, isolated from the Formosan soft coral *Lobophytum crassum*, exhibited anti-inflammatory activity through the suppression of COX-2 expression *in vitro* and *in vivo*^{19,20}. Therefore, we investigated the effect of lobohedleolide on HCV replication and further clarified whether the molecular mechanisms were down-regulation of HCV-induced COX-2 expression.

Results

Lobohedleolide decreases HCV replication in both HCV replicon cells and the infection system. To investigate the effect of LH on HCV protein synthesis and RNA replication, HCV replicon cells, Ava5, were incubated with lobohedleolide at different concentrations (5–40 μ M) for 3 days. The total cell lysate and cellular RNA were extracted and subjected to Western blotting and qRT-PCR, respectively. The results indicated that lobohedleolide dose-dependently reduced HCV protein and RNA levels, with an EC₅₀ value of $10 \pm 0.56 \mu$ M, compared with those of lobohedleolide -untreated cells (Fig. 1b,c). To rule out the possibility that the inhibitory effect of lobohedleolide on viral replication was due to the cytotoxic effect on the treated cells, the cell viability of lobohedleolide -treated cells was evaluated by MTS assay. As shown in Fig. 1c, no significant cytotoxicity was observed in the lobohedleolide-treated Ava5 cells at effective antiviral concentrations. Furthermore, lobohedleolide time-dependently reduced HCV protein synthesis and RNA replication (Fig. 1d). The anti-HCV activity of lobohedleolide was further evaluated in the HCV JFH-1 infection system, which showed an EC₅₀ value of $22 \pm 0.75 \mu$ M (Fig. 1e).

Lobohedleolide suppresses HCV replication by inhibiting HCV-induced COX-2 expression and its activity. Inhibition of COX-2 expression has been reported to interfere with HCV replication²¹. In the LPS-induced inflammatory model, lobohedleolide has been shown to exert an inhibitory effect on COX-2 expression²⁰. To investigate whether the suppression of COX-2 plays an important role in the activity of lobohedleolide against HCV replication, we first detected the amount of COX-2 protein in the HCV self-replicating cells, Ava5, and HCV JFH-1-infected Huh-7 cells after lobohedleolide treatment. The results of Western blotting indicated that lobohedleolide markedly suppressed the HCV-induced COX-2 protein levels in a dose-dependent manner compared with that in the parental Huh-7 cells, lobohedleolide-untreated Ava5 cells, and HCV-infected Huh-7 cells in the absence of lobohedleolide treatment (Fig. 2a,b). Next, we used a COX-2 promoter-based reporter assay to evaluate the inhibitory effect of lobohedleolide on COX-2 at the transcriptional level. The pCOX-2-Luc, a plasmid encoding firefly luciferase under the control of the COX-2 promoter, was transfected into Ava5 cells or HCV-infected Huh-7 cells, and then, these plasmid-transfected cells were incubated with lobohedleolide at increasing concentrations for 3 days. As shown in Fig. 2c,d, lobohedleolide dose-dependently decreased the HCV-elevated COX-2 promoter activity in Ava5 and HCV-infected cells. We further investigated the effect of lobohedleolide on COX-2 catalytic activity by monitoring the levels of PGE₂ in Huh-7 and Ava5 cells. The levels of HCV-elevated PGE₂ were significantly reduced with lobohedleolide treatment in a dose-dependent manner compared with that in the untreated cells (Fig. 2e).

To characterize whether the lobohedleolide -mediated down-regulation of COX-2 expression and its catalytic activity is involved in the suppression of HCV replication, we overexpressed exogenous COX-2 to evaluate the inhibitory activity of lobohedleolide on HCV replication. Ava5 cells were transfected with vehicle or various concentrations of pCMV-COX-2-Myc vector encoding the *cox-2* gene, and the plasmid-transfected cells were incubated with 40 μ M of lobohedleolide or 0.1% of DMSO, as a negative control. As shown in Fig. 2f, the lobohedleolide-reduced viral RNA level (top panel, lane 2) was gradually rescued following the increasing expression of exogenous COX-2-Myc (top panel, lanes 3–5) compared with that in the vehicle-transfected Ava5 cells incubated with DMSO (top panel, lane 1). Consistently, the exogenous expression of COX-2 attenuated the inhibitory effect of lobohedleolide on the HCV protein levels, as analyzed by Western blotting. These results clearly indicated that lobohedleolide inhibited HCV replication by down-regulating the virus-induced COX-2 expression.




Figure 1. The inhibitory effect of lobohedleolide on HCV replication. **(a)** The chemical structure of lobohedleolide. The reduction effect of lobohedleolide on HCV **(b)** protein synthesis and **(c)** RNA replication. Ava5 cells were treated with lobohedleolide at the indicated concentrations for 3 days. Cell viability was evaluated by MTS assay in the lobohedleolide-treated Ava5 cells. **(d)** Time-dependent inhibition of HCV protein synthesis and RNA replication by lobohedleolide. Ava5 cells were treated with 40 μM of lobohedleolide for the indicated time points. Total cell lysates were subjected to Western blotting with anti-HCV NS5B and anti-GAPDH antibodies. GAPDH served as an equal loading control in Western blotting. **(e)** Concentration-dependent inhibition of lobohedleolide in HCV JFH-1 infection system. Huh7.5 cells were infected with HCV JFH-1 at an MOI of 0.1 for 4 h, and the HCV-infected cells were treated with lobohedleolide at the indicated concentrations for another 3 days. Total cellular RNA was extracted, and the levels of HCV RNA were determined by qRT-PCR following the normalization of cellular *gadh* mRNA level. The efficiency of inhibition was calculated as the percentage of control (0.1% DMSO treatment). Data are presented as mean \pm SD of at least three independent experiments, with each measurement carried out in triplicate. Asterisks indicate significant difference in each sample of lobohedleolide-treated cells at the indicated concentration singly in comparison with DMSO-treated cells. * $P < 0.05$; ** $P < 0.01$.

Lobohedleolide inhibits C/EBP transcription factor activity. COX-2 induction is mediated by multiple transcription factor pathways, such as NF- κ B, C/EBP, and AP-1²². To identify which binding elements of transcription factors on the COX-2 promoter are responsible for modulating the lobohedleolide-mediated inhibition of COX-2 expression, we created a series of COX-2 promoter reporter constructs carrying the deleted promoter fragments, including wild-type (WT), Δ NF- κ B, and Δ NF- κ B/C/EBP, linked to firefly luciferase gene to elucidate the effect of lobohedleolide on COX-2 transcription, in which the WT promoter region contained NF- κ B, C/EBP, and AP-1 responsive elements (Fig. 3a). Huh-7 and Ava5 cells were transiently transfected with pCOX-2-Luc, pCOX-2(Δ NF- κ B)-Luc, or pCOX-2(Δ NF- κ B/C/EBP)-Luc, and the plasmid-transfected cells were respectively treated with lobohedleolide at 0, 20 and 40 μM . The relative COX-2 promoter activities were analyzed using luciferase activity assay. As shown in Fig. 3b, lobohedleolide dose-dependently reduced the HCV-induced luciferase activity driven by the COX-2/WT promoter and the COX-2/ Δ NF- κ B promoter, but exhibited no significant effect on that of luciferase activity driven by the COX-2/ Δ NF- κ B/C/EBP promoter, suggesting that the




Figure 2. The inhibitory effect of lobohedleolide on HCV-induced COX-2 expression at the protein and transcription levels. The inhibition of HCV-induced COX-2 expression by lobohedleolide in (a) HCV replicon cells and (b) HCV infection assay. Ava5 cells and HCV JFH-1-infected Huh7.5 cells were treated with lobohedleolide at the indicated concentrations for 3 days. Cell lysates were subjected to Western blotting with antibodies against COX-2 and GAPDH. GAPDH served as an equal loading control in Western blotting. The reduction of HCV-induced COX-2 promoter activity by lobohedleolide in (c) HCV replicon cells and (d) HCV infection system. Huh-7 cells and Ava5 cells were transfected with pCOX-2-Luc reporter plasmid encoding the firefly luciferase gene under the COX-2 promoter control. The pCOX-2-Luc-transfected Huh-7.5 cells were infected with JFH-1 virus at an MOI of 0.1 for 4 h. The transfected Ava5 cells and JFH-1-infected Huh-7.5 cells were treated with lobohedleolide at the indicated concentrations for 3 days. Cell lysates were subjected to luciferase activity assay. (e) Dose-dependent reduction of HCV-induced PGE₂ production by lobohedleolide. Ava5 cells were incubated with lobohedleolide at the indicated concentrations for 3 days. PGE₂ production was analyzed by PGE₂ ELISA kit. (f) Concentration-dependent restoration of HCV replication by exogenous COX-2 expression in lobohedleolide-treated Ava5 cells. Ava5 cells were transfected with the indicated amount of pCMV-COX-2-Myc encoding *cox-2* (lanes 3, 4, and 5) or pcDNA3.1 (1.5 μg , lanes 1 and 2) for 6 h, and the transfected cells were treated with DMSO (lane 1) or lobohedleolide at 40 μM for 3 days (lanes 2, 3, 4, and 5). Cell lysates were subjected to Western blotting with antibodies against NS5B, Myc, and GAPDH. GAPDH served as an equal loading control in Western blotting. Total cellular RNAs were extracted and subjected to qRT-PCR to evaluate the level of HCV RNA. Data are presented as mean \pm SD of at least three independent experiments, with each measurement carried out in triplicate. Asterisks indicate significant difference in each sample of lobohedleolide-treated cells at the indicated concentration singly in comparison with DMSO-treated cells. * $P < 0.05$; ** $P < 0.01$.




Figure 3. The effect of lobohedleolide on the transcriptional factor activity on COX-2 promoter. (a) Schematic diagrams of the deletion of COX-2 promoter constructs. (b) Effect of lobohedleolide on the deletion constructs of COX-2 promoter. Ava5 cells were cotransfected with 0.2 μ g of pCMV-Rellina-Luc and pCOX-2-Luc, pCOX-2(Δ NF- κ B)-Luc, or pCOX-2(Δ NF- κ B/C/EBP)-Luc for 6 h. The transfected cells were treated with DMSO or lobohedleolide at 20 and 40 μ M for 3 days. Cell lysates were subjected to luciferase activity assay. Data are presented as mean \pm SD of at least three independent experiments, with each measurement carried out in triplicate. Asterisks indicate significant difference between lobohedleolide- and DMSO-treated Ava5 cells. * $P < 0.05$; ** $P < 0.01$.

transcriptional element C/EBP was critical in the suppression of COX-2 promoter activity by lobohedleolide. To further verify which transcription factors are directly involved in the reduction effect of lobohedleolide on COX-2 promoter activity, Ava5 cells were transfected with individual luciferase reporter vector carrying each transcription element, including NF- κ B, C/EBP, and AP-1. The results of the reporter assay showed that lobohedleolide significantly reduced the HCV-induced luciferase activity mediated by C/EBP (Fig. 4b), but exhibited no significant effect on that of luciferase activity driven by NF- κ B or AP-1 (Fig. 4a,c). To confirm the inhibitory effect of lobohedleolide on the transcription factor activity of C/EBP in Ava5 cells, we created a COX-2 promoter reporter construct with a point mutation on C/EBP site alone, named as pCOX-2-(mC/EBP)-Luc (Fig. 4d). Ava5 cells were transfected with pCOX-2-Luc or pCOX-2-(mC/EBP)-Luc, and then the plasmid-transfected cells were incubated with lobohedleolide at concentrations ranging from 5 to 40 μ M. As shown in Fig. 4e, the COX-2/mC/EBP promoter activity was not affected by lobohedleolide compared with COX-2/WT promoter activity, confirming that the C/EBP transcription factor contributed to the suppression of HCV-induced COX-2 expression by lobohedleolide.

Lobohedleolide reduces HCV-induced C/EBP β binding and expression, c-Jun phosphorylation, and the activation of JNK.

To further examined whether lobohedleolide could reduce the binding of AP-1 and C/EBP β to the endogenous COX-2 promoter, we performed Chomatin Immunoprecipitation assay. The result showed that lobohedleolide decreased the C/EBP binding to COX-2 promoter (Fig. 5a), but lobohedleolide did not reduce the binding of AP-1 (Fig. 5b). A previous study has shown that the activated c-Jun could control the expression of C/EBP β . Therefore, we further examined the levels of C/EBP β protein and phosphorylated c-Jun in lobohedleolide-treated Ava5 cells, and the results of Western blotting showed that lobohedleolide dose-dependently reduced the C/EBP β protein level and the phosphorylation level of c-Jun (Fig. 5c). The activated JNK is the major upstream activator of c-Jun²⁴. To investigate whether the phosphorylation of JNK is involved in the inhibitory effect of lobohedleolide on COX-2 transcription, Ava5 cells were treated with lobohedleolide at 40 μ M for the indicated duration ranging from 0.5 to 6 h. The results of Western blotting showed that lobohedleolide significantly reduced the phosphorylation level of JNK in a time-dependent manner (Fig. 5d). In addition to the JNK/c-Jun-C/EBP β signaling pathway, COX-2 expression is also controlled by other transcriptional factors, including NF- κ B and MAPK²⁵. We found that lobohedleolide had no significant impact on the phosphorylation level of NF- κ B, ERK and p38 induced by HCV (Fig. 5e,f), suggesting that the JNK/c-Jun-C/EBP β signaling pathway is the most possible mechanism of lobohedleolide-mediated inhibition of COX-2 expression.

Combination treatment of lobohedleolide with clinically used anti-HCV agents synergistically reduces HCV replication. To examine the potential of lobohedleolide as a supplementary therapeutic agent, we investigated the antiviral effect using combination treatments of lobohedleolide with clinically used HCV




Figure 4. The reduction effect of loboheleolide on C/EBP transcription factor activity. Ava5 cells were cotransfected with 0.2 μg of pCMV-Rellina-Luc and (a) pNF-κB-Luc, (b) pC/EBP-Luc, or (c) pAP-1-Luc constructs for 6 h, and the transfected cells were treated with loboheleolide at the indicated concentrations for 3 days. Cell lysates were subjected to luciferase activity assay. (d) Schematic diagram of the COX-2 and mutagenized C/EBP of COX-2 promoter constructs. (e) The inhibitory effect of loboheleolide on COX-2 promoter activity was reduced by mutagenized C/EBP on COX-2 promoter. Ava5 cells were transfected with pCOX-2-Luc or pCOX-2-(mC/EBP)-Luc for 6 h, and the transfected cells were treated with loboheleolide at indicated concentrations for 3 days. Cell lysates were subjected to luciferase activity assay. Data are presented as mean ± SD of at least three independent experiments, with each measurement carried out in triplicate. Asterisks indicate significant difference between loboheleolide- and DMSO-treated Ava5 cells. *P < 0.05; **P < 0.01.

drugs, including IFN-α, the NS3/4A protease inhibitor telaprevir, the NS5B polymerase inhibitor sofosbuvir, and the NS5A inhibitor daclatasvir. The antiviral effects of the combination treatments were compared with the inhibitory effects of single treatments of the corresponding inhibitor in Ava5 cells, resulting in three CI values (ED₅₀, ED₇₅, and ED₉₀), through calculations as described in the Methods section. As shown in Table 1, the CI values are < 1 for all ED₅₀, ED₇₅, and ED₉₀ values, ranging from 0.12 to 0.59, showing that loboheleolide exhibited a synergistic effect with various inhibitors on HCV replication. There was no apparent cytotoxicity in all combination treatments (data not shown).

Discussion

Several studies have indicated that the inhibition of COX-2 by selective inhibitors or naturally available compounds could possibly be used as a strategy for inflammation treatment, cancer prevention, and suppression of virus replication^{26–28}. In the present study, we demonstrated that loboheleolide significantly reduced the HCV-induced COX-2 transcription, protein synthesis, and its metabolite PGE₂ production, resulting in the suppression of HCV replication (Figs 1 and 2). These observations indicate that loboheleolide could possibly be used for the treatment of HCV-related liver diseases with chronic inflammation in addition to a potential agent to conquer HCV infection.

Resistance to DAA agents is an incoming threat in treating chronic HCV infection with long-term treatment due to high mutation frequencies of RNA viruses. Therefore, host-targeting antiviral agents can be considered




Figure 5. The reduction effect of lobohedleolide on C/EBP β binding and expression, c-Jun phosphorylation, and JNK phosphorylation. **(a)** The reduction effect of lobohedleolide on the binding of C/EBP β to the endogenous COX-2 promoter. **(b)** Lobohedleolide did not significantly reduce the binding of AP-1 to the endogenous COX-2 promoter. Ava5 cells were treated with lobohedleolide at 20 and 40 μM for 3 days. Cells were fixed with formaldehyde. AP-1- or C/EBP β -bound DNA was isolated by chromatin immunoprecipitation with anti-AP-1 or anti-C/EBP β antibodies, respectively. The AP-1- or C/EBP-bound COX-2 promoter DNA was analyzed by PCR. **(c)** The suppression effect of C/EBP expression and c-Jun phosphorylation by lobohedleolide in Ava5 cells. Ava5 cells were treated with lobohedleolide at the indicated concentrations for 3 days, and the total cell lysate was subjected to Western blotting with relevant antibodies. **(d)** The reduction effect of lobohedleolide on JNK phosphorylation in Ava5 cells. **(e,f)** Lobohedleolide did not reduce the phosphorylation level of NF- κB , ERK, and p38. Ava5 cells were treated with lobohedleolide at 40 μM . Total cell lysates were extracted at the indicated time points and subjected to Western blotting with relevant antibodies. The relative blot intensities were quantified by densitometric scanning. GAPDH served as an equal loading control in Western blotting, with relevant antibodies. The relative blot intensities were quantified by densitometric scanning. GAPDH served as an equal loading control in Western blotting.

as an alternative strategy for overcoming the viral resistance due to the extremely low mutation rate of host genome in eukaryotic cells and the increasing drug susceptibility in all genotypes and serotypes²⁹. In addition, cocktail therapy has become a promising strategy to increase SVR and reduce drug resistance in HCV-infected patients^{30,31}. In the present study, we suggest lobohedleolide as a suitable anti-HCV agent based on its potential to target host COX-2 expression. Furthermore, lobohedleolide exhibited a synergistic anti-HCV effect with either IFN- α or other FDA-approved DAA agents (Table 1). Accordingly, lobohedleolide might be considered as a

Combination compound	Mean CI \pm SD at			
	ED50 [†]	ED75 [‡]	ED90 [§]	Influence
IFN- α	0.59 \pm 0.06	0.38 \pm 0.08	0.25 \pm 0.04	Synergistic
telaprevir	0.41 \pm 0.03	0.19 \pm 0.02	0.12 \pm 0.01	Synergistic
sofosbuvir	0.45 \pm 0.03	0.31 \pm 0.06	0.21 \pm 0.06	Synergistic
daclatasvir	0.53 \pm 0.04	0.34 \pm 0.02	0.26 \pm 0.05	Synergistic

Table 1. The synergistic reduction effect of lobohedleolide with IFN- α , telaprevir, or sofosbuvir on HCV replication. Standard deviations indicate the mean \pm SD of three independent experiments. [†]50% effective dose. [‡]75% effective dose. [§]90% effective dose.




Figure 6. The proposed model of lobohedleolide activity against HCV replication. Lobohedleolide reduced the HCV-induced COX-2 expression by downregulating the phosphorylation of JNK resulting in attenuation of the phosphorylation of c-Jun and the expression of C/EBP. Subsequently, the suppressed level of COX-2 and PGE₂ by lobohedleolide led to the inhibition of HCV replication.

dietary supplement for patients infected with HCV. Further investigation of the anti-HCV effect of lobohedleolide using a suitable animal model is required to analyze the potential use in HCV-infected patients.

An earlier study had demonstrated that lobohedleolide exhibited a reducing effect on LPS-induced iNOS and COX-2 expression in RAW264.7 macrophage cells²⁰, but the precise anti-inflammatory mechanism of lobohedleolide was not clearly investigated. In this study, we clearly demonstrated that lobohedleolide reduced the HCV-induced COX-2 transcription level by mediating the C/EBP regulatory element (Figs 3 and 4). The C/EBP family is a homodimeric DNA-binding basic-leucine zipper (bZIP) transcription factor and consists of six proteins divided into two subgroups, namely, C/EBP- α , - β , and - δ and C/EBP- γ , - ϵ , and - ζ .³² C/EBP- β and - δ are considered as the major mediators for regulating COX-2 expression^{33,34}. The effect of lobohedleolide on C/EBP isoform(s) involved in COX-2 expression in Avas cells remains to be investigated. In addition to C/EBP as an up-regulator for COX-2 expression during HCV infection, HCV-induced production of C/EBP mRNA causes an increase in the expression of inflammatory cytokines and chemokines, such as IL-1 α , TNF- α , and CXCL1³⁵, which contributes to chronic fibrogenesis and hepatocarcinogenesis³⁶. Clinically, a high expression level of IL-1 α , TNF- α , and IL-2 was observed in HCV-infected patients³⁷. In addition, the expression of a wide range of genes and microRNAs, such as miR-181c and miR-122, are also regulated by C/EBP, which participate in cell growth and proliferation in HCV-related liver diseases or HCC^{38,39}. Further investigation of the inhibitory effect of lobohedleolide on C/EBP-regulated inflammation in different chronic injuries could be performed, which would provide another potential therapeutic possibility of using lobohedleolide against HCV-related diseases. In contrast to the upregulator of COX-2 expression, the WT p53 has been considered as a suppressor of COX-2 transcription as it competes with the TATA-binding protein⁴⁰. Previous studies have shown that p53 expression is downregulated by the far upstream element (FUSE)-binding protein, leading to persistent HCV replication in most HCC tumors⁴¹. However, the relationship between p53 and COX-2 during HCV replication is not yet clearly verified. It will be useful to investigate the effect of lobohedleolide on p53-dependent inhibition on COX-2 expression during HCV infection.

In conclusion, our study results have revealed that lobohedleolide suppressed HCV replication by suppressing JNK phosphorylation, leading to the downregulation of c-Jun phosphorylation and C/EBP expression. Reduced COX-2 expression by lobohedleolide contributed to the suppression of PGE₂, finally causing a reduction on HCV RNA replication (Fig. 6). The synergistic anti-HCV effect of lobohedleolide with IFN, telaprevir, or sofosbuvir showed that lobohedleolide may serve as a therapeutic supplementary agent for increasing the treatment efficacy or decreasing the possibility of drug resistance in HCV-infected patients.

Materials and Methods

Cell culture and reagents. Ava5, an engineered HCV subgenomic replicon cell line, Huh-7 and Huh7.5 cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing 10% heat-inactivated fetal bovine serum, 1% antibiotic-antimycotic, and 1% nonessential amino acids at 37°C with a 5% CO₂ supplement. Ava5 cells have to be maintained in DMEM with 1 mg/ml G418 to maintain the stable expression of the replicon. Interferon alpha-2a (Roferon[®]-A) was purchased from Roche Ltd (Basel, Switzerland). Telaprevir was purchased from Legend Stat International Co., Ltd (Omdurman, Sudan). Daclatasvir and sofosbuvir was purchased from Shanghai Haoyuan Chemexpress Co., Ltd (Shanghai, China). Each experiment was incubated consistently in 0.1% dimethyl sulfoxide (DMSO). Telaprevir or sofosbuvir was stored as 10 mM in 100% DMSO.

Preparation of lobohedleolide. Lobohedleolide was isolated from the Formosan soft coral *L. crassum* following a method that has been described previously²⁰.

Western blotting. Ava5 cells were seeded into a 24-well plate at a density of 5×10^4 cells/well and into a 6-well plate at a density of 5×10^5 cells/well. After 12–16 h of incubation, cells were treated with the reagents for appropriate duration at the indicated concentrations. Then, the cells were washed with ice-cold PBS and lysed using RIPA lysis buffer. The insoluble protein was removed by centrifugation at 12,000 rpm for 30 min at 4°C, and the protein concentration of the soluble lysate was measured by Bio-Rad protein assay kit (Hercules, CA, USA). Immunoblotting analysis was performed as previously described. Briefly, an equal amount of protein was loaded onto 10% SDS-polyacrylamide gel electrophoresis and transferred to PVDF membranes. The levels of the protein of interest were measured using specific antibodies against HCV NS5B (1:5000; Abcam Cambridge, MA, USA), glyceraldehyde-3-phosphate dehydrogenase (GAPDH), COX-2 (1:1000; Cayman, ML, USA), Myc (1:1000; GeneTex, CA, USA), C/EBP, p-c-Jun, p-p38, t-p38, p-JNK, t-JNK, p-ERK, and t-ERK (1:1000; Cell Signaling Technology, Danvers, MA, USA). The blotting signal was developed using ECL detection kit (PerkinElmer, CT, USA) and was counted by the software Quantity One (Bio-Rad, CA, USA).

Quantitative real-time PCR (qRT-PCR) assay. The cells were incubated with lobohedleolide for appropriate duration at the indicated concentrations. Total cellular RNA was extracted using a Total RNA Miniprep Purification Kit (GMBiolab, Kaohsiung, Taiwan) according to the manufacturer's instructions. After being transcribed to cDNA by M-MLV reverse transcriptase (Promega, Madison, WI, USA) with HCV 3'UTR (5'-acttgatctgcagagaggcc-3') or oligo dT primer, the level of cDNA was determined through qRT-PCR with specific primers as previously described. The CT value of each sample was determined using the ABI Step One Real-Time PCR System, which was normalized to endogenous cellular *gapdh* gene. The PCR primers were as follows: GAPDH, 5'-gtcttcaccaccatggagaa-3' (forward), and 5'-atggcatggactgtggtcat-3' (reverse); NS5B, 5'-ggaaccaagctgccatca-3' (forward), and 5'-ctccacgatagaagtta-3' (reverse).

Cytotoxicity assay. Cell cytotoxicity was determined using colorimetric 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) assay (Promega, Madison, WI, USA) according to the manufacturer's instructions. Briefly, 5×10^3 Ava5 cells/well were seeded into a 96 well-plate and treated with lobohedleolide at the indicated concentrations for 3 days. The cells were then treated with 20 µl MTS solution and 80 µl PBS per well for 2–4 h at 37°C, and the relative cytotoxicity was quantified by measuring the absorbance at 490 nm using a 550 BioRad plate-reader (Bio-Rad, Hertfordshire, UK).

HCV particle preparation and infection assay. Full-length and linearized JFH-1 RNA was transfected into Huh7.5 cells for producing infectious HCV genomic type 2a JFH-1 virus particles⁴². After 3 days, the supernatant was collected and filtered through a 0.45-µm filter and stored at -70°C until use. The infectivity rate of HCV JFH-1 was determined through serial diluting and infecting the Huh7.5 cells for 3 days. Immunostaining was performed with anti-core antibodies. Huh7.5 cells were infected with HCV JFH-1 particles at a multiplicity of infection of 0.1 for 6 h, and the infected cells were washed with PBS and replaced with fresh medium. Lobohedleolide was applied at various concentrations for an additional 3 days. Total RNA was extracted and subjected to qRT-PCR as described above.

Transfection and luciferase activity assay. The Ava5 cells were seeded into a 24-well plate at a density of 5×10^4 cells/well and incubated for 16 h. Then, 0.2 µg of pCMV-Rellina-Luc and 1 µg of a series of COX-2 promoter reporter plasmids, including pCOX-2-Luc, pCOX-2-(ΔNF-κB)-Luc, and pCOX-2-(ΔNF-κB/C/EBP)-Luc were cotransfected into Ava5 cells by T-pro reagent (Ji-Feng Biotechnology CO., Ltd., Taipei, Taiwan) for 6 h following the manufacturer's instructions. To investigate the critical transcription factor involved in the inhibition of COX-2 expression by lobohedleolide, Ava5 cells were cotransfected with 0.2 µg of pCMV-Rellina-Luc and pAP-1-Luc, pNF-κB-Luc, or pC/EBP-Luc for 6 h. pCMV-Rellina-Luc served as the internal control for evaluating the transfection efficiency. The cells were exchanged with fresh medium containing lobohedleolide at the indicated concentrations. After 3 days of incubation, the cell extracts were subjected to luciferase activity assay using Dual-Glo Luciferase Assay System (Promega, Madison, WI, USA) in accordance with the manufacturer's instructions. pAP-1-Luc, pNF-κB-Luc and pC/EBP-Luc were purchased from Stratagene. Under exogenous gene expression condition, either vehicle vector pcDNA4/myc-His-A (Life Technologies, Carlsbad, CA, USA) or pcDNA4-COX-2-myc-His was transfected into Ava5 cells at various concentration (0.5, 1.0 and 1.5 µg) for 6 h. The cells were exchanged with fresh medium containing lobohedleolide at 40 µM for 3 days. The cell lysate and cellular RNA were analyzed by Western blotting and qRT-PCR as previously described, respectively.

Chromatin Immunoprecipitation assay. The ability of C/EBPβ to bind to COX-2 promoter upon lobohedleolide treatment was examined using Magna ChIP™ A/G kit according to the manufacturer's protocol

(Merck, Darmstadt, Germany). Briefly, Ava5 cells were treated with lobohedleolide for 3 days. Cells were then cross-linked with 1% formaldehyde for 10 minutes. After washing with PBS, cells were lysed with Cell Lysis Buffer containing protease inhibitor cocktail. The chromatin was sheared by sonication. 100 ng of DNA were used as input. The remaining chromatin fraction was immunoprecipitated with anti-C/EBP β (4 μ g; GeneTex, CA, USA) or anti-AP-1 antibodies (10 μ g; Sigma, MO, USA) overnight at 4 °C. The C/EBP β -bound COX-2 promoter DNA was analyzed by PCR with specific primers: 5'-cggagggtatgctcatgaaa-3' (forward), and 5'-caggctttaccacgcaaa-3' (reverse). PCR was performed at 94 °C for 30 s, 59 °C for 30 s, and 72 °C for 30 s, for 35 cycles.

Analysis of the drug synergism. Ava5 cells were seeded into a 24-well plate at a density of 5×10^4 cells/well and treated with serially diluted lobohedleolide at 2.5, 5, 10 and 20 μ M in combination with diluted IFN- α (7.5, 15, 30, and 60 U/mL), the HCV protease inhibitor telaprevir (0.075, 0.15, 0.3, and 0.6 μ M), or the RNA-dependent RNA polymerase nucleoside inhibitor sofosbuvir (10, 20, 40, and 80 nM). Each of the combination treatment was performed by adding lobohedleolide horizontally with various HCV inhibitors vertically in a checkerboard cross in the 24-well plate. After 3 days of treatment, total cellular RNA was extracted, and the RNA levels were quantified by qRT-PCR with specific primers as described above. The combination index (CI) values of each combination achieving 50%, 75%, or 95% reduction in the HCV RNA level were calculated using the CalcuSyn2TM computer program (Biosoft, Cambridge, UK), which was based on the Chou and Talalay analysis method^{43,44}. Primarily, CI values of 1, <1, and >1 indicate additive, synergistic, and antagonistic effects, respectively.

Statistical analysis. All data were obtained from at least three independent experiments and are presented as mean \pm SD. Statistical significance was determined using Student's *t* test for differences between two data groups (lobohedleolide-treated and -untreated cells). **P* < 0.05 or ***P* < 0.01 was considered to be statistically significant.

References

- Ke, P. Y. & Chen, S. S. Hepatitis C virus and cellular stress response: implications to molecular pathogenesis of liver diseases. *Viruses* **4**, 2251–2290, <https://doi.org/10.3390/v4102251> (2012).
- Poordad, F. *et al.* Boceprevir for untreated chronic HCV genotype 1 infection. *The New England journal of medicine* **364**, 1195–1206, <https://doi.org/10.1056/NEJMoa1010494> (2011).
- Reau, N., Hamzeh, F. M., Lentz, E., Zhou, X. & Jensen, D. Characterization of nonrapid virologic response patients infected with HCV genotype 1 who may relapse after standard therapy with peginterferon plus ribavirin. *Journal of viral hepatitis* **19**, 94–102, <https://doi.org/10.1111/j.1365-2893.2010.01422.x> (2012).
- Stedman, C. Sofosbuvir a NS5B polymerase inhibitor in the treatment of hepatitis C: a review of its clinical potential. *Therapeutic advances in gastroenterology* **7**, 131–140, <https://doi.org/10.1177/1756283X13515825> (2014).
- Poveda, E. *et al.* Update on hepatitis C virus resistance to direct-acting antiviral agents. *Antiviral research* **108**, 181–191, <https://doi.org/10.1016/j.antiviral.2014.05.015> (2014).
- Chhatwal, J., Kanwal, F., Roberts, M. S. & Dunn, M. A. Cost-effectiveness and budget impact of hepatitis C virus treatment with sofosbuvir and ledipasvir in the United States. *Annals of internal medicine* **162**, 397–406, <https://doi.org/10.7326/M14-1336> (2015).
- Lindenbach, B. D. & Rice, C. M. Unravelling hepatitis C virus replication from genome to function. *Nature* **436**, 933–938, <https://doi.org/10.1038/nature04077> (2005).
- Penin, F., Dubuisson, J., Rey, F. A., Moradpour, D. & Pawlotsky, J. M. Structural biology of hepatitis C virus. *Hepatology* **39**, 5–19, <https://doi.org/10.1002/hep.20032> (2004).
- Chen, W. C. *et al.* HCV NS5A Up-Regulates COX-2 Expression via IL-8-Mediated Activation of the ERK/JNK MAPK Pathway. *PLoS one* **10**, e0133264, <https://doi.org/10.1371/journal.pone.0133264> (2015).
- El-Bassiouny, A. E. *et al.* Expression of cyclooxygenase-2 and transforming growth factor-beta1 in HCV-induced chronic liver disease and hepatocellular carcinoma. *MedGenMed: Medscape general medicine* **9**, 45 (2007).
- Lin, Y. T. *et al.* Green tea phenolic epicatechins inhibit hepatitis C virus replication via cyclooxygenase-2 and attenuate virus-induced inflammation. *PLoS one* **8**, e54466, <https://doi.org/10.1371/journal.pone.0054466> (2013).
- Lin, C. K. *et al.* Betulinic acid exerts anti-hepatitis C virus activity via the suppression of NF-kappaB and MAPK-ERK1/2-mediated cyclooxygenase-2 expression. *British journal of pharmacology*, <https://doi.org/10.1111/bph.13233> (2015).
- Chen, K. J. *et al.* Aqueous extract of the edible *Gracilaria tenuistipitata* inhibits hepatitis C viral replication via cyclooxygenase-2 suppression and reduces virus-induced inflammation. *PLoS one* **8**, e57704, <https://doi.org/10.1371/journal.pone.0057704> (2013).
- Gee, J., Lee, I. L., Grossman, H. B. & Sabichi, A. L. Forced COX-2 expression induces PGE(2) and invasion in immortalized urothelial cells. *Urologic oncology* **26**, 641–645, <https://doi.org/10.1016/j.urolonc.2007.05.015> (2008).
- Lu, L. *et al.* NS3 protein of hepatitis C virus regulates cyclooxygenase-2 expression through multiple signaling pathways. *Virology* **371**, 61–70, <https://doi.org/10.1016/j.virol.2007.09.025> (2008).
- Inoue, H., Nanayama, T., Hara, S., Yokoyama, C. & Tanabe, T. The cyclic AMP response element plays an essential role in the expression of the human prostaglandin-endoperoxide synthase 2 gene in differentiated U937 monocytic cells. *FEBS letters* **350**, 51–54 (1994).
- Mestre, J. R. *et al.* Retinoids suppress phorbol ester-mediated induction of cyclooxygenase-2. *Cancer research* **57**, 1081–1085 (1997).
- Rashid, M. A., Gustafson, K. R. & Boyd, M. R. HIV-inhibitory cembrane derivatives from a Philippines collection of the soft coral *Lobophytum* species. *Journal of natural products* **63**, 531–533 (2000).
- Radhika, P., Rao, P. R., Archana, J. & Rao, N. K. Anti-inflammatory activity of a new sphingosine derivative and cembreoid diterpene (lobohedleolide) isolated from marine soft corals of *Sinularia crassa* TIXIER-DURIVAUULT and *Lobophytum* species of the Andaman and Nicobar Islands. *Biological & pharmaceutical bulletin* **28**, 1311–1313 (2005).
- Chao, C. H., Wen, Z. H., Wu, Y. C., Yeh, H. C. & Sheu, J. H. Cytotoxic and anti-inflammatory cembranoids from the soft coral *Lobophytum crassum*. *Journal of natural products* **71**, 1819–1824, <https://doi.org/10.1021/np8004584> (2008).
- Lin, C. K. *et al.* Betulinic acid exerts anti-hepatitis C virus activity via the suppression of NF-kappaB- and MAPK-ERK1/2-mediated COX-2 expression. *British journal of pharmacology*, <https://doi.org/10.1111/bph.13233> (2015).
- Prescott, S. M. & Fitzpatrick, F. A. Cyclooxygenase-2 and carcinogenesis. *Biochim Biophys Acta* **1470**, M69–78 (2000).
- Healy, Z. R., Zhu, F., Stull, J. D. & Konstantopoulos, K. Elucidation of the signaling network of COX-2 induction in sheared chondrocytes: COX-2 is induced via a Rac/MEKK1/MKK7/JNK2/c-Jun-C/EBPbeta-dependent pathway. *American journal of physiology. Cell physiology* **294**, C1146–1157, <https://doi.org/10.1152/ajpcell.00542.2007> (2008).
- Johnson, G. L. & Nakamura, K. The c-jun kinase/stress-activated pathway: regulation, function and role in human disease. *Biochim Biophys Acta* **1773**, 1341–1348, <https://doi.org/10.1016/j.bbamer.2006.12.009> (2007).
- Kim, A. R. *et al.* Phlorofucofuroeckol A inhibits the LPS-stimulated iNOS and COX-2 expressions in macrophages via inhibition of NF-kappaB, Akt, and p38 MAPK. *Toxicology in vitro: an international journal published in association with BIBRA* **25**, 1789–1795, <https://doi.org/10.1016/j.tiv.2011.09.012> (2011).

26. Leng, J., Han, C., Demetris, A. J., Michalopoulos, G. K. & Wu, T. Cyclooxygenase-2 promotes hepatocellular carcinoma cell growth through Akt activation: evidence for Akt inhibition in celecoxib-induced apoptosis. *Hepatology* **38**, 756–768, <https://doi.org/10.1053/jhep.2003.50380> (2003).
27. Kawamori, T., Rao, C. V., Seibert, K. & Reddy, B. S. Chemopreventive activity of celecoxib, a specific cyclooxygenase-2 inhibitor, against colon carcinogenesis. *Cancer research* **58**, 409–412 (1998).
28. Tseng, C. K. *et al.* Aqueous extract of *Gracilaria tenuistipitata* suppresses LPS-induced NF-kappaB and MAPK activation in RAW 264.7 and rat peritoneal macrophages and exerts hepatoprotective effects on carbon tetrachloride-treated rat. *PLoS one* **9**, e86557, <https://doi.org/10.1371/journal.pone.0086557> (2014).
29. Wohlfarth, C. & Efferth, T. Natural products as promising drug candidates for the treatment of hepatitis B and C. *Acta Pharmacol Sin* **30**, 25–30, <https://doi.org/10.1038/aps.2008.5aps20085> (2009).
30. Karayiannis, P. Direct acting antivirals for the treatment of chronic viral hepatitis. *Scientifica* **2012**, 478631, <https://doi.org/10.6064/2012/478631> (2012).
31. Gelman, M. A. & Glenn, J. S. Mixing the right hepatitis C inhibitor cocktail. *Trends in molecular medicine* **17**, 34–46, <https://doi.org/10.1016/j.molmed.2010.10.005> (2011).
32. Wang, W., Bergh, A. & Damber, J. E. Increased expression of CCAAT/enhancer-binding protein beta in proliferative inflammatory atrophy of the prostate: relation with the expression of COX-2, the androgen receptor, and presence of focal chronic inflammation. *Prostate* **67**, 1238–1246, <https://doi.org/10.1002/pros.20595> (2007).
33. Wadleigh, D. J., Reddy, S. T., Kopp, E., Ghosh, S. & Herschman, H. R. Transcriptional activation of the cyclooxygenase-2 gene in endotoxin-treated RAW 264.7 macrophages. *The Journal of biological chemistry* **275**, 6259–6266 (2000).
34. Zhu, Y., Saunders, M. A., Yeh, H., Deng, W. G. & Wu, K. K. Dynamic regulation of cyclooxygenase-2 promoter activity by isoforms of CCAAT/enhancer-binding proteins. *The Journal of biological chemistry* **277**, 6923–6928, <https://doi.org/10.1074/jbc.M108075200> (2002).
35. Xu, G., Zhang, Y., Zhang, L., Roberts, A. I. & Shi, Y. C/EBPbeta mediates synergistic upregulation of gene expression by interferon-gamma and tumor necrosis factor-alpha in bone marrow-derived mesenchymal stem cells. *Stem cells* **27**, 942–948, <https://doi.org/10.1002/stem.22> (2009).
36. Lu, X. L. *et al.* Chemopreventive effect of saikosaponin-d on diethylnitrosamine-induced hepatocarcinogenesis: involvement of CCAAT/enhancer binding protein beta and cyclooxygenase-2. *Molecular medicine reports* **5**, 637–644, <https://doi.org/10.3892/mmr.2011.702> (2012).
37. Kasprzak, A. *et al.* Expression of cytokines (TNF-alpha, IL-1alpha, and IL-2) in chronic hepatitis C: comparative hybridocytochemical and immunocytochemical study in children and adult patients. *The journal of histochemistry and cytochemistry: official journal of the Histochemistry Society* **52**, 29–38 (2004).
38. Mukherjee, A., Shrivastava, S., Bhanja Chowdhury, J., Ray, R. & Ray, R. B. Transcriptional suppression of miR-181c by hepatitis C virus enhances homeobox A1 expression. *Journal of virology* **88**, 7929–7940, <https://doi.org/10.1128/JVI.00787-14> (2014).
39. Zeng, C. *et al.* A novel GSK-3 beta-C/EBP alpha-miR-122-insulin-like growth factor 1 receptor regulatory circuitry in human hepatocellular carcinoma. *Hepatology* **52**, 1702–1712, <https://doi.org/10.1002/hep.23875> (2010).
40. Chun, K. S. & Surh, Y. J. Signal transduction pathways regulating cyclooxygenase-2 expression: potential molecular targets for chemoprevention. *Biochem Pharmacol* **68**, 1089–1100, <https://doi.org/10.1016/j.bcp.2004.05.031> (2004).
41. Dixit, U. *et al.* FUSE Binding Protein 1 Facilitates Persistent Hepatitis C Virus Replication in Hepatoma Cells by Regulating Tumor Suppressor p53. *Journal of virology* **89**, 7905–7921, <https://doi.org/10.1128/JVI.00729-15> (2015).
42. Kato, T. *et al.* Cell culture and infection system for hepatitis C virus. *Nature protocols* **1**, 2334–2339, <https://doi.org/10.1038/nprot.2006.395> (2006).
43. Chou, T. C. & Talalay, P. Generalized equations for the analysis of inhibitions of Michaelis-Menten and higher-order kinetic systems with two or more mutually exclusive and nonexclusive inhibitors. *European journal of biochemistry/FEBS* **115**, 207–216 (1981).
44. Chou, T. C. & Talalay, P. Quantitative analysis of dose-effect relationships: the combined effects of multiple drugs or enzyme inhibitors. *Advances in enzyme regulation* **22**, 27–55 (1984).

Acknowledgements

We are grateful to Dr. Charles Rice (Rockefeller University) and Apath, LLC, NY, USA for kindly supporting human hepatoma cell line (Huh-7 and Huh-7.5) and hepatitis C virus replicon cell line Ava5. We also thank T. Wakita (National Institute of Infectious Diseases, Japan) HCVcc plasmid. This work was supported by a grant from the Ministry Science and Technology of Taiwan (MOST-104-2320-B-037-025-MY3), a grant from the National Sun Yat-Sen University-KUM Joint Research Project of Taiwan (NSYSU-KMU 106-I007), and KMU-KMUH Co-Project of Key Research Grant (KMU-DK107011) from Kaohsiung Medical University of Taiwan

Author Contributions

Jin-Ching Lee, Jyh-Horng Sheu and Chih-Chuang Liaw performed the experimental design. Chun-Kuang Lin, Chin-Kai Tseng and Chiung-Yao Huang performed the research as described in the Material and Method. Jyh-Horng Sheu and Chiung-Yao Huang contributed essential reagents. Chun-Kuang Lin, Chin-Kai Tseng and Chih-Ku Wei analyzed the data. Jin-Ching Lee and Chun-Kuang Lin wrote the paper. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-26999-w>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018