





SCIENTIFIC REPORTS



OPEN

Publisher Correction: IMSindel: An accurate intermediate-size indel detection tool incorporating *de novo* assembly and gapped global-local alignment with split read analysis

Daichi Shigemizu^{1,2,3,4,5} , Fuyuki Miya^{2,3}, Shintaro Akiyama¹, Shujiro Okuda⁶ , Keith A. Boroevich³, Akihiro Fujimoto⁷, Hidewaki Nakagawa³, Kouichi Ozaki^{1,3}, Shumpei Niida¹, Yonehiro Kanemura^{8,9}, Nobuhiko Okamoto¹⁰, Shinji Saitoh¹¹ , Mitsuhiro Kato¹², Mami Yamasaki¹³, Tatsuo Matsunaga¹⁴, Hideki Mutai¹⁴, Kenjiro Kosaki¹⁵ & Tatsuhiko Tsunoda^{2,3,4,5} 

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-018-23978-z>, published online 04 April 2018

This Article contains an error in the Results section under the subheading ‘Performance comparison among call methods using simulation data’.

“We found that when the insertions were ≥ 2 bp length and contained repetitive sequence from the flanking region, *de novo* assembly (Inchworm) did not work well (Table S3).”

should read:

“We found that when the insertions were ≥ 25 bp length and contained repetitive sequence from the flanking region, *de novo* assembly (Inchworm) did not work well (Table S3).”

¹Department for Medical Genome Sciences, Medical Genome Center, National Center for Geriatrics and Gerontology, Aichi, Japan. ²Department of Medical Science Mathematics, Medical Research Institute, Tokyo Medical and Dental University (TMDU), Tokyo, Japan. ³RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. ⁴Medical Sciences Innovation Hub Program, Cluster for Science and Technology Hub, RIKEN, Yokohama, Japan. ⁵CREST, JST, Japan. ⁶Niigata University Graduate School of Medical and Dental Sciences, Niigata, Japan. ⁷Department of Drug Discovery Medicine, Graduate School of Medicine, Kyoto University, Kyoto, Japan. ⁸Division of Regenerative Medicine, Institute for Clinical Research, Osaka National Hospital, National Hospital Organization, Osaka, Japan. ⁹Department of Neurosurgery, Osaka National Hospital, National Hospital Organization, Osaka, Japan. ¹⁰Department of Medical Genetics, Osaka Medical Center and Research Institute for Maternal and Child Health, Osaka, Japan. ¹¹Department of Pediatrics and Neonatology, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan. ¹²Department of Pediatrics, Showa University School of Medicine, Tokyo, Japan. ¹³Department of Pediatric Neurosurgery, Takatsuki General Hospital, Osaka, Japan. ¹⁴Division of Hearing and Balance Research, National Institute of Sensory Organs, National Hospital Organization Tokyo Medical Center, Tokyo, Japan. ¹⁵Center for Medical Genetics, Keio University School of Medicine, Tokyo, Japan. Daichi Shigemizu and Fuyuki Miya contributed equally to this work. Correspondence and requests for materials should be addressed to D.S. (email: daichi@ncgg.go.jp) or T.T. (email: tsunoda.mesm@mri.tmd.ac.jp)



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018