

SCIENTIFIC REPORTS

There are amendments to this paper

OPEN

Conversion of tectonic and climatic forcings into records of sediment supply and provenance

Glenn R. Sharman¹, Zoltan Sylvester² & Jacob A. Covault²

Understanding how environmental forcings (e.g., tectonics, climate) are transformed by erosional landscapes into sedimentary signals is a critical component of inverting the stratigraphic record. Previous research has largely focused on sediment supply (Q_s) and grain size as the de facto sedimentary signals of changing forcing mechanisms. We use a numerical model to consider the paired response of sediment provenance (P_v), expressed as fractional sediment load, and Q_s to demonstrate that the same change in environmental forcing may have a different expression in the sedimentary record. While Q_s reflects integrated denudation across an erosional catchment, P_v is controlled by spatially variable erosion that occurs in transient landscapes. P_v from proximal sediment sources increases during upstream knickpoint migration, whereas P_v from distal sediment sources increases when bedrock channels incise to produce lower gradient profiles. Differences between the Q_s and P_v signals relate to distinct geomorphic processes that operate on different time scales and allow for a refined differentiation of the timing and mechanism of forcings than possible via analysis of either signal alone. Future efforts to integrate multiple sedimentary signals may thus yield a richer picture of underlying forcing mechanisms, facilitating efforts to invert the stratigraphic record.

Sedimentary deposits represent an important archive of how environmental conditions (e.g., tectonics, climate) have changed over geologic time. Using the stratigraphic record to differentiate and reconstruct such environmental forcing mechanisms is of critical importance, as understanding how the Earth's surface responded to past change is key to predicting the response of the Earth to ongoing and future change, such as a warming climate¹. A large body of research, based in part on analog and numerical experiments, has highlighted a number of challenges in using sedimentary data to reconstruct external forcing mechanisms, including the tendency for sedimentary signals to be non-unique, modified when the forcing timescale is less than the landscape response time, and shredded or attenuated during sediment transport and deposition^{2–8}. Others have found success in inverting the stratigraphic record, but under a limited set of scenarios (e.g., down-system grain size trends^{9–12}).

Previous research has largely considered sediment supply (Q_s), and to a lesser extent grain size, as the de facto sedimentary signal of changing forcing mechanisms^{4,5,11–16}. Although Q_s is an important sedimentary parameter (e.g., influences system progradation versus retrogradation) and is readily measured in numerical^{4,15,16} and analog¹⁷ experiments, Q_s is notoriously difficult to constrain in modern, and particularly ancient, sedimentary systems^{12,18}. Sedimentary provenance (P_v) can be interpreted in both modern and ancient systems via numerous, different proxies that are applicable to gravel, sand, and mud grain size fractions^{19,20}. We define P_v as the proportion of sediment mass eroded from the i^{th} source rock type:

$$P_{vi}(\%) = c_i Q_{si} / \sum_{j=1}^n c_j Q_{sj} \quad (1)$$

where c is a dimensionless coefficient that reflects the abundance of a given provenance indicator in a source area and n is the number of sediment sources within the catchment area²¹. Other sedimentary data types (e.g., sediment composition, thermochronology, stratigraphic architecture) may also signal changing forcing mechanisms, and how such signals are propagated and preserved in sedimentary systems is an active research topic⁵.

¹Department of Geosciences, University of Arkansas, Fayetteville, AR, 72701, USA. ²Bureau of Economic Geology, Jackson School of Geosciences, The University of Texas at Austin, Austin, TX, 78758, USA. Correspondence and requests for materials should be addressed to G.R.S. (email: gsharman@uark.edu)

Received: 21 September 2018

Accepted: 28 January 2019

Published online: 11 March 2019




Figure 1. (a) Landscape model geometry showing the aerial distribution of Source A and Source B. Variations on this model geometry were used in Scenarios 1.10–1.15 and 2.10–2.15 (Table DR1). (b) Explanation of experimental design and terms for a stepped increase and decrease in forcing (thick, dashed black line). X-axis displays elapsed time since initiation of experiment following establishment of steady-state. Scenarios 1.1 (uplift rate) and 2.1 (precipitation) are shown (see Tables DR1 and DR2). Minimum and maximum Q_s and P_{vA} values attained during stepped forcing are identified as Q_{s_min} , P_{vA_min} , Q_{s_max} , and P_{vA_max} respectively. Abbreviations: RT-response time.

This study explores the paired response of Q_s and P_v to temporally variable tectonic and climatic forcings on an erosional landscape. Specifically, we aim to answer the following questions. (1) Do Q_s and P_v differ in response to stepped and periodic changes in uplift and precipitation rates? (2) What are the underlying geomorphic causes for differences between Q_s and P_v ? (3) Can the paired $Q_s - P_v$ signal improve the reconstruction of environmental forcing type (tectonics versus climate) and timing?

Experimental Design and Model Parameters

Following the experimental design of previous workers¹⁶, we simulate erosion of a two-dimensional landscape using a detachment-limited bedrock channel erosion rate (E) of

$$E = K(rA)^m S^n \quad (2)$$

where K is a coefficient of erosion, r is precipitation rate, A is the upstream drainage area, S is the slope, and m and n are constants that reflect non-linearity between E , rA , and S ²². Together, the product of r and A is a proxy for water discharge that scales with precipitation for a catchment of a given size. Although this stream-power incision model is oversimplified with respect to natural rivers²³, it has been successfully applied to mountainous rivers in different climate regimes²⁴ and provides a minimum estimate of system response time from which to evaluate signal transmission from changing forcing mechanisms²⁵. Use of the detachment-limited stream-power incision model thus provides an optimistic (best-case) view of signal transmission, as signals are more readily transmitted from landscapes with short, or ‘rapid,’ equilibrium response times, defined as the time elapsed between a forcing perturbation and subsequent achievement of topographic and denudational steady-state^{5,13,14,25}. To carry out the landscape evolution modeling, we have used the open-source, Python-based Landlab toolbox²⁶, with Equation 2 implemented using the FastScapeEroder algorithm²⁷.

We follow model parameters of previous workers¹⁶ with a square grid with dimensions of 64 km² (Fig. 1; model parameters are presented in Table DR1). All four of the model’s bounding edges are closed boundaries with the exception of a single outlet point at the midpoint of the southern boundary that has a fixed value of 0 m elevation (Fig. 1). The model is further divided into two regions (Source A and Source B) positioned proximally and distally to the outlet point, respectively (Fig. 1). These regions represent different rock types in the landscape model for the purpose of modeling change in P_v during landscape evolution.

We conducted a series of experiments to determine the Q_s and P_v responses to a 10-fold increase and decrease in both uplift rate (scenario 1.1) and precipitation (scenario 2.1)^{4,16} (Fig. 2). Additional experimental runs were conducted to test the sensitivity of model results to absolute forcing magnitude (2- and 5-fold-changes in uplift and precipitation rates, scenarios 1.2–1.3 and 2.2–2.3); source area erodibility (K) (scenarios 1.4–1.5 and 2.4–2.5); provenance weighting coefficients (c) (scenarios 1.6–1.7 and 2.6–2.7); and model time step, size, and geometry (scenarios 1.8–1.15 and 2.8–2.15) (Table DR1). All model results are presented in Supplemental Fig. 1.

Each experiment contains three phases (Fig. 2). (1) An initially flat model surface (0 m elevation) is seeded with small, random variations to allow a stream network to develop. Uplift is added to the surface at a rate of 1 mm/yr, and the model is run until topographic steady-state is achieved (~2–2.5 Myr). (2) The model is then




Figure 2. Experimental results. **(a)** Forcing (uplift rate and precipitation) and the corresponding sedimentary signal. Top plot shows Q_s and P_{vA} (% Source A) and the bottom plot shows the Q_s from Source A and Source B (Q_{sA} and Q_{sB} , respectively). See Fig. 1 and text for an explanation. **(b)** Channel slope profiles, map of erosion rate, and elevation during periods of transient landscape response to increased (above) and decreased (below) forcing. Left and right panels correspond to changes in uplift rate and precipitation, respectively, shown in (a). The maps correspond to the channel slope profile colored red and to the red and blue stars in (a). Abbreviations: RT-response time.

subjected to a stepped increase and subsequent stepped decrease in either uplift rate or precipitation (Fig. 2). The time duration of each step of the initial phase is sufficiently long to allow the model to reach steady-state (Fig. 2). We follow^{4,16} in defining the signal response time (Q_s or P_v) as the time elapsed to return to within a threshold of the steady-state value (here defined as within 1%). (3) Following a similar procedure as previous workers¹⁶, the final phase consists of periodic, stepped forcing increase and decrease where the duration of each cycle is a percentage (100%, 75%, 50%, 25%, and 10%) of the Q_s response time determined during the second phase of the experiment (Fig. 2). The model is allowed to reach dynamic equilibrium, a condition where the Q_s and P_v signals are identical to the preceding cycle¹⁶, before the forcing period is changed (3 to 8 forcing cycles; Fig. 2). Because Q_s response time can differ for increasing versus decreasing forcing, periods of high and low forcing values are not necessarily of equal duration (Fig. 1b). This asymmetry is most pronounced for changes in precipitation, where the Q_s and P_v response times are much longer for a decrease in precipitation versus an increase (Fig. 2a).

Results

Signal Response to Stepped Forcing. Uplift Rate. Given a stepped change in uplift rate, Q_s increases to a new equilibrium value ($Q_{s,eq}$) such that $Q_{s,eq} = U_{new}A$, where U_{new} is the new rate of uplift and A is the catchment area. Maximum ($Q_{s,max}$) and minimum ($Q_{s,min}$) values are thus attained when the model is at steady-state during high and low uplift rates, respectively. In this case, $Q_{s,min} \approx 6 \text{ m}^3/\text{yr}$ and $Q_{s,max} \approx 60 \text{ m}^3/\text{yr}$, with a response time of $\sim 0.93 \text{ Myr}$ for both increased and decreased uplift rate (Figs 1b and 2; Table DR2).

Unlike Q_s , P_v does not tend towards a new equilibrium value, but instead exhibits a transient behavior where relative proximal erosion increases during an increase in uplift rate (peak change in P_{vA} , $P_{vA,max} \approx 70\%$ Source A) and relative distal erosion increases during a decrease in uplift rate ($P_{vA,min} \approx 38\%$ Source A), before returning to the 50% equilibrium value that reflects the areal distribution of rock types A and B (Figs 1b and 2). The lag time between uplift rate change and $P_{vA,max}$ is $\sim 0.2 \text{ Myr}$, which is considerably shorter than the lag time of $Q_{s,max}$ ($\sim 0.9 \text{ Myr}$). P_v also exhibits an asymmetry in response time: $\sim 0.9 \text{ Myr}$ for uplift rate increase and $\sim 1.2 \text{ Myr}$ for uplift rate decrease (Figs 1b and 2a).

Precipitation. As noted previously⁴, an increase in precipitation results in a rapid but temporary increase in Q_s followed by a return to $Q_{s,eq}$. In our model, $Q_{s,max}$ ($19 \text{ m}^3/\text{yr}$) is attained rapidly with a response time of $\sim 0.3 \text{ Myr}$. During a decrease in precipitation, $Q_{s,min}$ ($\sim 2 \text{ m}^3/\text{yr}$) is also achieved rapidly but with a more prolonged response time ($\sim 0.9 \text{ Myr}$).




Figure 3. Sedimentary signal (Q_s and P_{vA}) response in dynamic equilibrium (data position are shown in Fig. 2a). (a) Signal response normalized to forcing period. Dashed, black arrows indicate general direction of peak and trough translation as response time (RT) decreases. (b) Q_s versus P_{vA} .

The P_v signal responds to changes in precipitation in an opposite way as to changes in uplift rate; distal erosion increases ($P_{vA_max} \approx 42\%$ Source A) during precipitation increase and relative proximal erosion increases ($P_{vA_max} \approx 61\%$ Source A) during precipitation decrease (Fig. 2). The lag time between precipitation increase and P_{vA_max} is ~ 0.2 Myr, whereas the lag time associated with a decrease in precipitation is longer (~ 0.6 Myr). The P_v response times for a precipitation increase and decrease (~ 0.4 Myr and ~ 0.9 Myr, respectively) are both less than the equivalent values for changing uplift rate (Fig. 2).

Signal Response to Periodic Forcing. Our results corroborate previous research that suggests the Q_s signal is progressively modified as forcing periodicity becomes less than the Q_s response time^{3,14,16} and demonstrate that a similar pattern is characteristic of the P_v signal (Fig. 3).

Uplift Rate. As the periodicity of changing uplift rate becomes less than the Q_s response time, the relative amplitude of the Q_s signal is progressively damped such that the signal amplitude is $< \sim 10\%$ of the maximum possible signal ($Q_{s_max} - Q_{s_min}$) when the forcing period is 25% of the response time (Fig. 3a; Table DR3). At the same time, the relative timing of the peak Q_s migrates forward such that the peak is increasingly offset from when uplift rate increased (Fig. 3a). As noted by others¹⁶, the Q_s peak actually coincides with periods of low uplift rate for all scenarios where forcing periodicity is less than the Q_s response time.

Similar to the behavior of the Q_s signal, the amplitude of P_v is damped and the peak timing of P_v becomes increasingly offset as the forcing periodicity decreases (Fig. 3a). However, unlike Q_s , the P_{vA} peak does coincide with the time of high uplift rate for scenarios with forcing periods $> 50\%$ of the response time (Fig. 3a).

Precipitation. Model results show comparatively less damping of the Q_s signal as precipitation forcing periodicity decreases (Fig. 3). For example, when forcing periodicity is 25% of response time, variability in Q_s is still $\sim 55\%$ of its maximum variability ($Q_{s_max} - Q_{s_min}$) achieved during the stepped forcing experiment. The behavior of the P_v signal in dynamic equilibrium under changing precipitation is similar to the general pattern observed under changing uplift rate; P_v amplitude is damped and peak P_v signal is increasingly offset from the timing of forcing




Figure 4. Illustration of how changes in P_v (proximal erosion) and Q_s are predicted to differentiate geomorphic response (knickpoint migration versus lowering of channel profile gradient) and environmental forcing (uplift rate versus precipitation).

onset (Fig. 3). The shape of the Q_s versus P_v response becomes increasingly rectilinear as forcing period decreases as a consequence of Q_s responding more rapidly than P_v to a change in precipitation rate (Fig. 3b).

Sensitivity Analysis. The results from our sensitivity analysis (Scenarios 1.2–1.15 and 2.2–2.15) demonstrate similar Q_s and P_v signal responses to the base case under a range of model parameters (Table DR1 and Supplemental Fig. 1). The only significant exception is where the erodibility of the distal sediment source (Source B) is greater than the proximal sediment source (Source A) (Scenarios 1.5 and 2.5). In this case, the P_v signal includes both a peak and trough following a stepped change in forcing (Supplemental Fig. 1).

Discussion

Our results show that Q_s and P_v can have a variable response to the same change in forcing mechanism (Fig. 2). For instance, a plot of Q_s versus $P_{v,A}$ under dynamic equilibrium reveals strongly non-linear trends between these two sedimentary signals (Fig. 3b and Supplemental Fig. 1). The paired $Q_s - P_v$ response is distinct between changes in uplift rate and precipitation, suggesting that these forcing mechanisms can be differentiated from each other (Figs 3, 4). Although an increase in uplift rate and precipitation will both produce an increase in Q_s , the former is predicted to produce an increase in proximal erosion whereas the latter will produce an increase in distal erosion (Fig. 4). Similarly, although both increasing uplift rate and decreasing precipitation will increase proximal erosion, they will produce opposite trends in Q_s (Figs 2, 4). Because $Q_s = UA$ at steady-state, a stepped change in uplift rate will produce a permanent change in Q_s , whereas a stepped change in precipitation will produce only a transient change in Q_s ^{15,17}. These findings are consistent with previous research based on analog modeling that has suggested that tectonic and climatic forcings can be differentiated by knowledge of both Q_s and mean catchment elevation¹⁷. Successful application of the paired $Q_s - P_v$ response (Fig. 4) in natural systems will likely depend on the extent to which (1) Q_s can be estimated¹⁸, and (2) distinct rock types exist in the erosional catchment area. Identification of relative increases in proximal or distal erosion will be facilitated in circumstances where any contrasts in bedrock type are normal to the catchment, such that distal (high elevation) sediment sources yield distinct P_v versus proximal (low elevation) sources.

Further differentiation of tectonic versus climatic forcing mechanisms may be possible via analysis of the relative timing between peak Q_s and P_v responses. The Q_s response to precipitation change is predicted to be more rapid than the P_v response, with the opposite pattern for a change in uplift rate. Thus, our model results suggest that P_v may be a more sensitive signal of the timing of change in uplift rate, whereas Q_s is more sensitive to the timing of a change in precipitation. Signal response time also depends on forcing type and sign (increase or decrease). An increase in precipitation results in a signal (Q_s and P_v) that peaks and then decreases very quickly with a typical response time of ~ 0.4 Myr. However, a decrease in precipitation results in a much longer signal response time (~ 0.9 Myr), and changes in uplift rate may produce signals that have typically longer response times (0.9–1.2 Myr).

The divergent responses of Q_s and P_v suggest differing geomorphic drivers behind these sedimentary signals. For instance, our modeling results predict that relative proximal erosion should increase in response to upstream knickpoint migration, a scenario that can occur in response to an increase in uplift rate or decrease in precipitation^{28,29}. An increase in both proximal erosion and Q_s has been associated with an increase in normal fault uplift rate in the Central Apennines of Italy¹⁰. In this case, the region of increased proximal erosion was associated with a zone of pronounced convexity downstream of the knickpoint, as predicted by our model results¹⁰ (Fig. 4). Our model further predicts that distal erosion is favored in circumstances where streams incise to form lower gradient profiles, such as occur during a decrease in uplift rate or increase in precipitation¹⁷ (Fig. 4).

Although the modeling framework presented herein provides a first-order prediction of how the paired Q_s – P_v signal responds to tectonic and climatic forcings (Figs 3, 4), there are a number of potential caveats to applying model results to natural landscapes. (1) The stream-power erosional model (Equation 2) used herein does not allow for landsliding, a process that is known to exert an important control on sediment load and signal propagation in natural systems³⁰. (2) Modeled precipitation changes does not account for variations in precipitation event intensity or frequency. For example, development of a drier, but stormier, climate may buffer the predicted Q_s and P_v response^{31,32}. (3) Variations in bedrock erodibility may influence the P_v response, particularly when distal sources are more easily eroded than proximal sources (e.g., scenarios 1.5 and 2.5; Supplemental Fig. 1) and possibly in layered stratigraphy³³. (4) P_v may also be sensitive to geomorphic processes not modeled herein, such as drainage divide migration or stream capture^{34,35}.

Our results suggest that integration of multiple sedimentary parameters may provide greater resolving power than any single parameter alone (e.g. ref.²⁰). Although Q_s , grain size, and now P_v , have been investigated in the context of sedimentary signal propagation, future efforts to characterize the relationship between geomorphic processes, their underlying physical mechanisms³⁶, and resulting sediment characteristics may permit multi-dimensional analysis of signal transmission and propagation. For example, climate signals are hypothesized to be manifested as changes in sediment composition, such as the chemical index of alteration³⁷, and there is opportunity for more research to elucidate how a sediment composition or weathering signal is transmitted from erosional landscapes undergoing changing forcing mechanisms. Thus, to the extent that sedimentary signals are produced by different geomorphic processes that operate at distinct, characteristic timescales, use of multiple sedimentary signals with distinct origins may greatly facilitate inversion of the stratigraphic record.

References

1. IPCC. *Summary for Policymakers. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, 10.1017/CBO9781107415324 (2014).
2. Castelltort, S. & Van Den Driessche, J. How plausible are high-frequency sediment supply-driven cycles in the stratigraphic record? *Sediment. Geol.* **157**, 3–13 (2003).
3. Jerolmack, D. J. & Paola, C. Shredding of environmental signals by sediment transport. *Geophys. Res. Lett.* **37**, 1–5 (2010).
4. Armitage, J. J., Dunkely Jones, T., Duller, R. A., Whittaker, A. C. & Allen, P. A. Temporal buffering of climate-driven sediment flux cycles by transient catchment response. *Earth Planet. Sci. Lett.* **369–370**, 200–210 (2013).
5. Romans, B. W., Castelltort, S., Covault, J. A., Fildani, A. & Walsh, J. P. Environmental signal propagation in sedimentary systems across timescales. *Earth-Science Rev.* **153**, 7–29 (2016).
6. Foreman, B. Z. & Straub, K. M. Autogenic geomorphic processes determine the resolution and fidelity of terrestrial paleoclimate records. *Sci. Adv.* **3** (2017).
7. Schilgden, T. F. *et al.* Landscape response to late Pleistocene climate change in NW Argentina: Sediment flux modulated by basin geometry and connectivity. *J. Geophys. Res. Earth Surf.* **121**, 392–414 (2016).
8. Tofelde, S. *et al.* 100 kyr fluvial cut-and-fill terrace cycles since the Middle Pleistocene in the southern Central Andes, NW Argentina. *Earth Planet. Sci. Lett.* **473**, 141–153 (2017).
9. Duller, R. A. *et al.* From grain size to tectonics. *J. Geophys. Res. Earth Surf.* **115**, 1–19 (2010).
10. Whittaker, A. C., Attal, M. & Allen, P. A. Characterising the origin, nature and fate of sediment exported from catchments perturbed by active tectonics. *Basin Res.* **22**, 809–828 (2010).
11. Whittaker, A. C. *et al.* Decoding downstream trends in stratigraphic grain size as a function of tectonic subsidence and sediment supply. *Bull. Geol. Soc. Am.* **123**, 1363–1382 (2011).
12. D'Arcy, M., Whittaker, A. C. & Roda-Boluda, D. C. Measuring alluvial fan sensitivity to past climate changes using a self-similarity approach to grain-size fining, Death Valley, California. *Sedimentology* **64**, 388–424 (2017).
13. Paola, C., Heller, P. L. & Angevine, C. L. The large-scale dynamics of grain-size variation in alluvial basins, 1: Theory. *Basin Res.* **4**, 73–90 (1992).
14. Allen, P. A. Time scales of tectonic landscapes and their sediment routing systems. *Geol. Soc. London, Spec. Publ.* **296**, 7–28 (2008).
15. Armitage, J. J., Duller, R. A., Whittaker, A. C. & Allen, P. A. Transformation of tectonic and climatic signals from source to sedimentary archive. *Nat. Geosci.* **4**, 231–235 (2011).
16. Li, Q., Gasparini, N. M. & Straub, K. M. Some signals are not the same as they appear: How do erosional landscapes transform tectonic history into sediment flux records? *Geology* 2–5, <https://doi.org/10.1130/G40026.1> (2018).
17. Bonnet, S. & Crave, A. Landscape response to climate change: Insights from experimental modeling and implications for tectonic versus climatic uplift of topography. *Geology* **31**, 123–126 (2003).
18. Allen, P. A. *et al.* The Q_s problem: Sediment volumetric balance of proximal foreland basin systems. *Sedimentology* **60**, 102–130 (2013).
19. Weltje, G. J. & von Eynatten, H. Quantitative provenance analysis of sediments: review and outlook. *Sediment. Geol.* **171**, 1–11 (2004).
20. Jonell, T. N., Owen, L. A., Carter, A., Schwenniger, J. L. & Clift, P. D. Quantifying episodic erosion and transient storage on the western margin of the Tibetan Plateau, upper Indus River. *Quat. Res. (United States)* **89**, 281–306 (2018).
21. Sharman, G. R. & Johnstone, S. A. Sediment unmixing using detrital geochronology. *Earth Planet. Sci. Lett.* **477** (2017).
22. Whipple, K. X. & Tucker, G. E. Dynamics of the stream-power river incision model: Implications for height limits of mountain ranges, landscape response timescales, and research needs. *J. Geophys. Res. Solid Earth* **104**, 17661–17674 (1999).
23. Whipple, K. X. Implications of sediment-flux-dependent river incision models for landscape evolution. *J. Geophys. Res.* **107**, 2039 (2002).
24. D'Arcy, M. & Whittaker, A. C. Geomorphic constraints on landscape sensitivity to climate in tectonically active areas. *Geomorphology* **204**, 366–381 (2014).
25. Whipple, K. X. Fluvial landscape response time: How plausible is steady-state denudation? *Am. J. Sci.* **301**, 313–325 (2001).

26. Hobbey, D. E. J. *et al.* Creative computing with Landlab: An open-source toolkit for building, coupling, and exploring two-dimensional numerical models of Earth-surface dynamics. *Earth Surf. Dyn.* **5**, 21–46 (2017).
27. Braun, J. & Willett, S. D. A very efficient O(n), implicit and parallel method to solve the stream power equation governing fluvial incision and landscape evolution. *Geomorphology* **180–181**, 170–179 (2013).
28. Anderson, R. S. & Anderson, S. P. *Geomorphology The Mechanics and Chemistry of Landscapes*. (Cambridge University Press, 2010).
29. Burbank, D. W. & Anderson, R. S. *Tectonic Geomorphology*. (Blackwell Publishing, 2012).
30. Roda-Boluda, D. C., D'Arcy, M., McDonald, J. & Whittaker, A. C. Lithological controls on hillslope sediment supply: insights from landslide activity and grain size distributions. *Earth Surf. Process. Landforms* **43**, 956–977 (2018).
31. Miller, D. M. *et al.* Holocene landscape response to seasonality of storms in the Mojave Desert. *Quat. Int.* **215**, 45–61 (2010).
32. D'Arcy, M., Roda-Boluda, D. C. & Whittaker, A. C. Glacial-interglacial climate changes recorded by debris flow fan deposits, Owens Valley, California. *Quat. Sci. Rev.* **169**, 288–311 (2017).
33. Forte, A. M., Yanites, B. J. & Whipple, K. X. Complexities of Landscape Evolution During Incision Through Layered Stratigraphy with Contrasts in Rock Strength. *Earth Surf. Process. Landforms* n/a–n/a, <https://doi.org/10.1002/esp.3947> (2016).
34. Whipple, K. X., Forte, A. M., DiBiase, R. A., Gasparini, N. M. & Ouimet, W. B. Timescales of landscape response to divide migration and drainage capture: Implications for the role of divide mobility in landscape evolution. *J. Geophys. Res. Earth Surf.* **122**, 248–273 (2017).
35. Pelletier, J. D. Persistent drainage migration in a numerical landscape evolution model. *Geophys. Res. Lett.* **31**, 4–7 (2004).
36. Dietrich, W. E. *et al.* Geomorphic transport laws for predicting landscape form and dynamics. *Geophys. Monogr. Ser.* **135**, 103–132 (2003).
37. Hessler, A. M., Zhang, J., Covault, J. A. & Ambrose, W. Continental weathering coupled to Paleogene climate changes in North America. *Geology* **45**, 1–4 (2017).

Acknowledgements

We thank the sponsors of the Quantitative Clastics Laboratory (<http://www.beg.utexas.edu/qcl>) for sponsoring this research and the three reviewers that provided constructive feedback in peer-review.

Author Contributions

All authors designed the study. G.R. Sharman and Z. Sylvester conducted the analysis. G.R. Sharman led efforts in writing the manuscript, with J.A. Covault and Z. Sylvester contributing to reviewing and editing.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-019-39754-6>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019