

# SCIENTIFIC REPORTS

**OPEN**

## **Author Correction:** A randomized controlled trial of an ambulatory approach versus the hospital-based approach in managing suspected obstructive sleep apnea syndrome

David S. Hui<sup>1,2</sup>, Susanna S. Ng<sup>1</sup>, Kin-Wang To<sup>1</sup>, Fanny W. Ko<sup>1</sup>, Jenny Ngai<sup>1</sup>, Ken K. P. Chan<sup>1</sup>, Wing-Ho Yip<sup>1</sup>, Tat-On Chan<sup>1,2</sup>, Karen Yiu<sup>1,2</sup> & Wilson W. S. Tam<sup>3</sup>

Correction to: *Scientific Reports* <https://doi.org/10.1038/srep45901>, published online 04 April 2017

This Article contains errors in Table 4 where the values for ‘% of patients who used CPAP  $\geq$ 4 hrs per night [n(%))]’ and ‘% of patients who used CPAP  $\geq$ 70% of nights,  $\geq$ 4 hrs per night [n(%))]’ are incorrect in the ‘Home’ and ‘Hospital’ columns.

In addition, ‘Home (n = 62)’ should read ‘Home (n = 62, 72.1% of 86 patients accepted CPAP)’, and ‘Hospital (n = 69)’ should read ‘Hospital (n = 69, 80.2% of 86 patients accepted CPAP)\*’.

The correct Table 4 appears below as Table 1, accompanied by a footnote which gives details of the CPAP acceptance.

<sup>1</sup>Dept of Medicine & Therapeutics, The Chinese University of Hong Kong, Hong Kong, China. <sup>2</sup>SH Ho Sleep Apnoea Management Center, The Chinese University of Hong Kong, Hong Kong, China. <sup>3</sup>Alice Lee Centre for Nursing Studies, National University of Singapore, Singapore, Singapore. Correspondence and requests for materials should be addressed to D.S.H. (email: [dschui@cuhk.edu.hk](mailto:dschui@cuhk.edu.hk))

	Home (n=62, 72.1% of 86 patients accepted CPAP)	Hospital (n=69, 80.2% of 86 patients accepted CPAP)*	p-value
CPAP titration pressure (cmH <sub>2</sub> O)	13.3 (2.2)	12.8 (2.1)	0.252
CPAP usage at 3 months	5.0 (2.0)	3.9 (2.1)	<b>0.005</b>
% of patients who used CPAP $\geq$ 4 hrs per night [n(%)]	46 (74.2%)	39 (56.5%)	<b>0.033</b>
% of patients who used CPAP $\geq$ 70% of nights, $\geq$ 4 hrs per night [n(%)]	42 (67.7%)	37 (53.6%)	0.123
Time from 1st consultation to diagnostic sleep test (days)	54.2 (31.3)	258.9 (186.9)	<b>&lt;0.001</b>
Time from 1st consultation to CPAP titration (days)	109.0 (36.8)	274.0 (193.6)	<b>&lt;0.001</b>
Time from 1st consultation to CPAP treatment (days)	154.7 (69.5)	299.7 (198.7)	<b>&lt;0.001</b>

**Table 1.** CPAP outcomes & time to investigation and treatment (CPAP users only, n = 62 vs 69, TPP analysis). Footnote: \*62 out of 86 versus 69 out of 86 patients with AHI  $\geq$  15/hr in the home management group and hospital-based management group respectively accepted CPAP treatment and completed 3 months of follow up. No group difference on the rate of CPAP acceptance was found, P-value = 0.213.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019