

OPEN

Hadron production models' prediction for p_T distribution of charged hadrons in pp interactions at 7 TeV

S. Ullah¹, M. Ajaz¹, Z. Wazir², Y. Ali³, K. H. Khan⁴ & H. Younis³

The yield of transverse momentum (p_T) spectra of integrated hadrons and their ratios produced in pp collisions at 7 TeV are reported using DPMJET-III, EPOS1.99, EPOS-LHC, HIJING1.383, QGSJETII-04, and Sibyll2.3c models. The models' predictions are compared with the ALICE measurements obtained at mid rapidity and in the p_T range from 0.3–6 GeV/c, 0.2–6 GeV/c, and 0.1–3 GeV/c for protons, kaons and pions respectively. Compared to the experimental data, the EPOS1.99 and EPOS-LHC models predict the yield of pion very well while reproducing yields of kaon and proton at low p_T only. The DPMJET-III model predicts the yield of pions, kaons and protons only at intermediate value of p_T . The HIJING and Sibyll models describe the integrated yield of pion well at high p_T only. The QGSJETII model predicts the yield of kaon for almost the whole p_T range. EPOS1.99 and Sibyll models reproduce excellent prediction of the K/π ratio over the entire range of p_T whereas the EPOS-LHC, HIJING and DPMJET models underpredict with increasing order of discrepancy respectively. The EPOS1.99, EPOS-LHC, and Sibyll models predict the p/π ratio for low values of p_T , but overpredict at high values, while the DPMJET-III and HIJING models underpredict over the entire range of p_T . Although the models' predictions are mostly consistent with the ALICE measurements but none of the models completely describe the entire distribution for the integrated yield of hadron and their ratios.

The measurements of hadron production have an extended record in high energy nuclear and particle physics^{1–13}. In order to test soft parton interactions and hadronization processes predicted by non-perturbative quantum chromodynamics, the transverse momentum (p_T) spectra and absolute yields of hadrons are used by Monte Carlo (MC) event generators¹⁴. We get important information on multi-parton interactions and other final state effects from the dependence of the yield and p_T distribution on the hardness in pp collisions. The identified hadron distributions in pp collisions also provide an important reference to the findings of high-energy heavy-ion analysis, where the spectral shape and yield of hadron species are modified by the final state effect. The interactions responsible for pp collisions and subsequent hadronization process at the LHC can be understood using the quantum chromodynamics¹⁵. Phenomenological models must be employed, as precise calculations are difficult to perform in the non-perturbative regime. Monte Carlo event generators based on these models need to be tuned, to reproduce experimental observables.

The present analysis is focused on the study of p_T spectra of the integrated yield of π ($\pi^+ + \pi^-$), K ($K^+ + K^-$), proton (proton + anti-proton) production, and the K/π and p/π ratios in proton-proton (pp) interactions at 7 TeV using DPMJET-III, HIJING1.383, EPOS, QGSJETII-04, and Sibyll2.3c models, and their comparison with the data from the ALICE experiment¹¹. The analysis, in future, will be extended to heavy ion collision for the study of strongly interacting matter and to search for a signal on the onset of deconfinement.

¹Department of Physics, Abdul Wali Khan University Mardan, Mardan, 23200, Pakistan. ²Department of Physics, International Islamic University Islamabad, Islamabad, 44000, Pakistan. ³Department of Physics, COMSATS University, Islamabad, 44000, Pakistan. ⁴Department of Physics, Women University of Azad Jammu and Kashmir, Bagh, Pakistan. Correspondence and requests for materials should be addressed to M.A. (email: ajaz@awkum.edu.pk)

The Method and Models

p_T distributions of differential yields of integrated hadrons (sum of positively and negatively charged hadron) and their ratios, normalized by the total number of inelastic events, simulated for pp interactions at 7 TeV in the p_T range from 0.1–3 GeV/ c for pions, from 0.2–6 GeV/ c for kaons, and from 0.3–6 GeV/ c for protons with $0 < y < 0.5$ are presented here. Model simulations of 1M events using DPMJET, HIJING1.383, EPOS1.99, EPOS-LHC, QGSJETII-04 and Sibyll2.3c are compared with the measurements of experimental data from the ALICE experiment¹¹. Brief description of the models is given below while references given against each are referred for complete description of each model.

DPMJET-III¹⁶ is a Monte Carlo code constructed on the Dual Parton Model¹⁷ combining the features of DPMJET-II¹⁸ Dtunuc-2^{19,20}, and Phojet1.12²¹. DPMJET-III is a multipurpose code that allows for the simulation of photon-photon, photon-hadron, photon-nucleus, hadron-hadron (hh), hadron-nucleus (hA), and nucleus-nucleus (AA) interactions from a few GeV up to the highest energies cosmic rays¹⁶. In our analysis, we used DPMJETIII.17-1, which is an updated version of DPMJET-2.55 where few parameters were changed to reproduce TOTEM cross section²². The model is the Tevatron version while the new LHC version is not public yet.

HIJING is a hadron production model based on Monte Carlo simulation that allows high energy hh , hA , and AA collisions²³. The model is aimed particularly to study jets and related particle production based on QCD-inspired models. HIJING includes mechanisms such as soft excitation, production of multiple minijets, interactions of jets in dense hadronic matter, and nuclear shadowing of parton distribution functions²⁴. In our simulation, we used HIJING 1.383 with default parameters. The model is using Pythia 5.3 to generate kinetic variables and Jetset 7.2 for jet fragmentation.

EPOS is a hadronic interaction Monte Carlo event generator used for minimum bias interactions of heavy ion as well as cosmic ray air shower simulations^{25,26}. EPOS is also used for centrality dependent heavy-ion collision. Energy conservation is used by EPOS model at amplitude level. The EPOS1.99 model is tuned to data up to Tevatron energies. An updated version of EPOS for Large Hadron Collider (LHC) energies, is EPOS-LHC, where a different parametrization of flow has been introduced in pp collisions compared to heavy-ion producing large volume. EPOS-LHC model is tuned to LHC data up to 8 TeV but the 13 TeV data tune is still missing in it. The model, utilizing colour exchange mechanism of string excitation, is tuned to LHC data recorded in 2012. It is based on the Gribov Regge approach where the same flow parameterization is used as is used in EPOS1.99. EPOS-LHC can reproduce the minimum bias results for all those particles that are having transverse momentum, ranging from 0 to a few GeV/ c ^{27,28}. Beside EPOS-LHC, there is another parallel development of the EPOS model named EPOS 3, based on the same approach as EPOS1.99, used for collectivity in heavy ion collision.

QGSJET (Quark Gluon String model with JETs) is a Monte Carlo generator widely used in many collaborations, and relies on the physics picture of the Quark-Gluon String model²⁹. The Gribov's Reggeon calculus approach is used in the QGSJET model for hadronic and nuclear collisions as multiple scattering processes^{30,31}. Like EPOS-LHC, QGSJET is also tuned to several accelerator data including LHC data up to 8 TeV. No energy conservation in Sibyll and QGSJET is used for cross section calculations. Sibyll is a fast Monte Carlo event generator used for air shower simulations³². The model is able to describe features of hadronic particle production. It is a QCD inspired model that includes multiple interactions, hard as well a soft scattering, string fragmentation and saturation effects. Sibyll 2.1³³, that was tuned to TEVATRON data is updated to Sibyll 2.3c³⁴, to describe the NA49 data. The new version further includes charm production, baryon pair production and beam remnants. The model is tuned to NA61 data and to LHC data at 8 TeV.

The analysis is performed using the RIVET Monte Carlo analysis tool³⁵, which has a big code library to compare the results available on HEPDATA with the Monte Carlo event generators. It is based on the HepMC for the universal event record format³⁶, on the library of LHAPDF for parton densities^{37,38}, and on the Fastjet package for jet clustering^{39,40}.

Results and Discussion

The transverse momentum distribution of the integrated charged particles, π ($\pi^+ + \pi^-$), K ($K^+ + K^-$), proton (proton + anti-proton), from left to right respectively, normalized to the number of inelastic interactions are given in Fig. 1. Referring to legends in the figures, experimental data¹¹ is shown by a filled circle, while lines with different colours are used to represent model predictions. Solid red line is used to represent the prediction of DPMJET-III model, blue for EPOS1.99, green for EPOS-LHC, yellow for HIJING, purple for QGSJET while dash red line is used for Sibyll. Comparing to the experimental data, DPMJET-III model predicts the yield of pions, kaons and protons for $1.2 < p_T < 1.7$ GeV/ c , $3.3 < p_T < 5.2$, and $2.1 < p_T < 3.1$ respectively within 10%, 15% and 15%. The model overestimates the production of pions for $p_T < 1.2$ GeV/ c and $p_T > 1.7$ GeV/ c , underestimates for kaons in the region other than $3.3 < p_T < 5.2$ GeV/ c , and for protons it predicts lower yield at $p_T < 2.1$ GeV/ c and higher yield at $p_T > 3.1$ GeV/ c . EPOS1.99 model describes the p_T distribution of the integrated yield of pion within 5%, particularly at low values of p_T . The model reproduces the kaon and protons spectra only at low p_T , which is within 20% for kaon and 5% for proton in $1 < p_T < 2$ GeV/ c region. The model underestimates and overestimates the spectra for kaon and proton respectively in data at high p_T . The EPOS-LHC model predicts yield of pion within 20% at low p_T and within 10% at high p_T while predicts yields of kaon and protons only at low p_T . EPOS-LHC model underpredicts and overpredicts respectively yield of kaon and protons at high p_T . The HIJING model underpredicts the integrated yield of pion at low p_T , but describes the distribution within 10% at high p_T . The HIJING model underpredicts and overpredicts the kaons and protons yield respectively. The QGSJETII model gives higher yield of the pion as compared to the experimental data. The model predicts the kaon yield for $2.5 < p_T < 4.5$ GeV/ c , while describes protons yield within the uncertainty limit for $0.5 < p_T < 2.5$ GeV/ c but underestimates the data at high p_T . Sibyll model overestimates (20%) the pion yield at low p_T , but produce the




Figure 1. Transverse momentum spectra of the integrated yield of pion, kaons and protons by DPMJET-III, EPOS1.99, HIJING1.383, EPOS-LHC, QGSJETII-04 and Sibyll2.3c in pp collisions at 7 TeV are compared with the measurements of experimental data from the ALICE experiment. Experimental data is shown by filled circle while lines with different colours are used to represent models' predictions. Solid red line is used to represent the prediction of DPMJET, blue for EPOS1.99, green for EPOS-LHC, yellow for HIJING, purple for QGSJET while dash red line is used for Sibyll.




Figure 2. Ratio of the integrated yield of transverse momentum spectra of kaons to pions (left) and protons to pions (right) by DPMJET-III, EPOS1.99, HIJING1.383, EPOS-LHC, QGSJETII-04 and Sibyll2.3c in pp collisions at 7 TeV are compared with the measurements of experimental data from the ALICE experiment. Experimental data is shown by filled circle while lines with different colours or texture are used to represent models' predictions as shown in legend.

results within 5% at high p_T . The model describes Kaon distribution within 10% for $p_T < 1.2$ GeV/c but underestimates at high p_T while overestimating yield of proton over the entire p_T range.

The ratios of the K yield relative to π ($(K^+ + K^-)/(\pi^+ + \pi^-)$), and p yield relative to π ($(p + p)/(\pi^+ + \pi^-)$) are shown in Fig. 2. DPMJET-III model underpredicts both K/π and p/π ratios. EPOS1.99 model produces good prediction within 5% of the K/π ratio over the entire range of p_T , but could predict only $1.2 < p_T < 1.7$ of p/π ratio while overestimate the rest of the p_T region. EPOS-LHC model underpredicts the K/π ratio within 15%. The EPOS-LHC model reproduces p/π ratio within 10% at $p_T < 1.5$ GeV/c, but overestimates at levels beyond 1.5 GeV/c. The HIJING model underpredicts (within 20% at low and 30% at high p_T) the K/π ratio but overshoots the p/π ratio. QGSJETII model underpredicts both the K/π and p/π ratios. The Sibyll model predicts the K/π ratio very well within the uncertainty limit up to $p_T \sim 2$ GeV/c and within 10% at high p_T . The model reproduces within 20% the p/π ratio for $p_T < 1$ GeV/c but overshoots the ratio afterwards.

Measurements performed in¹¹ are compared with three different tunes of Pythia^{41–43}, EPOS-LHC²⁵ and PHOJET⁴⁴ models. Their¹¹ conclusion is summarized as follows. EPOS prediction for p_T spectrum of pion is reported to be within 15% with highest deviation at 1.2 GeV/c, which is more prominent for kaons and protons than pions. All three tunes of the Pythia give similar predictions for pions as given by EPOS. No one tune predicts the shape of the proton spectrum in the full p_T range while gives a good prediction for the yield in the range $1 < p_T < 2$ GeV/c, but they overshoot the measurements at lower and higher p_T by up to 40%. The PHOJET model

fails to describes the spectrum for all the particle species. All models overshoot the kaon spectra. All models undervalue the experimental data at high momenta for the ratios K/π versus p_T with EPOS showing closest prediction. Pythia6-Z2 and Pythia8 provide good prediction of the p/π ratio, while all other show a large deviation at high momenta.

Conclusion

The study compares the predictions of the transverse momentum distribution of charged particles and their ratios obtained using DPMJET-III, HIJING1.383, EPOS-LHC, QGSJETII-04 and Sibyll2.3c with the ALICE measurements obtained at mid rapidity. DPMJET-III model predicts the yield of pion, kaons and protons at mid value of p_T only. The EPOS1.99 and EPOS-LHC models predict the yield of pion very well while predict the yield of kaon and proton only at low p_T . The HIJING model describes yield of pion well only at high p_T . The QGSJETII model predicts the kaon and proton yield within the uncertainty limit of experimental data for $2.5 < p_T < 4.5$ and $0.5 < p_T < 2.5$ respectively. Sibyll model reproduces the yield of pion at high p_T but overestimate the results at low p_T . EPOS1.99 and Sibyll models with in 5% limit reproduce the K/π ratio over the entire range of p_T whereas the EPOS-LHC, HIJING and DPMJET model underpredict the ratio with increasing order of discrepancy respectively. The EPOS1.99, EPOS-LHC and Sibyll2.3c models predict the p/π ratio for low values of p_T whereas the DPMJET-III and HIJING models underpredict over the entire range of p_T . Although the models' predictions are mostly compatible with the ALICE measurements but no model completely describe all the distributions.

References

1. Abgrall, N. *et al.* Measurements of cross sections and charged pion spectra in proton-carbon interactions at 31 GeV/c. *Phys. Rev. C* **84**, 034604 (2011).
2. Abgrall, N. *et al.* Measurements of π^\pm , K^\pm , K_s^0 , Λ and proton production in proton-carbon interactions at 31 GeV/c with the NA61/SHINE spectrometer at the CERN SPS. *Eur. Phys. J. C* **76**, 84 (2016).
3. Nigmatov, T. S. *et al.* Forward neutron production at the Fermilab Main Injector. *Phys. Rev. D* **83**, 012002 (2011).
4. Abgrall, N. *et al.* Measurements of π^\pm , K^\pm and proton double differential yields from the surface of the T2K replica target for incoming 31 GeV/c protons with the NA61/SHINE spectrometer at the CERN SPS. *Eur. Phys. J. C* **79**, 100 (2019).
5. Ullah, S., Ali, Y., Ajaz, M., Tabassam, U. & Ali, Q. π^\pm , K^\pm , protons and antiprotons production in proton-carbon interactions at 31 GeV/c using hadron production models. *Int. J. Mod. Phys. A* **33**, 1850108 (2018).
6. Ullah, S., Ajaz, M. & Ali, Y. Spectra of strange hadrons and their role in neutrinos flux prediction. *EPL* **123**, 31001 (2018).
7. Adare, A. *et al.* Identified charged hadron production in p+p collisions at $\sqrt{s} = 200$ and 62.4 GeV. *Phys. Rev. C* **83**, 064903 (2011).
8. Alt, C. *et al.* Inclusive production of charged pions in p+p collisions at 158 GeV/c beam momentum. *Eur. Phys. J. C* **45**, 343 (2006).
9. Ajaz, M., Bilal, M., Ali, Y., Suleymanov, M. K. & Khan, K. H. Models prediction of hadrons production ratios in pp collisions at $\sqrt{s} = 7$ TeV. *Mod. Phys. Lett. A* **34**, 1950090 (2019).
10. Ajaz, M., Tufail, M. & Ali, Y. Production of light flavored charged hadron in pp collisions at 900 GeV with hadron production models. *Mod. Phys. Lett. A* **34**, 1950100 (2019).
11. Adam, J. *et al.* Measurement of pion, kaon and proton production in proton-proton collisions at $\sqrt{s} = 7$ TeV. *Eur. Phys. J. C* **75**, 226 (2015).
12. Abelev, B. *et al.* Production of charged pions, kaons and protons at large transverse momenta in pp and Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. *Phys. Lett. B* **736**, 196 (2014).
13. Aaij, R. *et al.* Measurement of prompt hadron production ratios in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV. *Eur. Phys. J. C* **72**, 2168 (2012).
14. Chatrchyan, S. *et al.* Study of the inclusive production of charged pions, kaons, and protons in pp collisions at $\sqrt{s} = 0.9, 2.76$, and 7 TeV. *Eur. Phys. J. C* **72**, 2164 (2012).
15. Ellis, R. K., Stirling, W. J. & Webber, B. R. QCD and Collider Physics, Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology, Volume 8 (Cambridge University Press, 1996).
16. Roesler, S. *et al.* *Advanced Monte Carlo for Radiation Physics, Particle Transport Simulation and Applications*. (Springer, Berlin, Heidelberg, 2001).
17. Capella, A. *et al.* Dual parton model. *Phys. Rept* **236**, 225 (1994).
18. Ranft, J. The dual parton model at cosmic ray energies. *Phys. Rev. D* **51**, 64 (1995).
19. Engel, R., Ranft, J. & Roesler, S. Photoproduction off nuclei and pointlike photon interactions. I. Cross sections and nuclear shadowing. *Phys. Rev. D* **55**, 6957 (1997).
20. Roesler, S., Engel, R. & Ranft, J. Photoproduction off nuclei: Particle and jet production. *Phys. Rev. D* **57**, 2889 (1998).
21. Engel, R. & Ranft, J. Hadronic photon-photon interactions at high energies. *Phys. Rev. D* **54**, 4244 (1996).
22. Csorgo, T. *et al.* Elastic Scattering and Total Cross-Section in p+p reactions measured by the LHC Experiment TOTEM at $\sqrt{s} = 7$ TeV. *Prog. Theor. Phys. Suppl.* **193**, 180–183 (2012).
23. Wang, X.-N. Role of multiple minijets in high-energy hadronic reactions. *Phys. Rev. D* **43**, 104 (1991).
24. Wang, X.-N. & Gyulassy, M. HIJING: A Monte Carlo model for multiple jet production in pp, pA, and AA collisions. *Phys. Rev. D* **44**, 3501 (1991).
25. Werner, K. *et al.* Parton ladder splitting and the rapidity dependence of transverse momentum spectra in deuteron-gold collisions at the BNL Relativistic Heavy Ion Collider. *Phys. Rev. C* **74**, 044902 (2006).
26. Pierog, T. & Werner, K. EPOS Model and Ultra High Energy Cosmic Rays. *Nucl. Phys. Proc. Suppl.* **196**, 102 (2009).
27. Pierog, T. *et al.* EPOS LHC: Test of collective hadronization with data measured at the CERN Large Hadron Collider. *Phys. Rev. C* **92**, 034906 (2015).
28. Pierog, T. & Werner, K. Hadronic Interaction Model EPOS and Air Shower Simulations: New Results on Muon Production. Proceedings of the 30th International Cosmic Ray Conference, Mexico **4**, 629–632 (2008).
29. Ostapchenko, S. Monte Carlo treatment of hadronic interactions in enhanced Pomeron scheme: QGSJET-II model. *Phys. Rev. D* **83**, 014018 (2011).
30. Gribov, N. A Reggeon Diagram Technique. *Sov. Phys. JETP* **26**, 414 (1968).
31. Gribov, V. N. Glauber corrections and the interaction between high-energy hadrons and nuclei. *Sov. Phys. JETP* **29**, 483 (1969).
32. Fletcher, R. S., Gaisser, T. K., Lipari, P. & Stanev, T. SIBYLL: An event generator for simulation of high energy cosmic ray cascades. *Phys. Rev. D* **50**, 5710 (1994).
33. Ahn, E.-J., Engel, R., Gaisser, T. K., Lipari, P. & Stanev, T. Cosmic ray interaction event generator SIBYLL 2.1. *Phys. Rev. D* **80**, 094003 (2009).
34. Riehn, F. *et al.* The hadronic interaction model Sibyll 2.3c and Feynman scaling. *PoS ICRC2017*, 301 (2017).
35. Buckley, A. *et al.* Rivet user manual. *Comput. Phys. Commun.* **184**, 2803–2819 (2013).
36. Dobbbs, M. & Hansen, J. B. The HepMC C++ Monte Carlo event record for High Energy Physics. *Comput. Phys. Commun.* **134**, 41–46 (2001).

37. Whalley, M. R., Bourilkov, D. & Group, R. C. The Les Houches accord PDFs (LHAPDF) and lhapglue. arXiv:hep-ph/0508110 (2005).
38. Bourilkov D., Group, R. C. & Whalley, M. R. LHAPDF: PDF use from the Tevatron to the LHC. arXiv:hep-ph/0605240 (2006).
39. Cacciari, M. & Salam, G. P. Dispelling the N^3 myth for the k_T jet-finder. *Phys. Lett. B* **641**, 57–61 (2006).
40. Cacciari, M. *et al.* FastJet user manual. *Eur. Phys. J. C* **72**, 1896 (2012).
41. Skands, P. Z. Tuning Monte Carlo generators: The Perugia tunes. *Phys. Rev. D* **82**, 074018 (2010).
42. Sjöstrand, T., Mrenna, S. & Skands, P. Z. A Brief Introduction to PYTHIA 8.1. *Comput. Phys. Commun.* **178**, 852–867 (2008).
43. Corke, R. & Sjöstrand, T. Multiparton Interactions with an x-dependent Proton Size. *JHEP* **1105**, 009 (2011).
44. Engel, R., Ranft, J. & Roesler, S. Hard diffraction in hadron-hadron interactions and in photoproduction. *Phys. Rev. D* **52**, 1459–1468 (1995).

Acknowledgements

This work is supported by the Higher Education Commission (HEC) of Pakistan by the Grant No. 20-3379/NRPU/R&D/HEC/2014.

Author Contributions

Mr. Shaid Ullah, Dr. Muhammad Ajaz and Dr. Yasir Ali. These authors contributed equally to this work. These authors wrote the main manuscript and carried out the research work. Dr. Muhammad Ajaz, Dr. Zafar Wazir, Dr. Kamal Hussain Khan and Dr. Hannan Younis. These authors jointly supervised this work. Draw all the figures and proofread the manuscript.

Additional Information

Competing Interests: Mr. Shahid Ullah work has been funded by Dr. Muhammad Ajaz in his project received from Higher Education Commission of Pakistan under National Research Program for Universities. Dr Yasir Ali, Dr. Zafar Wazir, Dr. Kamal Hussain Khan and Dr. Hannan Younis declare no potential conflict of interest.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019