SCIENTIFIC REPORTS

natureresearch

Corrected: Publisher Correction

OPEN

TCF7L2 polymorphisms are associated with amygdalar volume in elderly individuals with Type 2 Diabetes

Ithamar Ganmore^{1,2,4,5*}, Abigail Livny^{2,3,5}, Ramit Ravona-Springer^{2,4,5}, Itzik Cooper², Anna Alkelai⁶, Shahar Shelly^{1,2,7}, Galia Tsarfaty^{3,5}, Anthony Heymann^{5,8}, Michal Schnaider Beeri^{2,9} & Lior Greenbaum^{2,5,10}

The association between several Single Nucleotide Polymorphisms (SNPs) within the transcription factor 7-like 2 (TCF7L2) gene and Type 2 Diabetes (T2D) as well as additional T2D-related traits is well established. Since alteration in total and regional brain volumes are consistent findings among T2D individuals, we studied the association of four T2D susceptibility SNPS within TCF7L2 (rs7901695, rs7903146, rs11196205, and rs12255372) with volumes of white matter hyperintensities (WMH), gray matter, and regional volumes of amygdala and hippocampus obtained from structural MRI among 191 T2D elderly Jewish individuals. Under recessive genetic model (controlling for age, sex and intracranial volume), we found that for all four SNPs, carriers of two copies of the T2D risk allele (homozygous genotype) had significantly smaller amygdalar volume: rs7901695- CC genotype vs. CT + TT genotypes, p = 0.002; rs7903146-TT vs. TC + CC, p = 0.003; rs11196205- CC vs. CG + GG, p = 0.0003; and rs12255372-TT vs. TG + GG, p = 0.003. Adjusting also for T2D-related covariates, body mass index (BMI), and ancestry did not change the results substantively (rs7901695, p = 0.003; rs7903146, p = 0.005; rs11196205, p = 0.001; and rs12255372, p = 0.005). Conditional analysis demonstrated that only rs11196205 was independently associated with amygdalar volume at a significant level. Separate analysis of left and right amygdala revealed stronger results for left amygdalar volume. Taken together, we report association of TCF7L2 SNPs with amygdalar volume among T2D elderly Jewish patients. Further studies in other populations are required to support these findings and reach more definitive conclusions.

Type 2 diabetes (T2D) is a multifactorial disease, with a complex polygenic architecture^{1–3}. The association of the transcription factor 7-like 2 (*TCF7L2*) gene (chromosome 10q25.2-q25.3) with T2D is one of the most reproducible and robust finding in T2D genetics, as supported by Genome-wide association studies (GWAS), multiple replication studies and meta-analyses^{4,5}. Several single nucleotide polymorphisms (SNPs) within *TCF7L2* were independently associated with T2D susceptibility and related traits (e.g. insulin secretion and blood glucose levels)^{4–8}.

The protein encoded by *TCF7L2* gene is a transcription factor, involved in the Wnt/beta-catenin signaling pathway, which plays a role in cell proliferation and differentiation^{9,10}. It is related to beta-cells and other pancreatic cells functions¹¹⁻¹³, as well as to the development and function of adipose tissue¹⁴. In addition, *TCF7L2* is expressed in multiple brain regions and characterized by existence of several alternative splicing variants in

¹Department of Neurology, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel. ²The Joseph Sagol Neuroscience Center, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel. ³Department of Diagnostic Imaging, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel. ⁴Memory clinic, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel. ⁵Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel. ⁶Institute for Genomic Medicine, Columbia University, New York, NY, USA. ⁷Department of Neurology, Mayo Clinic Rochester, Minnesota, USA. ⁸Maccabi Healthcare Services, Tel Aviv, Israel. ⁹Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ¹⁰The Danek Gertner Institute of Human Genetics, Sheba Medical Center, Tel Hashomer, Ramat Gan, Israel. *email: itamargan@gmail.com

Characteristics	Values
Sex (males/females, N [%])	118/73 (61.8%/38.2%)
Age at IDCD study baseline (Mean \pm SD, years)	70.79 ± 4.22
Follow up in the registry (Mean \pm SD, years)	9.33 ± 4.58
HbA1c (Mean \pm SD, %)	6.66 ± 0.76
BMI (Mean \pm SD, kg/m ²)	28.37 ± 4.32
Systolic blood pressure (Mean \pm SD, mmHg)	133.05 ± 9.32
Diastolic blood pressure (Mean \pm SD, mmHg)	76.77 ± 4.94
T2D medications consumption (yes/no, N [%])	164/27 (85.9%/14.1%)
Ashkenazi origin (yes/no, N [%])	127/64 (66.5%/33.5%)

Table 1. Demographic and clinical description of the analyzed sample (N = 191). Abbreviations: N- numberof subjects; SD- standard deviation; BMI- body mass index; HbA1c- hemoglobin A1c; T2D- type 2 diabetes;IDCD- the Israel Diabetes and Cognitive Decline.

SNP	Position	Location	Minor, Major alleles	MAF	T2D risk allele
rs7901695	114754088	Intron 4	С, Т	0.403	С
rs7903146	114758349	Intron 4	Т, С	0.388	Т
rs11196205	114807047	Intron 5	G, C	0.487	С
rs12255372	114808902	Intron 5	T, G	0.382	Т

Table 2. TCF7L2 SNPs included in the analysis. SNPs' position refers to GRCh37.p13. Abbreviations: SNP-single nucleotide polymorphism; MAF-minor allele frequency; T2D- type 2 diabetes.

different species¹⁵⁻¹⁸. In humans, a unique splice variant was found in the brain, pancreatic islets and gut and therefore named the "neuroendocrine form"¹⁹. SNPs in *TCF7L2* have been associated with psychiatric disorders, such as schizophrenia and bipolar disorder²⁰⁻²².

Individuals with T2D have higher risk for cognitive dysfunction and dementia than those without T2D²³. The most consistent findings in neuroimaging of T2D patients are higher number of infarcts and white matter lesions, as well as general cerebral and hippocampal atrophy^{24–26}. Few studies have found regional atrophy in different brain structures^{27–29}, but these results were not consistent. Greater atrophy of the amygdala and hippocampus had been associated not only with T2D³⁰, but also to high plasma glucose levels within the normal range³¹. However, the underlying etiology of brain volume differences in T2D is still unknown. At the genetic level, previous studies in the Israel Diabetes and Cognitive Decline (IDCD) study found that apolipoprotein $\varepsilon 4$ (*APOE4*) genotype³² and haptoglobin (Hp) 1–1 genotype³³, affect the relationship between neuroimaging phenotypes (White matter hyperintensities [WMHs] and Hippocampal volume, respectively) and glycemic control among T2D patients.

We examined the association of four *TCF7L2* SNPs (rs7901695, rs7903146, rs11196205, and rs12255372) with WMH, gray matter, and regional volumes of the amygdala and hippocampus obtained from structural brain magnetic resonance imaging (MRI) in Jewish T2D elderly patients. These SNPs are reported in the literature as robustly associated with T2D^{4,5}. We hypothesized that *TCF7L2* risk alleles for T2D would be associated with smaller regional brain volume and larger WMH volume among T2D patients.

Results

Demographic and medical characteristics. The final analysis included 191 T2D individuals, all of them were IDCD participants. The demographic and clinical description of the sample is detailed in Table 1. All participants were both genotyped for the *TCF7L2* SNPs (rs7901695, rs7903146, rs11196205, and rs12255372; Table 2) and had brain MRI scans. Genotyping success rate per SNP was 97.38–100%

Association of *TCF7L2* SNPs with the volume of different brain regions. We found that two SNPs, rs7903146 and rs7901695 were highly correlated to each other ($r^2 = 0.94$) and can be viewed essentially as a single signal (Table 3). From the analyzed SNPs, only rs7903146 showed slight deviation form Hardy-Weinberg equilibrium (p = 0.045), but nevertheless we included it in the analysis.

The mean volumes of the different brain regions, according to the four analyzed SNPs and genotypes are described in Table 4. As shown in Table 5, by employing linear regression, we found a significant association in all four *TCF7L2* SNPs with amygdalar volume in the recessive genetic model (regression model B - adjusting for sex, age and total intracranial volume [TICV]): rs7901695- CC genotype vs. CT + TT genotypes, $\beta = -0.21$, p = 0.002; rs7903146-TT vs. TC + CC, $\beta = -0.20$, p = 0.003; rs11196205- CC vs. CG + GG, $\beta = -0.247$, p = 0.0003; and rs12255372- TT vs. TG + GG, $\beta = -0.20$, p = 0.003. These results withstood our threshold for multiple testing correction (p = 0.0042).

Controlling also for T2D-related covariates (time in the diabetes registry, mean hemoglobin A1c (HbA1C) levels and use of T2D medication [yes/no]), body mass index (BMI) and ancestry (regression model B) did not change the results substantively (rs7901695 $\beta = -0.2$, p = 0.003; rs7903146 $\beta = -0.19$, p = 0.005; rs11196205 $\beta = -0.23$, p = 0.001; and rs12255372 $\beta = -0.19$, p = 0.005), although rs12255372 and rs7903146 no longer

		r^2						
SNP		rs7901695	rs7903146	rs11196205	rs12255372			
	rs7901695		0.94	0.62	0.81			
D'	rs7903146	1		0.58	0.76			
D	rs11196205	0.97	0.97		0.61			
	rs12255372	0.94	0.88	1				

Table 3. Correlation (r^2) and D' values between analyzed *TCF7L2* SNPs.

TCF7L2 SNPs [T2D risk allele]	Genotype	N	Amygdala*	Hippocampus*	Gray matter*	WMH*
	CC	35	0.466 ± 0.082	0.496 ± 0.038	507.784 ± 60.71	13.696 ± 15.73
rs7901695 [C]	СТ	84	0.515 ± 0.087	0.508 ± 0.053	521.343 ± 47.95	12.542 ± 13.45
	TT	72	0.517 ± 0.079	0.508 ± 0.047	507.962 ± 47.37	13.272 ± 13.61
	CC	77	0.515 ± 0.078	0.504 ± 0.048	508.301 ± 47.33	13.847 ± 13.88
rs7903146 [T]	СТ	75	0.514 ± 0.089	0.510 ± 0.052	520.995 ± 48.36	12.242 ± 13.34
	TT	35	0.466 ± 0.082	0.496 ± 0.038	507.784 ± 60.71	13.696 ± 15.73
	CC	49	0.469 ± 0.086	0.494 ± 0.041	504.668 ± 59.42	14.490 ± 14.78
rs11196205 [C]	CG	93	0.521 ± 0.080	0.512 ± 0.049	522.145 ± 45.96	12.709 ± 14.54
	GG	44	0.516 ± 0.084	0.506 ± 0.050	506.499 ± 48.84	13.038 ± 11.89
	GG	77	0.512 ± 0.081	0.501 ± 0.052	507.759 ± 48.14	13.679 ± 13.42
rs12255372 [T]	GT	82	0.518 ± 0.086	0.516 ± 0.047	523.465 ± 48.63	11.705 ± 13.43
	TT	32	0.466 ± 0.081	0.492 ± 0.038	503.657 ± 57.52	14.878 ± 16.09

Table 4. Volumes (mean \pm SD) of amygdala, hippocampus, gray matter and WMH according to *TCF7L2* SNPs genotypes. *Mean of total volumes in cubic centimeters \pm SD ; Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes; WMH- white matter hyperintensities.

.....

withstood the threshold for multiple testing correction (Table 5). In all four SNPs, individuals who were homozygous of the T2D risk allele had ~9.5% smaller amygdalar volume compared to the carriers of the non-risk allele (Table 6). Adjusting also to systolic and diastolic blood pressure values did not change results (regression model C, Supplementary Table 1). In addition, *TCF7L2* rs11196205 showed a significant association with amygdalar volume in the additive genetic model (regression model B- p = 0.005; model B- p = 0.008) but did not remain significant when implementing multiple testing correction (Table 5).

In order to analyze the potential distinct effect of the four *TCF7L2* SNPs (under recessive model), we performed a conditional analysis, including a second SNP as a covariate in the regression model. In the joint analysis of rs11196205 and each of the other three SNP (separately), the association with amygdalar volume was still significant (regression model A- p = 0.039, p = 0.044, p = 0.033 – controlling for rs7901695, rs7903146 or rs12255372 respectively), or approaching significance (model B- p = 0.08, p = 0.086, p = 0.067 – controlling for rs7901695, rs7903146 or rs12255372 respectively) (Table 7). However, in the joint analysis of the highly correlated SNP rs7901695/rs7903146 with rs11196205 or rs12255372, the association of rs7901695 (model A, p = 0.61 and p = 0.3, respectively) or rs7903146 (model A, p = 0.624 and p = 0.322, respectively) with amygdalar volume became non-significant. Similarly, the joint analysis of rs12255372 with any of the other SNPs (Table 7) was not significant. Therefore, we assume that none of the three SNPs rs7901695, rs7903146 and rs12255372 contributed independently to the association with amygdalar volume, beyond the effect of the most highly significant SNP rs11196205.

Due to a variable levels of linkage disequilibrium (LD) between the four *TCF7L2* SNPs, we performed haplotype analysis. Two haplotype blocks were found: 1. rs7901695 and rs7903146 (first block); 2. rs11196205 and rs12255372 (second block). Consistent with the recessive model, participants with two copies of the first block CT haplotype (rs7901695-C and rs7903146-T, N = 32) or participant with two copies of the second block CT haplotype (rs11196205-C and rs12255372-T, N = 35) had significantly smaller amygdalar volume compared to participants with other haplotype combinations in the same block (Supplementary Table 2). These results were essentially identical to the association of rs7903146 and rs12255372 alone, respectively (model A- p = 0.003; model B- p = 0.005). Combining all SNPs, carriers of two copies of the CTCT haplotype (rs7901695-C, rs7903146-T, rs11196205-C and rs12255372-T, N = 29) had significantly smaller amygdalar volume compared to participants with other haplotypes combinations (model A- p = 0.013; model B- p = 0.014). Information regarding the haplotype analysis, including haplotypes frequencies, is presented in Supplementary Tables 2 and 3.

A posteriori power estimates (based on the observed regression coefficients) for rs11196205 association with amygdalar volume in our sample (recessive model, minor allele frequency of 0.487, required p = 0.05), ranged from 94% (adjusting for only age, sex and TICV) to 89% (adjusting also for additional covariates). A posteriori power estimates for the other three SNPs (rs7901695, rs7903146 and rs12255372) association with amygdalar volume (recessive model, minor allele frequency of 0.382–0.403, required p = 0.05), were 77–83% (model A) and 68–78% (model B).

Interestingly, we also found a significant association of rs11196205 with hippocampal and gray matter volume under the recessive genetic model at a nominal significance level ($\beta = -0.141$, p = 0.035 and $\beta = -0.085$,

		Regression model A			Regression model B				
		Unstandar	rdized ts	Standardized Coefficients		Unstandardized Star Coefficients Coe		Standardized Coefficients	
TCF7L2 SNPs [T2D risk allele]	Genetic model	В	Std. Error	6	p value	В	Std. Error	6	p value
Amvgdala				1	1			1-	
18	Additive	-0.018	0.008	-0.155	0.0270	-0.016	0.0080	-0.140	0.0450
rs7901695 [C]	Recessive	-0.046	0.015	-0.208	0.0020	-0.043	0.015	-0.198	0.003
	Dominant	-0.011	0.012	-0.060	0.395	-0.008	0.012	-0.043	0.541
	Additive	-0.018	0.008	-0.157	0.025	-0.017	0.008	-0.147	0.036
rs7903146 [T]	Recessive	-0.044	0.015	-0.202	0.003	-0.042	0.015	-0.192	0.005
	Dominant	-0.013	0.012	-0.073	0.307	-0.011	0.012	-0.064	0.367
	Additive	-0.023	0.008	-0.194	0.005	-0.022	0.008	-0.181	0.008
rs11196205 [C]	Recessive	-0.048	0.013	-0.247	0.0003	-0.044	0.013	-0.226	0.001
	Dominant	-0.013	0.014	-0.066	0.343	-0.013	0.014	-0.067	0.333
	Additive	-0.016	0.008	-0.134	0.054	-0.014	0.008	-0.118	0.090
rs12255372 [T]	Recessive	-0.046	0.015	-0.203	0.003	-0.043	0.015	-0.191	0.005
	Dominant	-0.007	0.012	-0.039	0.585	-0.004	0.012	-0.021	0.762
Hippocampus									
	Additive	-0.003	0.004	-0.039	0.559	-0.002	0.004	-0.027	0.685
rs7901695 [C]	Recessive	-0.010	0.008	-0.077	0.246	-0.009	0.008	-0.071	0.275
	Dominant	0.000	0.007	0.005	0.944	0.002	0.007	0.020	0.770
	Additive	0.000	0.004	-0.002	0.978	0.000	0.004	0.004	0.950
rs7903146 [T]	Recessive	-0.009	0.008	-0.069	0.300	-0.008	0.008	-0.066	0.317
	Dominant	0.005	0.007	0.055	0.425	0.006	0.007	0.061	0.363
	Additive	-0.006	0.005	-0.085	0.205	-0.005	0.004	-0.069	0.287
rs11196205 [C]	Recessive	-0.015	0.007	-0.141	0.035	-0.013	0.007	-0.121	0.064
	Dominant	0.001	0.008	0.005	0.938	0.001	0.007	0.010	0.885
	Additive	0.001	0.005	0.015	0.822	0.002	0.004	0.032	0.628
rs12255372 [T]	Recessive	-0.015	0.008	-0.116	0.079	-0.013	0.008	-0.101	0.121
	Dominant	0.011	0.007	0.115	0.090	0.013	0.007	0.130	0.052
Gray Matter									
	Additive	-2.602	2.974	-0.037	0.383	-2.509	2.955	-0.036	0.397
rs7901695 [C]	Recessive	-7.397	5.451	-0.057	0.176	-7.223	5.366	-0.056	0.180
	Dominant	-0.937	4.497	-0.009	0.835	-0.759	4.481	-0.007	0.866
	Additive	-1.939	2.902	-0.028	0.505	-1.997	2.884	-0.029	0.490
rs7903146 [T]	Recessive	-6.473	5.419	-0.050	0.234	-6.448	5.364	-0.050	0.231
	Dominant	-0.216	4.413	-0.002	0.961	-0.333	4.388	-0.003	0.940
	Additive	-4.048	3.034	-0.056	0.184	-3.580	2.987	-0.050	0.232
rs11196205 [C]	Recessive	-9.779	4.830	-0.085	0.044	-8.626	4.767	-0.075	0.072
	Dominant	-0.667	5.099	-0.006	0.896	-0.640	5.019	-0.005	0.899
10055050 [75]	Additive	-2.323	2.987	-0.033	0.438	-2.085	2.969	-0.030	0.483
rs12255372[1]	Recessive	-9.199	5.640	-0.068	0.105	-8.193	5.564	-0.061	0.143
3473.671	Dominant	0.500	4.415	0.005	0.910	0.495	4.400	0.005	0.911
WMH	A .] .]:6:100	0.140	1 429	0.008	0.017	0.042	1 441	0.002	0.077
m7001605 [C]	Additive	-0.149	1.428	-0.008	0.917	-0.042	1.441	-0.002	0.9//
187901095 [C]	Dominant	0.558	2.035	0.016	0.855	0.759	2.035	0.021	0.775
	Additive	-0.709	2.147	-0.025	0.742	-0.011	1 130	-0.021	0.779
re7903146 [T]	Recessive	-1.822	2 139	-0.063	0.315	-1.755	2 128	-0.061	0.318
13/ 505140 [1]	Dominant	_1.022	1 699	-0.062	0.373	-1.423	1.692	-0.063	0.401
	Additive	0.220	1.099	0.014	0.415	0.238	1.092	0.015	0.401
rs11196205 [C]	Recessive	0.497	1.171	0.020	0.791	0.355	1.100	0.013	0.851
1011170205[0]	Dominant	0.072	1 949	0.003	0.970	0.279	1 940	0.011	0.886
<u> </u>	Additive	-0.667	1.150	-0.043	0.563	-0.711	1.157	-0.046	0.540
rs12255372 [T]	Recessive	-0.511	2.208	-0.017	0.817	-0.548	2.202	-0.018	0.804
10122000/2 [1]	Dominant	-1.118	1.673	-0.050	0.505	-1.184	1.682	-0.053	0.483

Table 5. Association of *TCF7L2* SNPs with amygdala, hippocampus, gray matter and white matter hyperintensities (WMH) volume. Each SNP was analyzed for association with the four phenotypes under additive, recessive and dominant genetic models, using linear regression. The basic regression model (A) included adjusting for age, sex and TICV (except of WMH). In addition to the covariates included in model A, the second model (B), was also adjusted for T2D related characteristics, BMI and ancestry. Significant associations are in bold. Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes; WMH- white matter hyperintensities.

<i>TCF7L2</i> SNPs [T2D risk allele]	Genotype group	N	Amygdala*
	CC	35	0.466 ± 0.082
rs7901695 [C]	CT + TT	156	0.516 ± 0.083
	Volume difference	e	9.7%
	CC+CT	152	0.514 ± 0.083
rs7903146 [T]	TT	35	0.466 ± 0.082
	Volume difference	9.3%	
	CC	49	0.469 ± 0.086
rs11196205 [C]	CG+GG	137	0.519 ± 0.081
	Volume difference	9.6%	
	GG+GT	159	0.515 ± 0.083
rs12255372 [T]	TT	32	0.466 ± 0.081
	Volume difference	9.5%	

Table 6. Amygdalar volumes (mean \pm SD) according to *TCF7L2* SNPs genotypes groups (recessive genetic model). *Mean of total volumes in cubic centimeters \pm SD ; Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes.

Amygdala Regression Regression model A- p Adjusted for SNP model B- p SNP value value rs11196205 0.609 0.501 rs7901695* rs12255372 0.304 0.297 0 529 rs11196205 0.624 rs7903146* rs12255372 0.322 0.321 rs7901695 0.039 0.08 rs11196205 rrs7903146 0.044 0.086 rs12255372 0.033 0.067 0.435 0.495 rs7901695 rs12255372 rrs7903146 0 4 5 2 0.505 rs11196205 0.762 0.651

Table 7. Conditional analysis for the association of *TCF7L2* SNPs with amygdalar volume under recessive model. The basic regression model (A) included adjusting for age, sex and TICV. In addition to the covariates included in model A, the second model (B), was also adjusted for T2D related characteristics, BMI and ancestry. Significant associations are in bold. *Conditional analysis for the rs7903146 and rs7901695 SNP pair was not performed due to the high correlation between them ($r^2 = 0.94$). Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes;

.....

p = 0.044, respectively), but these results became marginal when the second set of covariates was added (model B- $\beta = -0.121$, p = 0.064 and $\beta = -0.075$, p = 0.072, respectively) (Table 5). All other associations (amygdalar volume in the additive and dominant models, as well as all models for WMH, hippocampal and gray matter volumes) did not reach the required level of significance after correction for multiple testing.

Association of *TCF7L2* SNPs with left versus right amygdalar volume. In a secondary analysis we applied similar linear regression for left and right amygdala separately, in accordance to previous evidence in the literature showing differences between the two sides³⁴. As shown in Table 8, association results of all four SNPS are stronger (at the significance level achieved) for the left amygdala under recessive model (Model B- rs7901695 $\beta = -0.19$, p=0.005; rs7903146 $\beta = -0.19$, p=0.006; rs11196205 $\beta = -0.24$, p=0.0004; and rs12255372 $\beta = -0.21$, p=0.002), while weaker for the right amygdala (Model B- rs7901695 $\beta = -0.18$, p=0.012; rs7903146 $\beta = -0.17$, p=0.018; rs11196205 $\beta = -0.16$, p=0.025). Looking at the left amygdala separately, results of conditional (Table 9) and haplotype (Supplementary Table 2) analyses are similar to that of the total amygdalar volume.

Discussion

The well-established association of *TCF7L2* with T2D and the link between T2D and brain imaging changes, have motivated us study the association of this gene with neuroimaging phenotypes in our sample of elderly T2D Jewish patients. We have found a consistent association of *TCF7L2* SNPs with amygdalar volume. In the four investigated SNPs (rs7901695, rs7903146, rs11196205, and rs12255372), carriers of two copies of the T2D risk allele had smaller amygdalar volume, compared to carriers of the non-risk allele (recessive model), while controlling for sex, age and TICV. Adjusting also for T2D related covariates, BMI, ancestry and blood pressure did

	Regression			Regression model B				
	Unstandare Coefficient	dized s	Standardized Coefficients		Unstandardized Coefficients		Standardized Coefficients	
<i>TCF7L2</i> SNPs [T2D risk allele]	В	Std. Error	β	p value	В	Std. Error	β	p value
Amygdala- Left								
rs7901695 [C]	-0.053	0.0180	-0.199	0.003	-0.050	0.018	-0.188	0.005
rs7903146 [T]	-0.052	0.018	-0.196	0.004	-0.049	0.018	-0.185	0.006
rs11196205 [C]	-0.061	0.016	-0.257	0.0001	-0.056	0.016	-0.236	0.0004
rs12255372 [T]	-0.062	0.018	-0.224	0.001	-0.059	0.018	-0.212	0.002
Amygdala- Right				·				
rs7901695 [C]	-0.047	0.0180	-0.183	0.0090	-0.045	0.0180	-0.176	0.0120
rs7903146 [T]	-0.044	0.018	-0.175	0.013	-0.042	0.018	-0.168	0.018
rs11196205 [C]	-0.047	0.016	-0.211	0.003	-0.043	0.016	-0.192	0.006
rs12255372 [T]	-0.044	0.018	-0.167	0.017	-0.042	0.018	-0.158	0.025

Table 8. Association of TCF7L2 SNPs with left and right amygdalar volume (recessive genetic model). The basic linear regression model (A) included adjusting for age, sex and TICV. In addition to the covariates included in model A, the second model (B), was also adjusted for T2D related characteristics, BMI and ancestry. Significant associations are in bold. Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes.

Left amygdala								
SNP	Adjusted for SNP	Regression model A- p value	Regression model B- p value					
rc7001605*	rs11196205	0.943	0.828					
137 901095	rs12255372	0.754	0.752					
07002146	rs11196205	0.952	0.86					
187903146**	rs12255372	0.775	0.784					
rs11196205	rs7901695	0.012	0.027					
	rrs7903146	0.013	0.027					
	rs12255372	0.038	0.076					
	rs7901695	0.119	0.139					
rs12255372	rrs7903146	0.121	0.14					
	rs11196205	0.588	0.488					

Table 9. Conditional analysis for the association of *TCF7L2* SNPs with left amygdalar volume under recessive model. The basic regression model (A) included adjusting for age, sex and TICV. In addition to the covariates included in model A, the second model (B), was also adjusted for T2D related characteristics, BMI and ancestry. Significant associations are in bold. *Conditional analysis for the rs7903146 and rs7901695 SNP pair was not performed due to the high correlation between them ($r^2 = 0.94$). Abbreviations: SNP- single nucleotide polymorphism; T2D- type 2 diabetes.

.....

not change the results substantially. Further examination of the left and right amygdala separately, revealed that the association is derived mainly due to the left amygdalar volume (p = 0.0004-0.006) than the right amygdalar volume (p = 0.006-0.025). On conditional analysis, we found that rs7901695, rs12255372 or rs7903146 SNPs associations with amygdalar volume were not independent of the most highly significant SNP rs11196205, and therefore only one association signal was detected in region. No associations of hippocampal, gray matter and WMH volumes with *TCF7L2 SNPs* withstood Bonferroni adjustment for multiple testing correction.

Several limitations of this study should be considered. Our sample size (N = 191 individuals) is considered small in the context of a genetic association study. Nevertheless, the sample is unique, since it includes only T2D elderly, a population at risk for cognitive decline and dementia. All participants had clinical (including measures of glycemic control), neuroimaging and genetic data. Some of the associations survived Bonferroni correction for multiple testing, indicating robustness of the results and therefore reducing the likelihood of false positive results. The mere nominal level of association with gray matter and hippocampal volume, which did not withstand the Bonferroni correction, might be due to a small sample size, and larger size would have been an advantage in terms of statistical power. In addition, the cross-sectional design of the study impedes reaching conclusions of causality. The longitudinal component of the IDCD is ongoing and may assist in shedding light on functional effect of this association in the future.

Previous neuroimaging genetics studies did not find association of *TCF7L2* with amygdalar volume. Of particular interest is a recent GWAS meta-analysis study of ~30,000 participants (mostly European origin) conducted by the Enhancing Neuro Imaging Genetics through Meta-Analysis (ENIGMA) consortium³⁵ (which did not include specific cohorts of T2D patients in particular). No significant associations (additive model, p < 0.05) were found in this study between amygdalar volume and *TCF7L2* SNPs rs7901695, rs7903146 and rs12255372. Although rs11196205 was not tested directly in the ENIGMA GWAS, its proxy SNP rs10885409 (D' = 1, $r^2 > 0.95$), found by using the Broad institute SNP proxy search (http://archive.broadinstitute.org/mpg/snap/ldsearch.php), was not significantly associated with amygdalar volume as well. Several explanations for the discrepancy in results are plausible in addition to different genetic model, including that the observed association of *TCF7L2* with amygdalar volume is specific to the Jewish population, or alternatively is specific to T2D affected individuals. Our study did not include control participants without T2D, and therefore we cannot address the generalization of this finding to non-T2D individuals. It is also possible that the association is influenced by older age and potentially not found in younger population (this sample include individuals aged 65 years and older). Taken together, at the current stage, our findings should be considered as preliminary and caution is required in their interpretation. Further studies are required in various populations to validate it.

The amygdala, one of the limbic system's components, has been implicated in several functions - mainly emotional processing and responses (e.g. fear, anxiety, and aggression), decision-making, associative learning and memory³⁶. Amygdalar aberrant function or structure is common in neurodevelopmental disorders³⁷. Previous studies in various populations have reported association of variation in several genes with amygdalar volume, including *STMN1* and *SLC6A4*³⁸, *CACNA1C*^{39,40} and the oxytocin receptor *OXTR*^{41,42}.

Consistent with our results, previous reports have demonstrated an association between structural brain changes and T2D, e.g. lower brain volumes and greater brain atrophy in T2D patients²⁴⁻²⁶, including amygdala³⁰. Indeed, greater amygdalar atrophy had been associated with high plasma glucose levels within the normal range³¹.

TCF7L2 is expressed in many brain regions, including the amygdala in mice⁴³ and at a relatively low level in human amygdala (Genotype Tissue expression portal, GTex, Broad Institute; https://gtexportal.org/home/ gene/TCF7L2/). As part of the Wnt/beta-catenin signaling pathway, *TCF7L2* plays role in the activation of lymphoid enhancer-binding factor 1/T cell factor (LEF1/TCF) transcription factors complexes. The Wnt/ beta-catenin signaling is involved in neuroplasticity, adult neurogenesis and CNS development⁴⁴⁻⁴⁶, as well as in amygdala-dependent learning and long-term memory formation⁴⁷. Decreased levels of beta-catenin were found in the amygdala of rats that showed behavioral sensitization to administration of drugs of abuse⁴⁸. In humans, polymorphisms in *TCF7L2* were associated with schizophrenia and bipolar disorder²⁰⁻²². At the behavioral level in animal models, *TCF7L2* deficient mice demonstrated altered anxiety like behavior and fear learning⁴⁹, and this gene mediated cellular and behavioral response to lithium treatment in mice and zebrafish⁵⁰. Combined, these evidences implicate a role of *TCF7L2* in brain function and behavioral phenotypes.

To conclude, our results in a sample of T2D elderly demonstrate for the first-time associations of four *TCF7L2 SNPs* with amygdalar volume. Confirmation of these results in additional cohorts is required in order to reach more definitive conclusions.

Methods

Sample. Participants were recruited from the Israel Diabetes and Cognitive Decline (IDCD) study, a collaboration of the Icahn School of Medicine, Mount Sinai, NY, USA, Sheba Medical Center, Israel, and the Maccabi Health Services (MHS), Israel. The IDCD study design has been previously described in detail⁵¹. Briefly, community-dwelling Israeli elderly individuals with T2D (265 years old) were recruited from the MHS diabetes registry. Criteria for enrolment into the IDCD study were: (1) having T2D (defined as any of the following- (A) HbA1c > 7.25%; (B) Glucose blood levels of 200 mg/dl on two examinations more than 3 months apart; (C) purchase of diabetic medication twice within 3 months; or (D) diagnosis of T2D (International Classification of Diseases [ICD9] code) by a general practitioner, internist, endocrinologist, ophthalmologist, or diabetes advisor, supported by a HbA1c > 6.5% or glucose > 125 mg/dl within half a year); (2) normal cognition at entry to the IDCD study; (3) being free of any neurological (e.g., Parkinson's disease, stroke), psychiatric (e.g. schizophrenia) or other diseases (e.g., alcohol or drug abuse) that might affect cognition; (4) having an informant; (5) fluency in Hebrew; (6) living in the area of Tel Aviv. The Diabetes Registry has collected detailed laboratory, medication, and diagnoses information since 1998⁵². Based on self-report, the IDCD individuals are unrelated to each other (at least at first- and second-degree level). The HbA1c and blood pressure (systolic and diastolic) values were calculated for each participant as the means of all measurements in the diabetes registry.

MRI acquisition. A randomly recruited sub-sample of the IDCD cohort underwent MRI scan, performed at the Diagnostic Imaging Department, Sheba Medical Center using a 3 Tesla scanner (GE, Signa HDxt, v16VO2). High-resolution (1 mm³) images were acquired by using a 3D inversion recovery prepared fast spoiled gradient-echo (FSPGR) T1-weighted sequence (TR/TE = 7.3/2.7 s, 20° flip angle, TI 450 ms). In addition, a T2-weighted fluid-attenuated inversion recovery (FLAIR) sequence was acquired with the following parameters: Repetition time/Echo time (TR/TE) 9500/123 ms, axial slices, slice-width/gap 3/0.4 mm, 22 cm FOV, 64×64 matrix, 90° flip-angle.

MRI analysis. For volumetric analysis, the voxel based morphometry (VBM⁵³) toolbox, (http://www.fil.ion. ucl.ac.uk/spm/ext/#VBMtools) implemented in Statistical Parametric Mapping (SPM8) software was used on the T1 weighted anatomical images. This procedure included automated iterative skull stripping, segmentation of the images into gray matter, white matter (WM), cerebrospinal fluid probability images, and spatial normalization of the gray matter images to a customized gray matter template in standard MNI (Montreal Neurological Institute) space. Finally, the gray matter maps were smoothed using an 8 mm Gaussian kernel. Gray matter probability maps were thresholded at 0.2 to minimize inclusion of incorrect tissue types. Total intracranial volume (TICV) was calculated by summing the segmented and thresholded images (TICV = gray matter + white matter + cerebrospinal fluid). Based on our a-priori hypothesis, we used a region of interest (ROI) approach centered on the amygdala and hippocampus, identified with the 'Human Automated Anatomical Labelling (AAL) atlas²⁵⁴ within the Wake

Forest University PickAtlas (http://www.rad.wfubmc.edu/fmri) and extracted using the MarsBaR ROI toolbox⁵⁵ as implemented in SPM12. All reported volumes are total regional volumes.

For WMH quantification we used the Lesion segmentation toolbox (LST) (implemented in SPM8), following previously described methods³². The default LST settings were used with the exception of κ (k), a value indicating the threshold for the initial lesion mask. Visual inspection of the probability maps across participants by using various k values, to maximize sensitivity while reducing false positive results, indicated that a k = 0.15 was the optimal value for our sample images. This procedure generated one binary lesion image per participant from which a total lesion volume (in milliliters) map was extracted.

SNPs selection and genotyping. Four intronic *TCF7L2* SNPs (rs7901695, rs7903146, rs11196205, and rs12255372) were selected for this study (Table 2), based on ample evidence of their confirmed association with T2D and related traits^{4,5}. These SNPs were genotyped with the Sequenom MassARRAY system, at the Washington University Human Genetics Division Genotyping Core, St. Louis, USA. Quality control measures were implemented.

Statistical analysis. We employed hierarchical linear regression to study the association of the *TCF7L2* SNPs with amygdalar, hippocampal, gray matter and WMH volumes, under three genetic models (additive, dominant and recessive – referring to the effect of the T2D risk allele). In the basic regression model (model A), we controlled for sex, age at IDCD baseline recruitment and TICV (this covariate was included in the models for gray matter, hippocampus and amygdala). In the second step (model B), we included in the regression model all the covariates from model A, in addition to a set of T2D related characteristics (time in the MHS diabetes registry [an approximation to T2D duration⁵⁶], mean HbA1C levels, use of T2D medication [yes/n0]), mean body mass index (BMI), and ancestry (Ashkenazi vs. Non-Ashkenazi, based on self-report and land of birth data). In the third step (model C), we included in the regression model all the covariates from models A and B, in addition to mean systolic and mean diastolic blood pressure.

The analysis was conducted for each SNP and brain region separately. For each brain region, a two-sided p value of 0.0042 (0.05/12) was considered statistically significant following employment of Bonferroni correction for multiple testing (0.05/[4 SNPs included in the final analysis \times 3 genetic models]). For statistical analysis, we used SPSS version 21.0 (SPSS Inc., Chicago, IL, USA). Hardy–Weinberg calculations, SNPs pairwise correlation and linkage disequilibrium (LD) values were obtained with PLINK (http://pngu.mgh.harvard.edu/purcell/plink)⁵⁷. For the WMH, we applied square-root transformation to obtain normal distribution. Power calculation for *TCF7L2* SNPs association with amygdalar volume was carried by Quanto v1.2.4 software⁵⁸. To assess a potential distinct contribution of the four *TCF7L2* SNPs on amygdalar volume, we performed conditional analysis for each SNP (adjusting for a second SNPs within the regression model, coded recessively).

For haplotype analysis, we used PLINK to determine haplotypes blocks and frequencies. We employed hierarchical linear regression to study the association of the *TCF7L2* haplotypes with amygdalar volume under recessive model (comparing carriers of two copies of haplotype of interest, to carriers of all other haplotypes combinations), adjusting for covariates.

Study approval and informed consent. All participants provided informed consent, and all experimental protocols were approved by the institutional review boards (IRBs) of all three collaborating institutions (Icahn School of Medicine, Mount Sinai, NY, USA, Sheba Medical Center, Israel, and MHS, Israel). In addition, all the methods were carried out in accordance with the relevant guidelines and regulations.

Data availability

The datasets generated and analyzed during the current study are available from the corresponding author on reasonable request.

Received: 9 August 2018; Accepted: 8 August 2019;

Published online: 01 November 2019

References

- 1. Ahlqvist, E., Ahluwalia, T. S. & Groop, L. Genetics of type 2 diabetes. Clin Chem 57, 241-254 (2011).
- Billings, L. K. & Florez, J. C. The genetics of type 2 diabetes: what have we learned from GWAS? Ann N Y Acad Sci 1212, 59–77 (2010).
- 3. Sparso, T. *et al.* Combined analysis of 19 common validated type 2 diabetes susceptibility gene variants shows moderate discriminative value and no evidence of gene-gene interaction. *Diabetologia* **52**, 1308–1314 (2009).
- Peng, S. et al. TCF7L2 gene polymorphisms and type 2 diabetes risk: a comprehensive and updated meta-analysis involving 121,174 subjects. Mutagenesis 28, 25–37 (2013).
- 5. Tong, Y. *et al.* Association between TCF7L2 gene polymorphisms and susceptibility to type 2 diabetes mellitus: a large Human Genome Epidemiology (HuGE) review and meta-analysis. *BMC Med Genet* **10**, 15 (2009).
- 6. Palmer, N. D. et al. Resequencing and analysis of variation in the TCF7L2 gene in African Americans suggests that SNP rs7903146 is the causal diabetes susceptibility variant. *Diabetes* **60**, 662–668 (2011).
- 7. Ferreira, M. C., da Silva, M. E. R., Fukui, R. T., Arruda-Marques, M. D. C. & Dos Santos, R. F. TCF7L2 correlation in both insulin secretion and postprandial insulin sensitivity. *Diabetol Metab Syndr* **10**, 37 (2018).
- 8. Gjesing, A. P. *et al.* Carriers of the TCF7L2 rs7903146 TT genotype have elevated levels of plasma glucose, serum proinsulin and plasma gastric inhibitory polypeptide (GIP) during a meal test. *Diabetologia* **54**, 103–110 (2011).
- 9. Ip, W., Chiang, Y. T. & Jin, T. The involvement of the wnt signaling pathway and TCF7L2 in diabetes mellitus: The current understanding, dispute, and perspective. *Cell Biosci* 2, 28 (2012).
- 10. MacDonald, B. T., Tamai, K. & He, X. Wnt/beta-catenin signaling: components, mechanisms, and diseases. Dev Cell 17, 9-26 (2009).
- 11. Migliorini, A. & Lickert, H. Beyond association: A functional role for Tcf7l2 in beta-cell development. Mol Metab 4, 365–366 (2015).
- 12. Lyssenko, V. *et al.* Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *J Clin Invest* **117**, 2155–2163 (2007).

- 13. Sakhneny, L. et al. Pancreatic Pericytes Support Beta-Cell Function in a Tcf7l2-Dependent Manner. Diabetes (2017).
- 14. Chen, X. et al. The Diabetes Gene and Wnt Pathway Effector TCF7L2 Regulates Adipocyte Development and Function. Diabetes (2018).
- Lee, S., Lee, C. E., Elias, C. F. & Elmquist, J. K. Expression of the diabetes-associated gene TCF7L2 in adult mouse brain. J Comp Neurol 517, 925–939 (2009).
- Nazwar, T. A., Glassmann, A. & Schilling, K. Expression and molecular diversity of Tcf7l2 in the developing murine cerebellum and brain. J Neurosci Res 87, 1532–1546 (2009).
- 17. Nagalski, A. *et al.* Postnatal isoform switch and protein localization of LEF1 and TCF7L2 transcription factors in cortical, thalamic, and mesencephalic regions of the adult mouse brain. *Brain Struct Funct* (2012).
- Murray, K. D., Choudary, P. V. & Jones, E. G. Nucleus- and cell-specific gene expression in monkey thalamus. *Proc Natl Acad Sci USA* 104, 1989–1994 (2007).
- Prokunina-Olsson, L. & Hall, J. L. Evidence for neuroendocrine function of a unique splicing form of TCF7L2 in human brain, islets and gut. *Diabetologia* 53, 712–716 (2010).
- 20. Hansen, T. et al. At-risk variant in TCF7L2 for type II diabetes increases risk of schizophrenia. Biol Psychiatry 70, 59–63 (2011).
- 21. Alkelai, A. *et al.* Association of the type 2 diabetes mellitus susceptibility gene, TCF7L2, with schizophrenia in an Arab-Israeli family sample. *PLoS One* 7, e29228 (2012).
- Winham, S. J. et al. Genome-wide association study of bipolar disorder accounting for effect of body mass index identifies a new risk allele in TCF7L2. Mol Psychiatry 19, 1010–1016 (2014).
- 23. Ganmore, I. & Beeri, M. S. Magnitude and Trajectories of Cognitive Dysfunction in Type 2 Diabetes Mellitus. In: Srikanth V and Arvanitakis Z (eds) Type 2 Diabetes and Dementia. Elsevier pp 29-47 (2018).
- 24. McCrimmon, R. J., Ryan, C. M. & Frier, B. M. Diabetes and cognitive dysfunction. Lancet 379, 2291-2299 (2012).
- Li, W., Risacher, S. L., Huang, E., Saykin, A. J. & Alzheimer's Disease Neuroimaging, I. Type 2 diabetes mellitus is associated with brain atrophy and hypometabolism in the ADNI cohort. *Neurology* 87, 595–600 (2016).
- Biessels, G. J. & Reijmer, Y. D. Brain changes underlying cognitive dysfunction in diabetes: what can we learn from MRI? *Diabetes* 63, 2244–2252 (2014).
- Moulton, C. D., Costafreda, S. G., Horton, P., Ismail, K. & Fu, C. H. Meta-analyses of structural regional cerebral effects in type 1 and type 2 diabetes. *Brain Imaging Behav* 9, 651–662 (2015).
- Chen, Z., Li, L., Sun, J. & Ma, L. Mapping the brain in type II diabetes: Voxel-based morphometry using DARTEL. Eur J Radiol 81, 1870–1876 (2012).
- 29. Moran, C. *et al.* Brain atrophy in type 2 diabetes: regional distribution and influence on cognition. *Diabetes Care* **36**, 4036–4042 (2013).
- 30. den Heijer, T. *et al.* Type 2 diabetes and atrophy of medial temporal lobe structures on brain MRI. *Diabetologia* **46**, 1604–1610 (2003).
- Cherbuin, N., Sachdev, P. & Anstey, K. J. Higher normal fasting plasma glucose is associated with hippocampal atrophy: The PATH Study. Neurology 79, 1019–1026 (2012).
- Livny, A. et al. Long-term Variability in Glycemic Control Is Associated With White Matter Hyperintensities in APOE4 Genotype Carriers With Type 2 Diabetes. Diabetes Care 39, 1056–1059 (2016).
- Livny, A. et al. Haptoglobin 1-1 Genotype Modulates the Association of Glycemic Control With Hippocampal Volume in Elderly Individuals With Type 2 Diabetes. Diabetes 66, 2927–2932 (2017).
- Markowitsch, H. J. Differential contribution of right and left amygdala to affective information processing. Behav Neurol 11, 233–244 (1998).
- 35. Hibar, D. P. et al. Common genetic variants influence human subcortical brain structures. Nature 520, 224-229 (2015).
- 36. Janak, P. H. & Tye, K. M. From circuits to behaviour in the amygdala. Nature 517, 284-292 (2015).
- Schumann, C. M., Bauman, M. D. & Amaral, D. G. Abnormal structure or function of the amygdala is a common component of neurodevelopmental disorders. *Neuropsychologia* 49, 745–759 (2011).
- Stjepanovic, D., Lorenzetti, V., Yucel, M., Hawi, Z. & Bellgrove, M. A. Human amygdala volume is predicted by common DNA variation in the stathmin and serotonin transporter genes. *Transl Psychiatry* 3, e283 (2013).
- Lancaster, T. M., Foley, S., Tansey, K. E., Linden, D. E. & Caseras, X. CACNA1C risk variant is associated with increased amygdala volume. Eur Arch Psychiatry Clin Neurosci 266, 269–275 (2016).
- Wolf, C. et al. CACNA1C genotype explains interindividual differences in amygdala volume among patients with schizophrenia. Eur Arch Psychiatry Clin Neurosci 264, 93–102 (2014).
- Furman, D. J., Chen, M. C. & Gotlib, I. H. Variant in oxytocin receptor gene is associated with amygdala volume. Psychoneuroendocrinology 36, 891–897 (2011).
- 42. Wang, J. et al. Neural mechanisms of oxytocin receptor gene mediating anxiety-related temperament. Brain Struct Funct 219, 1543–1554 (2014).
- Weaver, C., Turner, N. & Hall, J. Review of the neuroanatomic landscape implicated in glucose sensing and regulation of nutrient signaling: immunophenotypic localization of diabetes gene Tcf7l2 in the developing murine brain. J Chem Neuroanat 45, 1–17 (2012).
- 44. Backman, M. *et al.* Effects of canonical Wnt signaling on dorso-ventral specification of the mouse telencephalon. *Dev Biol* **279**, 155–168 (2005).
- 45. Budnik, V. & Salinas, P. C. Wnt signaling during synaptic development and plasticity. Curr Opin Neurobiol 21, 151–159 (2011).
- Brinkmeier, M. L., Potok, M. A., Davis, S. W. & Camper, S. A. TCF4 deficiency expands ventral diencephalon signaling and increases induction of pituitary progenitors. *Dev Biol* 311, 396–407 (2007).
- 47. Maguschak, K. A. & Ressler, K. J. Wnt signaling in amygdala-dependent learning and memory. J Neurosci 31, 13057-13067 (2011).
- Cuesta, S., Severin, M. J., Batuecas, J., Rosso, S. B. & Pacchioni, A. M. Wnt/beta-catenin pathway in the prefrontal cortex is required for cocaine-induced neuroadaptations. *Addict Biol* 22, 933–945 (2017).
- 49. Savic, D. et al. Modulation of Tcf7l2 expression alters behavior in mice. PLoS One 6, e26897 (2011).
- Misztal, K. et al. TCF7L2 mediates the cellular and behavioral response to chronic lithium treatment in animal models. Neuropharmacology 113, 490–501 (2017).
- 51. Beeri, M. S. *et al.* The Israel Diabetes and Cognitive Decline (IDCD) study: Design and baseline characteristics. *Alzheimers Dement* **10**, 769–778 (2014).
- Heymann, A. D. et al. The implementation of managed care for diabetes using medical informatics in a large Preferred Provider Organization. Diabetes Res Clin Pract 71, 290–298 (2006).
- 53. Ashburner, J. & Friston, K. J. Voxel-based morphometry-the methods. Neuroimage 11, 805-821 (2000).
- Tzourio-Mazoyer, N. *et al.* Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* 15, 273–289 (2002).
- 55. Brett, M., Anton, J. L., Valabregue, R. & Poline, J. B. Region of interest analysis using an SPM toolbox [abstract]. Presented at the 8th International Conference on Functional Mapping of the Human Brain. Sendai, Japan Available on CD-ROM in NeuroImage, Vol 16, No 2, abstract 497 (2002).
- West, R. K. et al. The association of duration of type 2 diabetes with cognitive performance is modulated by long-term glycemic control. Am J Geriatr Psychiatry 22, 1055–1059 (2014).

- 57. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559–575 (2007).
- 58. Gauderman, W. J. Sample size requirements for association studies of gene-gene interaction. Am J Epidemiol 155, 478-484 (2002).

Acknowledgements

This research was supported by NIA grants R01 AG034087 and R21 AG043878 for Dr. Beeri and P50 AG05138 for Dr. Mary Sano; the Bader Philanthropies and the Leroy Schecter Foundation, as well as a New Investigator Award in Alzheimer's Disease from the American Federation for Aging Research for Dr. Cooper.

Author contributions

I.G. and L.G. researched data, performed statistical analysis and wrote the manuscript; A.L. performed MRI data acquisition, and participated in analysis; R.R.S. and M.S.B. contributed to research design and reviewed the manuscript; I.C., A.A., S.S., G.T. and A.H. reviewed the manuscript and contributed to discussion.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is avilable for this paper at https://doi.org/10.1038/s41598-019-48899-3.

Correspondence and requests for materials should be addressed to I.G.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2019