OPEN

Author Correction: Structural Features of a Bacteroidetes-Affiliated Cellulase Linked with a Polysaccharide Utilization Locus

A. E. Naas, A. K. MacKenzie, B. Dalhus, V. G. H. Eijsink & P. B. Pope

Correction to: *Scientific Reports* <https://doi.org/10.1038/srep11666>, published online 02 July 2015

This Article contains an error, where the protein name corresponding to 4IM4 (pdbid) is incorrectly given as “Lic26A-Cel5E”, and should read “CelE”.

In addition, three references were omitted from Table 2, and are given below as References 1–3.

The correct Table 2 and Figure 4 are given below as Table 1 and Figure 1 respectively.

Finally, the updated Supplementary Information file, containing the correct protein name in Figure S1, is given below.

Published online: 08 April 2020

	Structural comparison			Enzymatic activities							Reference
	DALI Z-score	RMSD (Å)	% id	CMC (U/mg)	β -glucan (U/mg)	Filter paper (U/mg)	Avicel (U/mg)	Lichenan (U/mg)	Xylan (U/mg)	Xyloglucan (U/mg)	
<i>Bacteroidetes AC2a</i> Cellulase Cel5A	—	—	—	216.8	1471.4	0.152 ^a	0.115 ^a	839.9	nd	trace	<i>This study and</i> ⁷
<i>Paenibacillus pabuli</i> Xyloglucanase XG5 (2JEP)	49.2	1.6	34	nd	nd	—	nd	nd	nd	8700	³³
<i>Clostridium cellulovorans</i> Cellulase EngD (3NDY)	47.0	1.7	33	15	42	—	0.017	—	0.5	36	¹⁵
<i>Bacteroides ovatus</i> Xyloglucanase BoGH5A (3ZMR)	46.6	1.9	32	nd	nd	—	—	nd	—	514.2	⁴⁴
<i>Clostridium thermocellum</i> Cellulase CelE (4IM4)	46.0	1.6	32	Active	75.4	Active	—	—	Active	—	^{1–3} , b
<i>Clostridium cellulolyticum</i> Cellulase CelCCA (1EDG)	45.9	1.9	32	101.3	104.3	—	0.028 (0.124) ^d	79.2	10.1	—	^{45,46} , c
<i>Piromyces rhizinflata</i> Cellulase CelAcd (3AYR)	43.2	1.8	30	344.9	576	0.64	1.39	542.5	106.2	—	⁴⁷

Table 1. Structural comparison of AC2aCel5A with its six closest structural homologues identified using the DALI server¹⁷, along with reported enzyme activities. RMSD; root-mean-square deviation of C-alpha atoms. The structural homologues are sorted based on the Z-score obtained in the DALI search. One Unit of enzyme activity was defined as the amount of enzyme releasing 1 μ mol of reducing sugar equivalents per minute. “nd” means not detected, whereas a hyphen, “—”, indicates “not tested”. ^a μ mol reducing sugar equivalents calculated as μ mol cellotriose + μ mol cellobiose + μ mol glucose, quantified by HPAEC-PAD⁷. ^bGH5 domain of CelE. ^cU/mg calculated from U/ μ mol reported in⁴⁵. ^dFor this enzyme data for both the full length protein (in parentheses), and the catalytic domain only were published.

Figure 1.

References

1. Hall, J., Hazlewood, G. P., Barker, P. J. & Gilbert, H. J. Conserved reiterated domains in *Clostridium thermocellum* endoglucanases. *Gene* **69**, 29–38 (1988).
2. Durrant, A. J., Hall, J., Hazlewood, G. P. & Gilbert, H. J. The non-catalytic C-terminal region of endoglucanase E from *Clostridium thermocellum* contains a cellulose-binding domain. *Biochem. J.* **273**, 289–293 (1991).
3. Takasuka, T. E. *et al.* Cell-free translation of biofuel enzymes. *Methods Mol. Biol.* **1118**, 71–95 (2014).

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-62786-2>.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020