OPEN

Author Correction: Salidroside promotes peripheral nerve regeneration based on tissue engineering strategy using Schwann cells and PLGA: *in vitro* and *in vivo*

Hui Liu, Peizhen Lv, Yongjia Zhu, Huayu Wu, Kun Zhang, Fuben Xu, Li Zheng & Jinmin Zhao

Correction to: *Scientific Reports*, <https://doi.org/10.1038/srep39869>, published online 05 January 2017

This Article contains errors.

In Figure 1E the image for PLGA+S-0.2 at 4d is incorrect and is duplicating an image for PLGA+S-0.1 at 6d, which was already correct at the time of publication. The correct Figure 1 appears below.

In Figure 2A the images for PLGA at 2d and TCP at 4d are incorrect. The correct Figure 2 appears below.

Published online: 09 September 2020

Figure 1. (A) Chemical structure of salidroside; (B) Preliminary drug screening analysis of RSC 96 treated on PLGA scaffold with different concentrations of salidroside after 3 days ($n = 3$, mean $\pm$ SEM); (C) Proliferative effects of salidroside on RSC96 on PLGA scaffold measured by MTT assay ($n = 3$, mean $\pm$ SEM). Different letters denote significances with $P < 0.05$ and the same letter shows no significant differences ($P \geq 0.05$); (D) Quantitative data of the mean number of SCs. Data of each bar are shown as the mean of three independent experiments $\pm$ SD. Different letters denote significances with $P < 0.05$ and the same letter shows no significant differences ($P \geq 0.05$); (E) Cell viability was measured by FDA/PI staining under microscope. (PLGA means cultured with 0 mM SDS, PLGA+S-0.1 means cultured with 0.1 mM SDS, PLGA+S-0.2 means cultured with 0.2 mM SDS, PLGA+S-0.4 means cultured with 0.4 mM SDS, TCP means cultured on TCP alone, TCP+S-0.2 means cultured with 0.2 mM SDS on TCP).

Figure 2. (A) Hematoxylin-eosin staining images showing the morphology of RSC 96 cultured in vitro in six groups at 2, 4 and 6 days. (PLGA means cultured with 0 mM SDS, PLGA+S-0.1 means cultured with 0.1 mM SDS, PLGA+S-0.2 means cultured with 0.2 mM SDS, PLGA+S-0.4 means cultured with 0.4 mM SDS, TCP means cultured on TCP alone, TCP+S-0.2 means cultured with 0.2 mM SDS on TCP). (B) Quantitative data of the mean number of SCs. Data of each bar are shown as the mean of three independent experiments $\pm$ SD. Different letters denote significances with $P < 0.05$ and the same letter shows no significant differences ($P \geq 0.05$).

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020