



OPEN

Modelling the effects of CO₂ on C₃ and C₄ grass competition during the mid-Pleistocene transition in South Africa

Michaela Ecker^{1,2}, Douglas Kelley³ & Hiromitsu Sato⁴

Palaeoenvironmental reconstructions of the interior of South Africa show a wetter environment than today and a non-analogous vegetation structure in the Early Pleistocene. This includes the presence of grasses following both C₃ and C₄ photosynthetic pathways, whereas C₃ grasses decline after the mid-Pleistocene transition (MPT, c. 1.2–0.8 Ma). However, the local terrestrial proxy record cannot distinguish between the potential drivers of these vegetation changes. In this study we show that low glacial CO₂ levels, similar to those at the MPT, lead to the local decline of C₃ grasses under conditions of decreased water availability, using a vegetation model (LPX) driven by Atmosphere–Ocean coupled General Climate Model climate reconstructions. We modelled vegetation for glacial climates under different levels of CO₂ and fire regimes and find evidence that a combination of low CO₂ and changed seasonality is driving the changes in grass cover, whereas fire has little influence on the ratio of C₃:C₄ grasses. Our results suggest the prevalence of a less vegetated landscape with limited, seasonal water availability, which could potentially explain the much sparser mid-Pleistocene archaeological record in the southern Kalahari.

South Africa can be divided into three seasonal rainfall zones (winter, summer and year-round rainfall), which in turn strongly influence the distribution of vegetation¹ (Supplementary Fig. S3). Palaeoenvironmental studies hypothesized that areas that are in the summer rainfall zone of the savanna biome might have been under the influence of an expanding winter rainfall zone during Pleistocene glacial conditions^{2–4}. This could have led to shifts in the vegetation structure of the savanna biome, which today consists broadly of tropical and subtropical grasses following the C₄ photosynthetic pathway with scattered small trees and bushes following the C₃ photosynthetic pathway¹. Rainfall seasonality, in turn, is a major control of fire⁵, and regular wildfire is common in African savannas, enabling turnover in the grass layer. C₄ grasses, on the whole, tend to recover faster after fires than C₃ grasses. This may be due to rapid regrowth due to generally higher photosynthetic rate, especially with the removal of shading from higher canopies post-fire⁶. The importance of fire in maintaining modern ecosystems has been shown in modelling studies⁷. Fire intensity and regularity is challenging to reconstruct for the Pleistocene, particularly when microcharcoal records are absent. However, there can be considerable overlap between C₃ and C₄ species of grass⁸. Atmospheric CO₂ levels are another major control of plant photosynthesis. C₃ plants, which can also include temperate grasses, thrive under high CO₂ levels, whereas C₄ grasses are more efficient under low CO₂ levels (Fig. 1). In laboratory studies, for example, glacial CO₂ has been found to increase the stomatal conductance of C₄ grasses to greater effect than C₃ grasses, while maintaining its high water-use efficiency⁹. This difference in water-use efficiency is a likely mechanism for C₄ grasses to outcompete C₃ grasses in arid regions. The vegetation response to changing CO₂ is therefore heavily dependent on local subsurface conditions¹⁰.

Approximately 900,000 years ago, ice ages switched from occurring every 41 kyr to every 100 kyr, called the mid-Pleistocene transition (MPT), which in turn had consequences for the global CO₂ record. Boron isotope reconstructions of the global CO₂ records during the MPT (c. 1.2–0.8 Ma) show glacial (c. 160–200 ppm) CO₂ levels to be particularly low; lower than in the preceding Early Pleistocene (c. 185–250 ppm). At the same time

¹Institute of Pre- and Protohistoric Archaeology, University of Kiel, Kiel, Germany. ²Archaeology Centre, University of Toronto, Toronto, Canada. ³UK Centre for Ecology and Hydrology, Wallingford, UK. ⁴Department of Earth Sciences, University of Toronto, Toronto, Canada. ✉email: mecker@ufg.uni-kiel.de




Figure 1. Prediction of atmospheric CO₂ and daytime growing season temperature conditions that favour C₃ or C₄ grasses. Black box covers CO₂ levels and potential temperatures during glacial in the central interior of South Africa before the Mid-Pleistocene transition ('40 k world') and during the Mid-Pleistocene transition ('100 k world') (adapted after¹¹ and¹²).

interglacial CO₂ levels are similar before and after the MPT (c. 250–320 ppm), but on average lower during the MPT (c. 200–300 ppm)¹³. CO₂ reconstructions from the oldest ice in Antarctica confirm lower glacial CO₂ concentrations during the MPT in comparison to the 40 kyr world before and the 100 kyr world afterwards¹⁴. In total, the glacial-interglacial span of CO₂ increased substantially. In accordance with their physiology, going from, on average, glacial 185–250 ppm to 160–200 ppm has a much larger effect on photosynthesis in C₄ plants than on C₃ plants (Fig. 1).

Recent studies suggested early human evolution during the Pliocene and Pleistocene in non-analogous, productive C₃ plant dominated environments in parts of Africa^{15,16}. In the central interior of South Africa, Wonderwerk Cave is the site with the longest and most fine-grained record of palaeoenvironmental change, covering the last two million years¹⁷. This site and other local multi-proxy records showed an environment during the Early Pleistocene (c. 1.96–0.78 Ma) that included the presence of persistent standing bodies of water and grasses following both C₃ and C₄ photosynthetic pathways^{18,19,20} (Supplementary Table S1). After c. 800 kyr, C₃ grasses show a decline in abundance at Wonderwerk Cave and are not a part of the local vegetation during the Holocene, as is common in African savannas. Grazer enamel isotopes and short cell grass phytoliths records show changing proportions of C₃ and C₄ grasses but cannot distinguish between the drivers and their interactions.

Dynamic Global Vegetation Models (DGVMs), which simulate climate-driven vegetation processes, can generate spatially-continuous reconstructions of past environments to complement discrete and dispersed proxy records. They also allow insight into past biosphere–atmosphere interactions using the body of ecophysiological theory embedded in DGVMs. However, it is critical that model output be regarded in the context of independent palaeoecological records to build confidence in their findings. After knowing the state of past environments, it is imperative that we identify and firmly understand the processes that led to their formation. In this study, we use the LPX-DGVM²¹ driven by Atmosphere–Ocean coupled General Climate Model (AOGCM) climate output²², to test three possible drivers forcing vegetation change that have been suggested in the Wonderwerk Cave palaeoenvironmental study¹⁸: (1) atmospheric CO₂ levels, (2) rainfall seasonality, and (3) disturbance through fire. Under identical glacial climate reconstructions from four different AOGCM outputs, we modulated CO₂ levels to determine its effect on vegetation cover, particularly the expansion or contraction of C₃ and C₄ grasses. We test if the proposed drivers were influencing the changes that have been reconstructed from proxy records for the Early- and mid-Pleistocene in the southern Kalahari at c. 27° southern latitude. Model runs were performed under 150 ppm (post-MPT scenario) and 250 ppm (pre-MPT scenario) glacial CO₂ conditions, with and without the possibility of natural fires occurring (Supplementary Fig. S1 and S2). We used extreme values on both ends for the CO₂ values to explore the maximum sensitivity of the system. From the results, we estimate the effect of CO₂ concentration on C₃/C₄ grass cover, determine the climatic niches of C₃/C₄ grasses and elucidate ecophysiological mechanisms to explain low CO₂-mediated C₄ grass expansion.

Consistent with our hypothesis, model reconstructions suggest that CO₂ levels have a clear influence on C₄ grass cover over Africa. However, C₄ grass cover is most significantly impacted over East Africa and to a smaller degree over Southern Africa (Fig. 2). For all four LGM climate reconstructions, the low CO₂ (150 ppm) scenario, tends to expand and increase the density of C₄ grass cover. This trend is consistent for runs with and without fire, suggesting that fire has little influence on the competition between grass types within the model in this context (Supplementary Fig. S1 and S2). The large difference in C₄ grass cover particularly around the central African regions is a result of reduced tree cover instead of direct competition with C₃ grass, as tree cover is sensitive to fire regime changes. There is significant inter-climate model variation in Southern Africa, where the FGOALS-g1.0 reconstructions show clearest an increase in the western half of Southern Africa in C₃ grass cover and decrease in the eastern half of Southern Africa in the pre-MPT scenario compared to the post-MPT scenario (Fig. 2). In contrast, the other models show a more uniform increase in C₃ grass cover due to low CO₂, except for the eastern coastal areas (Fig. 2). These differences can be traced back to details in rainfall and temperature differences in the models, as shown in Supplementary Fig. S3. Figure 3 shows grass cover in the climate space of mean annual temperature (MAT) and mean annual precipitation (MAP), showing that C₄ cover tends to increase when MAT is above 15 °C and MAP is above 500 mm/year in areas dominated by concentrated summer rainfall. At low CO₂, this border becomes more prominent, while at the same time, less aridity is needed for a shift to occur.




Figure 2. Foliage projective cover of (a) C_3 and (b) C_4 grasses at 150 ppm and 250 ppm, and their difference derived from vegetation reconstructions simulated by LPX-DGVM²¹. In the difference plot, green indicates increases and purple indicates decreases. All models are shown with the impact of fire on in the model. The figure was constructed using raster2.8–19 (<https://CRAN.R-project.org/package=raster>) and mapproj1.2.6 (<https://CRAN.R-project.org/package=mapproj>) in R 3.5.2 (<https://www.R-project.org/>). The present-day coastline was obtained from mapsv3.3.0 (<https://CRAN.R-project.org/package=maps>), while LGM coastlines are based on data provided by PMIP2²².

The modelling results match well with the palaeoenvironmental changes seen in interior South Africa¹⁸. In the early Pleistocene, dry and wet phases with both C_3 and C_4 grasses exist. After the MPT, evidence for increased seasonality focussed on summer rainfall¹⁸ and significant groundwater changes in the Kuruman hills area^{19,20} indicate increasing aridity. In these arid, low glacial CO_2 conditions C_3 grasses, as with all C_3 plants, were more stressed than C_4 plants. On long timescales, C_4 grasses may have outcompeted C_3 grasses, which do not return in significant numbers during the Pleistocene^{24,25}. Low levels of CO_2 during glacials alone may not have led to major changes in vegetation in the interior of South Africa during the Pleistocene. However, the combined effects of low CO_2 in addition to changes in the growing season could have pushed vegetation through certain bioclimatic thresholds (Fig. 3). C_4 plants require on average over c. 22 °C growing season (warmest month) temperature to effectively compete against C_3 plants, limiting their distribution to warm tropical and subtropical areas^{11,26,27}. Below this ‘crossover temperature’, C_3 photosynthesis has a higher quantum yield¹¹. In glacial periods during the Pleistocene, when temperatures were lower in all seasons, the growing season in South Africa might have shifted when winter rainfall influences extended east^{2–4}. This would increase water availability at Wonderwerk Cave year-round, which could then support a mix of C_3 and C_4 grasses, as exists today in a very small area in the southern Karoo, close to the year-round rainfall zone²⁸. In addition, lower temperatures mean less evapotranspirative demands on plants, resulting in lower water stress. Changes in rainfall seasonality or amount alone without CO_2 changes cannot explain the long-term change to C_4 grass savannas as there is no return of a substantial amount of C_3 grasses in the humid late Pleistocene at Wonderwerk Cave²⁴. Our results have implications for hominin responses during the later phase of the Early Stone Age as seen in the rich Acheulean cultural record in the southern Kalahari, which contrasts with a much sparser mid- and late Pleistocene record^{24,29}. The much sparser archaeological record could be a reaction to the less vegetated landscape (less C_3 plants) and limited,




Figure 3. C₄ grass cover, with fire on, in climate space as simulated by LPX-DGVM²¹. Left is 150 ppm runs, right 250 ppm runs. (a–d) MAP (x-axis) versus MAT (y-axis), (e–f) seasonal concentration (distance from centre) and phase (direction) based on²³. Higher concentration means shorter rain season. Colours are: (a,b) C₄ grass coverage as % of land (c–f) % of grass that is C₄.

seasonal water availability. Major changes in the cultural record of the region, for example, the transition from the Early to the Middle Stone Age (c. 500–200 kyr), occur not at the start but during and after this ongoing environmental change.

Our study is an example of testing local terrestrial proxy reconstructions, as other areas of Africa clearly follow different trajectories (Fig. 2). Eastern parts of South Africa^{30,31} and East Africa^{32,33}, have a dominant C₄ grass component before the MPT. The impact of low CO₂ on the already existing C₄ grass cover here is shown as substantial in our modelling results. Our study shows how vegetation models are able to test hypotheses generated from local palaeoecological research targeting specific drivers and their interactions. However, any model output is always a reflection of larger patterns rather than a site-specific environmental reconstruction. It is noteworthy that in all reconstructions, C₄ grasses tend not to appear south of ~18°S, where C₃ grasses dominate regardless of CO₂ level (Fig. 2), even though C₄ grasses dominate the landscape there today²⁶. One reason is possibly the different winter rainfall expansion between the models, and C₃ grasses being a result of winter rainfall influence. However, the models only simulate winter rainfall for the far south-west of the continent (Fig. 4, Supplementary Fig. S3). Another could be the models lower temperature limits for C₄ photosynthesis (see “Methods”) interacting with possible local temperature biases in climate model simulation. More attention has to be paid in future work on such discrepancies between vegetation models and modern environmental data. In turn, this can be used to refine the models for use in past as well as in future climate change predictions.

Methods

Model description The Land surface Processes and eXchanges (LPX) is a coupled, process-based fire-enabled dynamic global vegetation model (DGVM). Full details of the dynamic vegetation component can be found in Sitch et al. (2003)³⁵, while the fire model is described in Thonicke et al. (2010)³⁶ and Prentice et al.²¹. LPX simulates atmosphere-to-biosphere interactions and ecosystem structure and function; computing spatially and temporally resolved estimates of potential vegetation cover and height, biomass, soil carbon and water and energy fluxes. LPX uses nine Plant Functional Types (PFT) to represent potential vegetation. PFTs compete within grid cells, where their differential ecophysiological responses to driving climate data, background turnover (or mortality) and resistance to fire disturbance determine their relative abundances. PFTs can be either tree or grasses. Trees are split by climate range (tropical, temperate, boreal), leaf type (broadleaf, evergreen), and phenological response (evergreen, raingreen, summergreen) and grasses are split between by C₃ and C₄ photosynthetic pathways.

To represent CO₂ fertilization, LPX explicitly couples CO₂ assimilation with transpiration using the photosynthesis-water balance scheme^{37,38} adapted from the Farquhar model^{39,40}. Available CO₂ reduces potential water stress on a plant by lowering the required stomatal conductance (g_c) for a given photosynthetic rate. Maximum potential stomatal conductance (g_{cMax}) for when water is not limiting depends on the maximum potential daytime pathway-specific photosynthetic assimilation rate (A_{max}) and minimum canopy conductance parameters, and atmospheric CO₂ concentration. If g_{cMax} requires a transpiration rate (D) that is greater than the available water supply (S , which is a function of soil water content and soil properties), then g_c (and therefore photosynthesis and production) is reduced in such a way as to be consistent with an empirical formulation derived from Monteith (1995)⁴¹. At higher atmospheric CO₂ concentrations, g_{cMax} decreases while A_{max} remains the same, and the value of S that induces water stress is, therefore, lower and maximum production rates can occur at lower moisture availability. Furthermore, when S is less than D , g_c (and therefore production) requires less down-regulation. Increased CO₂ thus leads to a fertilization effect, with increases production in drier conditions.

The distribution of C₃ versus C₄ plants is dynamically determined by light-use competition between the two photosynthetic pathways³⁵. The ratio of internal to ambient CO₂ concentrations optimal for photosynthesis is lower in C₄ compared to C₃, thereby reducing photorespiration and making photosynthesis more efficient. It also means that C₄ productivity is less sensitive to the lowering of atmospheric CO₂ concentrations. Leaf respiration as a fraction of Rubisco capacity is, however, higher for C₄ photosynthesis.

Under plentiful water supply and present-day CO₂, C₄ plants tend to have higher Gross Primary Production (GPP) than C₃ plants at temperatures above ~20 °C³⁶. Increased water stress, induced by dry conditions or reduced CO₂ concentrations, tends to reduce this crossover temperature. Under cooler temperatures, the simulated higher metabolic costs of C₄ photosynthesis outway advantages in photorespiration, and C₃ gains a competitive edge. Additionally, C₃ has no lower temperature limit for survival—through reduced productivity will effectively prevent establishment at very low temperatures, this is well those considered in this study. C₄ grasses, however, do not survive temperatures of less than 15.5 °C.

Fires occur from lightning ignitions, and the probability of an ignition event causing a fire depends on local fuel and atmospheric moisture content. Fire spread, flame height and residence time are based on weather conditions and fuel moisture, calculated using the Rothermel equations⁴². The area affected by fires is the product of the number of fires and the average spread of fire. LPX simulates fire mortality through two pathways: leaf and crown scorching, affecting all PFTs and cambial damage, affecting just tree PFTs. The amount of crown scorching depends on the height and intensity of the fire in relation to the height of the local vegetation. The probability of mortality from crown scorch increases as flame height increases beyond the canopy height of each PFT. While the fire mortality in this scheme will kill the shorter grass PFTs more than tree PFTs, grass PFTs tend to recover faster. Depending on other environmental conditions, water supply and CO₂ concentration, differing growth rates will preferentially select either C₃ and C₄ grass PFTs.

Modelling Protocol We follow the same modelling protocol as⁴³. Four distinct reconstructions of LGM climate (MIROC3.2, HadCM3M2, FGOALS-g1.0, CNRM-CM33)⁴⁴, generated by four atmosphere–ocean general circulation models (AOGCM) derived from the paleoclimate model intercomparison project 2 (PMIP2), were used to drive the Land surface Processes and eXchanges (LPX) DGVM²¹. Vegetation was ‘spun-up’ from bare ground




Figure 4. Mean annual precipitation, mean annual temperature and temperature during peak rainfall. Peak rainfall is equivalent to the growing season. Present day climate conditions taken from CRU TS 4.01³⁴. Peak rainfall is based on the month of the phase (Supplementary Fig. S3). Contours at mean annual temperatures of 15.5 °C and peak rainfall temperatures of 20 °C. The figure was constructed using raster 2.8–19 (<https://CRAN.R-project.org/package=raster>) and mapproj1.2.6 (<https://CRAN.R-project.org/package=mapproj>) in R 3.5.2 (<https://www.R-project.org/>). The present-day coastline was obtained from maps3.3.0 (<https://CRAN.R-project.org/package=maps>), while LGM coastlines are based on data provided by PMIP2²².

for 4000 years to reach equilibrium from which model runs were set for an additional 1380 years. Raw model output was then post-processed to show distributions of vegetation and climate-vegetation relationships as per⁴³.

Seasonal comparison We use season phase and concentration metrics from²³. Each month, m , was represented by a vector with direction (θ_m) corresponding to the time of year and the length corresponding to the magnitude of the variable for that month. A mean vector was calculated the average of x and y vectors:

$$\begin{aligned}\theta_m &= 2(m - 1)/12 \\ x &= \sum_m [v_m \times \cos(\theta_m)] \\ y &= \sum_m [v_m \times \sin(\theta_m)]\end{aligned}$$

The ratio of the mean vector length to annual average described the seasonal concentration (C) of the variable, timing is described by the mean direction (P):

$$C = \sqrt{x^2 + y^2} / \sum_m v_m \quad (1)$$

$$P = \arctan(x/y) \quad (2)$$

C is equal to 1 when the variable is 0 in all but 1 month (i.e., there is only 1 month of rainfall when assessing rainfall seasonality) and P is equal to that month. C is 0 and P is undefined when the variable is evenly distributed throughout the year.

Data availability

Available from the authors on request.

Code availability

Available from the authors on request.

Received: 26 May 2020; Accepted: 2 September 2020

Published online: 01 October 2020

References

- Mucina, L. & Rutherford, M. C. *The Vegetation of South Africa, Lesotho and Swaziland* (South African National Biodiversity Institute, Pretoria, 2006).
- van Zinderen Bakker, E. M. The evolution of late Quaternary paleoclimates of Southern Africa. *Palaeoecol. Afr.* **9**, 160–202 (1976).
- Cockcroft, M. J., Wilkinson, M. J. & Tyson, P. D. The application of a present-day climatic model to the late Quaternary in southern Africa. *Clim. Change* **10**, 161–181 (1987).
- Chase, B. M. & Meadows, M. E. Late Quaternary dynamics of southern Africa's winter rainfall zone. *Earth Sci. Rev.* **84**(3), 103–138 (2007).
- Bistinas, I., Harrison, S. P., Prentice, I. C. & Pereira, J. M. C. Causal relationships vs. emergent patterns in the global controls of fire frequency. *Biogeosciences* **11**, 5087–5101 (2014).
- Hoetzel, S., Dupont, L., Schefuß, E., Rommerskirchen, F. & Wefer, G. The role of fire in Miocene to Pliocene C 4 grassland and ecosystem evolution. *Nat. Geosci.* **6**(12), 1027–1030 (2013).
- Bond, W. J., Woodward, F. I. & Midgley, G. F. The global distribution of ecosystems in a world without fire. *New Phytol.* **165**(2), 525–538 (2005).
- Ripley, B. *et al.* Fire ecology of C3 and C4 grasses depends on evolutionary history and frequency of burning but not photosynthetic type. *Ecology* **96**(10), 2679–2691 (2015).
- Pinto, H., Sharwood, R. E., Tissue, D. T. & Ghannoum, O. Photosynthesis of C₃, C₃-C₄, and C₄ grasses at glacial CO₂. *J. Exp. Bot.* **65**(13), 3669–3681 (2014).
- Roth-Nebelsick, A. & Konrad, W. Habitat responses of fossil plant species to palaeoclimate—possible interference with CO₂?. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **467**, 277–286 (2017).
- Ehleringer, J. R., Cerling, T. E. & Helliker, B. R. C₄ photosynthesis, atmospheric CO₂, and climate. *Oecologia* **112**(3), 285–299 (1997).
- Edwards, E. J., Osborne, C. P., Strömberg, C. A., Smith, S. A. & C4 Grasses Consortium. The origins of C4 grasslands: integrating evolutionary and ecosystem science. *Science* **328**(5978), 587–591 (2010).
- Hönisch, B., Hemming, N. G., Archer, D., Siddall, M. & McManus, J. F. Atmospheric carbon dioxide concentration across the mid-Pleistocene transition. *Science* **324**(5934), 1551–1554 (2009).
- Yan, Y. *et al.* Two-million-year-old snapshots of atmospheric gases from Antarctic ice. *Nature* **574**(7780), 663–666 (2019).
- Faith, J. T., Rowan, J. & Du, A. Early hominins evolved within non-analog ecosystems. *Proc. Natl. Acad. Sci.* **116**(43), 21478–21483 (2019).
- Sealy, J., Naidoo, N., Hare, V. J., Brunton, S. & Faith, J. T. Climate and ecology of the palaeo-Agulhas Plain from stable carbon and oxygen isotopes in bovid tooth enamel from Nelson Bay Cave, South Africa. *Quat. Sci. Rev.* **235**, 105974 (2019).
- Horwitz, L. K. & Chazan, M. Past and present at Wonderwerk Cave (Northern Cape Province, South Africa). *Afr. Archaeol. Rev.* **32**(4), 595–612 (2015).
- Ecker, M. *et al.* The palaeoecological context of the Oldowan-Acheulean in southern Africa. *Nat. Ecol. Evol.* **2**(7), 1080–1086 (2018).
- Matmon, A. *et al.* New chronology for the southern Kalahari Group sediments with implications for sediment-cycle dynamics and early hominin occupation. *Quat. Res.* **84**(1), 118–132 (2015).
- Vainer, S., Erel, Y. & Matmon, A. Provenance and depositional environments of Quaternary sediments in the southern Kalahari Basin. *Chem. Geol.* **476**, 352–369 (2018).
- Prentice, I. C. *et al.* Modeling fire and the terrestrial carbon balance. *Glob. Biogeochem. Cycles* **25**(3), 2–13 (2011).
- Braconnot, P. *et al.* Results of PMIP2 coupled simulations of the Mid-Holocene and Last Glacial Maximum-Part 1: experiments and large-scale features. *Clim. Past* **3**(2), 261–277 (2007).
- Kelley, D. I. *et al.* A comprehensive benchmarking system for evaluating global vegetation models. *Biogeosciences* **10**(5), 3313–3340 (2013).

24. Chazan, M. *et al.* Archaeology, paleoenvironment and chronology of the early middle stone age component of Wonderwerk cave in the interior of southern Africa. *J. Palaeolithic Archaeol.* <https://doi.org/10.1007/s41982-020-00051-8> (2020).
25. Lee-Thorp, J. A. & Beaumont, P. B. Vegetation and seasonality shifts during the late Quaternary deduced from 13C/12C ratios of grazers at Equus Cave, South Africa. *Quat. Res.* **43**, 426–432 (1995).
26. Vogel, J. C. The geographical distribution of Kranz species in southern Africa. *South Afr. J. Sci.* **75**, 209–215 (1978).
27. Zhou, H., Helliker, B. R., Huber, M., Dicks, A. & Akçay, E. C₄ photosynthesis and climate through the lens of optimality. *Proc. Natl. Acad. Sci.* **115**(47), 12057–12062 (2018).
28. Rubin, F., Palmer, A. R. & Tyson, C. Patterns of endemism within the Karoo National Park, South Africa. *Bothalia* **31**(1), 117–133 (2001).
29. Walker, S. J., Lukich, V. & Chazan, M. Kathu townlands: a high density earlier stone age locality in the interior of South Africa. *PLoS ONE* **9**(7), e103436 (2014).
30. Lee-Thorp, J. A., Sponheimer, M. & Luyt, J. Tracking changing environments using stable carbon isotopes in fossil tooth enamel: an example from the South African hominin sites. *J. Hum. Evol.* **53**(5), 595–601 (2007).
31. Codron, D., Brink, J. S., Rossouw, L. & Clauss, M. The evolution of ecological specialization in southern African ungulates: competition- or physical environmental turnover?. *Oikos* **117**, 344–353 (2008).
32. Plummer, T. W. *et al.* The environmental context of Oldowan hominin activities at Kanjera South, Kenya. In *Interdisciplinary approaches to the Oldowan* (eds Hovers, E. & Braun, D. R.) 149–160 (Springer, Berlin, 2009).
33. Cerling, T. E. *et al.* Dietary changes of large herbivores in the Turkana Basin, Kenya from 4 to 1 Ma. *Proc. Natl. Acad. Sci.* **112**(37), 11467–11472 (2015).
34. Harris, I., Osborn, T. J., Jones, P. & Lister, D. Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Sci. Data.* <https://doi.org/10.1038/s41597-020-0453-3> (2020).
35. Sitch, S. *et al.* Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Glob. Change Biol.* **9**(2), 161–185 (2003).
36. Thonicke, K. *et al.* The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model. *Biogeosciences* **7**(6), 1991–2011 (2010).
37. Haxeltine, A. & Prentice, I. C. BIOME3: an equilibrium terrestrial biosphere model based on ecophysiological constraints, resource availability, and competition among plant functional types. *Glob. Biogeochem. Cycles* **10**(4), 693–709 (1996).
38. Haxeltine, A. & Prentice, I. C. A general model for the light-use efficiency of primary production. *Funct. Ecol.* **10**, 551–561 (1996).
39. Farquhar, G. D., Von Caemmerer, S. & Berry, J. A. A biochemical model of photosynthetic CO₂ assimilation in leaves of C₃ plants. *Planta* **149**, 78–90 (1980).
40. Farquhar, G. D. & Von Caemmerer, S. Modelling of photosynthetic response to environmental conditions. In *Physiological Plant Ecology II: Water Relations and Carbon Assimilation* (eds Nobel, P. S. *et al.*) 549–587 (Springer, Berlin, 1982).
41. Monteith, J. L. A reinterpretation of stomatal responses to humidity. *Plant Cell Environ.* **18**, 357–364 (1995).
42. Rothermel, R. C. A Mathematical Model for Predicting Fire Spread in Wildland Fuels (Vol. 115). Intermountain Forest and Range Experiment Station, Forest Service, US Department of Agriculture (1972).
43. Sato, H., Kelley, D. I., Mayor, S. J., Cowling S. A., Calvo, M. M. & Prentice, I. C. Fire and low CO₂ opened dry corridors in South America during the Last Glacial Maximum. Under Review for Nature Geosciences: NGS-2019-07-01558B (2020).
44. Prentice, I. C., Harrison, S. P. & Bartlein, P. J. Global vegetation and terrestrial carbon cycle changes after the last ice age. *New Phytol.* **189**(4), 988–998 (2011).

Acknowledgements

M.E. has received funding from the European Union's Framework Programme for Research and Innovation Horizon 2020 (2014–2020) under the Marie Skłodowska-Curie Grant Agreement No. 837730. The contribution by D.K. was supported by the UK Natural Environment Research Council through The UK Earth System Modelling Project (UKESM, Grant No. NE/N017951/1).

Author contributions

M.E. and H.S. designed research. H.S. and D.K. performed research. M.E., H.S. and D.K. wrote the paper.

Funding

Open Access funding provided by Projekt DEAL.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://doi.org/10.1038/s41598-020-72614-2>.

Correspondence and requests for materials should be addressed to M.E.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020