



OPEN

Author Correction: MC1R is dispensable for the proteinuria reducing and glomerular protective effect of melanocortin therapy

Yingjin Qiao, Anna-Lena Berg, Pei Wang, Yan Ge, Songxia Quan, Sijie Zhou, Hai Wang,
Zhangsuo Liu & Rujun Gong

Correction to: *Scientific Reports*, <https://doi.org/10.1038/srep27589>, Published online 08 June 2016

This Article contains errors.

In Figure 8a, at time 0 h, the images for Control e^{-}/e^{-} , LPS WT, and LPS e^{-}/e^{-} , were inadvertently taken from the Control WT condition. The correct Figure 8a is now included below as Figure 1.

The errors do not affect the results or conclusions of the study.

Published online: 21 September 2020




Figure 1. (a) Phase-contrast micrographs were taken immediately after wounding (0 h) and after migration for 24 h. Scale bar = 100 μm . Cell morphology at 24 h was taken under high-power fields. LPS injury resulted in marked podocyte shrinkage and this effect was abrogated by NDP-MSH. Scale bar = 20 μm .



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020