OPEN

Author Correction: A System of Cytokines Encapsulated in ExtraCellular Vesicles

Wendy Fitzgerald, Michael L. Freeman , Michael M. Lederman, Elena Vasilieva, Roberto Romero & Leonid Margolis

Published online: 29 October 2020

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-018-27190-x>, published online 12 June 2018

In this Article, the legend of Figure 2 is incorrect:

“Distribution of cytokines between the surface and the inner volume of EVs. Fractions of total EV-associated cytokines $\pm$ SEM. Samples collected as in Fig. 1. (a) Placental villous explants n = 10; (b) Amnion explants, n = 10; (c) Tonsillar explants n = 5; (d) Cervical explants n = 6; (e) PPP from healthy donors n = 52; (f) Amniotic fluid from 3 donors (g) T cells n = 6; (h) Monocytes n = 6. *Indicates significant difference $p < 0.05$ between surface and encapsulated cytokines.”

should read:

“Distribution of cytokines between the surface and the inner volume of EVs. Fractions of total EV-associated cytokines $\pm$ SEM. Samples collected as in Fig. 1. (a) Placental villous explants n = 10; (b) Amnion explants, n = 10; (c) Tonsillar explants n = 5; (d) Cervical explants n = 6; (e) PPP from healthy donors n = 52; (f) T cells n = 6; (g) Monocytes n = 6; (h) Amniotic fluid from 3 donors. *Indicates significant difference $p < 0.05$ between surface and encapsulated cytokines.”

Consequently, in the Results section under the subheading ‘Cytokines associated with EVs: surface-bound vs. encapsulated’, the referrals to Figure 2 are incorrect.

“In contrast to plasma, in amniotic fluid out of all EV-associated cytokines only two, IL-17 and IFN- γ , were preferentially on the surface of EVs and only small fractions of another three, IL-1 α , IL-22, and ITAC, were present on the surface of EVs; whereas 27 other cytokines were almost exclusively inside EVs (Fig. 2f).

For T cells (Fig. 2g) and monocytes (Fig. 2h) fewer cytokines were on the surface of EVs, whereas for both types of cells EV-associated cytokines were predominantly inside the vesicles (25 and 20 respectively).”

should read:

“In contrast to plasma, in amniotic fluid out of all EV-associated cytokines only two, IL-17 and IFN- γ , were preferentially on the surface of EVs and only small fractions of another three, IL-1 α , IL-22, and ITAC, were present on the surface of EVs; whereas 27 other cytokines were almost exclusively inside EVs (Fig. 2h).

For T cells (Fig. 2f) and monocytes (Fig. 2g) fewer cytokines were on the surface of EVs, whereas for both types of cells EV-associated cytokines were predominantly inside the vesicles (25 and 20 respectively).”

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2020