scientific reports

Check for updates

OPEN

Author Correction: Anisomycin prevents OGD-induced necroptosis by regulating the E3 ligase CHIP

Published online: 28 December 2021

Mi-bo Tang, Yu-sheng Li, Shao-hua Li, Yuan Cheng, Shuo Zhang, Hai-yang Luo, Cheng-yuan Mao, Zheng-wei Hu, Jonathan C. Schisler, Chang-he Shi & Yu-ming Xu

Correction to: *Scientific Reports* https://doi.org/10.1038/s41598-018-24414-y, published online 23 April 2018

This Article contains an error in Figure 1A. As a result of a mistake in figure assembly, the band for p-MLKL is incorrect. The correct Figure 1 and accompanying legend appear below.

This change does not affect the conclusions of the Article.



Figure 1. OGD challenge induces necroptotic cell death. (**A–B**) Representative western blot, with β-actin used for normalization. RIPK1, RIPK3, p-RIPK3 and p-MLKL in control cells, cells challenged by OGD which were pre-treated with or without Nec-1 (50 μM) were tested. The bars represent the mean ± SEM of five independent experiments. Significant differences $^{*}p < 0.05$ vs. Control, $^{\#}p < 0.05$ vs. OGD. Related blots are shown in Supplementary Fig. S1.