




OPEN

# Author Correction: Both higher fitness level and higher current physical activity level may be required for intramyocellular lipid accumulation in non-athlete men

Nozomu Yamasaki, Yoshifumi Tamura, Kageumi Takeno, Saori Kakehi, Yuki Someya, Takashi Funayama, Yasuhiko Furukawa, Hideyoshi Kaga, Ruriko Suzuki, Daisuke Sugimoto, Satoshi Kadowaki, Motonori Sato, Takashi Nakagata, Miho Nishitani-Yokoyama, Kazunori Shimada, Hiroyuki Daida, Shigeki Aoki, Hiroaki Satoh , Ryuzo Kawamori & Hirotaka Watada

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-020-61080-5>, published online 05 March 2020

The original version of this Article contained a typographical error in the spelling of the author Hiroaki Satoh, which was incorrectly given as Hiroaki Sato. This has now been corrected in the PDF and HTML versions of the Article.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021