



OPEN **Publisher Correction:** The impact of smoking different tobacco types on the subgingival microbiome and periodontal health: a pilot study

Sausan Al Kawas, Farah Al-Marzooq, Betul Rahman, Jenni A. Shearston, Hiba Saad, Dalenda Benzina & Michael Weitzman

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-020-80937-3>, published online 13 January 2021

This Article contains errors in Figure 4 where the horizontal and vertical axis labels are missing. The correct Figure 4 appears below as Figure 1.

Published online: 03 May 2021

Folds change in the relative abundance





Figure 1. Significant log₂ fold change of genera abundances in cigarette, shisha, or medwakh smoking groups compared to non-smokers (log₂ tobacco vs log₂ non-smokers). Red indicates an increase and blue indicates a decrease in the relative abundance of each genus compared to non-smokers. *Significant difference between each tobacco group and non-smokers; #significant difference exclusively in patients with severe periodontitis; M medwakh smokers, C cigarette smokers, SH shisha smokers.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021