



OPEN

Production of basil (*Ocimum basilicum* L.) under different soilless cultures

El-Sayed Khater^{1✉}, Adel Bahnasawy¹, Wael Abass², Osama Morsy², Hossam El-Ghobashy³, Yousry Shaban³ & Mohsen Egela³

The main aim of this paper was to investigate the possibility of growing basil under three soilless systems (aeroponic, hydroponic and peatmoss slab systems). A model was developed to predict the nutrients consumption by basil plants. Shoot and root height, fresh and dry mass of whole plant, nutrients uptake, and oil content were studied during the growth period (after 4 and 7 weeks from transplanting). The results indicated that the shoot lengths of basil plants were 71.67 ± 2.89 , 65.67 ± 1.15 and 62.33 ± 2.31 cm at the end of growth period for aeroponic, hydroponic and peatmoss slabs, respectively. The highest value of root height of basil plants was 37.67 ± 6.66 cm for aeroponic system. The dry mass of shoot of basil plants ranged from 28.48 ± 0.91 to 44.77 ± 0.97 and 72.98 ± 0.83 to 117.93 ± 1.40 g plant⁻¹ after 4 and 7 weeks from transplanting, respectively. The highest values of the N, P, K, Ca and Mg uptakes were 753.99 ± 5.65 , 224.88 ± 3.05 , 449.75 ± 4.59 , 529.12 ± 6.63 and 112.44 ± 1.67 mg plant⁻¹ at the end of experimental period, respectively. The basil oil content ranged from 1.129 (1.11%) to 2.520 (1.80%) and 2.664 (1.42%) to 6.318 (1.44%) g plant⁻¹ after 4 and 7 weeks from transplanting, respectively at the same previous order. The production costs of basil plant were 2.93, 5.27 and 6.24 EGP kg⁻¹ of plant. The model results were in a reasonable agreement with the experimental ones.

There is an increasing interest recently for growing sweet basil (*Ocimum basilicum* L.) in greenhouse soilless culture, which offer a suitable condition for maximization of production^{1,2}. It is cultivated commonly in an open field with a variability in productivity and quality³.

Basil has high nutritional contents with low caloric values. It is used as a pharmacological raw material. Also, it contains vitamins A, B₆ and C as well as carotene besides calcium, potassium, phosphorus, magnesium, iron. Therefore, it needs a warm climate and high temperature and soil should be fecundity^{4,5}.

The advantages of the soilless culture are the earliest growth and higher yield compared to traditional culture. Also, this system assures an equal supply of nutrient solution, so it can obtain a homogeneous crop. The mineral elements concentration and composition are well adjusted. Also, the buffer capacity of nutrient solution is low. pH and mineral composition of solution are easily changeable. Soilless culture decreases the time of adjusting solution⁶.

Soilless cultivation systems provide plant management under controlled water and minerals supply of the nutrient solution with or without medium. There are three systems of soilless cultivation namely, system with solid medium, in a liquid medium and aerated medium^{7,8}.

Hydroponic system is a way plants without soil in water having a nutritional solution. The soil is used in traditional cultivation as a medium to add water and minerals in it, this soil is not needed in hydroponic because the minerals are added directly to water where the plants grow. It is more efficient to control water which can be reused after adjustment. It decreases the use of pesticides. It is used for many crops such as beets, radishes, carrots, potatoes, cereal crops, fruits, ornamentals and seasonal flowers can be grown on inert supporting medium instead of soil⁹⁻¹¹.

QI¹² reported that the aeroponic system is a type of growing plants in air or mist environment without using any soil. In hydroponic, plant's roots are growing in water with nutrients. But for aeroponic, the nutrients are added through mist spray by sprinkles to plant's roots. The aeroponic system consist of a pump, nozzles, and

¹Agricultural and Biosystems Engineering Department, Faculty of Agriculture, Benha University, P.O. Box 13736, Moshthor, Toukh, Kalubia, Egypt. ²Basic and Applied Science Department, College of Engineering and Technology, Arab Academy for Science and Technology and Maritime Transport (AASTMT), P.O. Box 2033, Cairo, Egypt. ³Institute of Agricultural Engineering Research, Agriculture Research Center, Doki, Giza, Egypt. ✉email: alsayed.khater@fagr.bu.edu.eg

growing chamber. There are a few types of aeroponic like low pressure type, high pressure type and commercial system. Basil is used as fresh and dried leaves a medicinal herb^{13,14} for its diuretic and stimulating properties and also used in perfume compositions¹⁵. Basil is growing better in soilless systems than conventional systems and many studies have used basil as aquaponic or hydroponic crop¹⁶.

The most severe problem in the hydroponic system and soilless is the root rot which is due to the low oxygen level in the nutrient solution, therefore, proper aeration is required to overcome this problem. Aeroponic system is the proper solution to provide the plant with the required oxygen and nutrients. Besides, demand of organic production is increasing day after day. Therefore, this study aimed to improve the basil production under three soilless systems.

Materials and methods

The experiment was conducted at Agricultural and Bio-Systems Engineering Department, Faculty of Agriculture Moshtohor, Benha University, Egypt (latitude 30° 21' N and 31° 13' E), during the period of May to July, 2019 season under the university guidelines and legislation. Basil seedlings were sown in the plastic cups (7 cm diameter and 7 cm height) filled with peat moss. The cups were irrigated daily using water with nutrient solution ($\text{Ca}(\text{NO}_3)_2$, 236 g L⁻¹, KNO_3 , 101 g L⁻¹, K_2SO_4 , 115 g L⁻¹, KH_2PO_4 , 136 g L⁻¹, MgSO_4 246 g L⁻¹ and chelates for trace elements into preacidified groundwater (from the following ppm concentration are achieved in this formulation: N = 210, P = 31, K = 234, Ca = 200, Mg = 48, S = 64, Fe = 14, Mn = 0.5, Zn = 0.05, Cu = 0.02, B = 0.5, Mo = 0.01)). Two weeks old basil seedlings were planted at 9.0 plant m⁻² in the experimental tanks. These seedlings were planted according to the permission of Benha university rules and legislation.

Culture systems description. Figure 1a,b show the experimental setup. It shows the system which consists of hydroponic system, aeroponic system, soilless substrate, solution system and pumps.

The hydroponic system (Deep Water Culture (DWC)) consists of three rectangular polyethylene tanks that used for basil plants culture. Dimensions of each tank are 80 cm long, 40 cm wide and 30 cm high. The slope of hydroponic tanks was 2% and stand 1 m high above the ground. The hydroponic tanks were covered with foam boards to support the plants. Each hydroponic tank provided with an air blower (Model NS 780—Flow Rate 850 L h⁻¹—Head 1.5 m—Power 15 W, China) to increase dissolved oxygen concentrations. The solution was circulated by a pump (Model First QB60—Flow Rate 30 L min⁻¹—Head 25 m—Power 0.5 hp, China) from the solution tank to the upper ends of the hydroponic tanks. Small tubes (16 mm) were used to provide tanks with solution in a closed system.

Aeroponic system consists of three rectangular polyethylene tanks that used for basil plants culture. Dimensions of each tank are 80 cm long, 40 cm wide and 50 cm high. The aeroponic tanks were established 1 m above the ground. Each aeroponic tank was divided into two parts, the lower part was made from polyethylene and the upper part was made from wood. The aeroponic tanks were covered with foam boards to support the plants. Each aeroponic tank was provided with two fog nozzles (Model M3MJWT5M – Orifice 2 mm – Discharge 8 L h⁻¹, India) located at the bottom of the tank sprayed nutrient solution into the tank in order to keep the roots wet. Small tubes (16 mm) were used to provide aeroponic tank with solution in a closed system.

Soilless substrates consist are placed in three rows are 2 m long. Each row consists standard peat moss slabs (1.00 m × 0.20 m × 0.075 m). Basil plants were placed on row peat moss slabs with a drip irrigation system. There were three plants per slab giving a mean density of 9.0 plant m⁻². Each plant was fed by a single drip.

The circular polyethylene tank of the nutrient solution system 500 L capacity was used for collecting the drained solution by gravity from the ends of the three systems. The nutrient solutions were prepared manually once per ten days^{17,18} by dissolving appropriate amounts of $\text{Ca}(\text{NO}_3)_2$, 236 g L⁻¹, KNO_3 , 101 g L⁻¹, K_2SO_4 , 115 g L⁻¹, KH_2PO_4 , 136 g L⁻¹, MgSO_4 246 g L⁻¹ and chelates for trace elements into preacidified groundwater (from the following ppm concentration are achieved in this formulation: N = 210, P = 31, K = 234, Ca = 200, Mg = 48, S = 64, Fe = 14, Mn = 0.5, Zn = 0.05, Cu = 0.02, B = 0.5, Mo = 0.01). pH and Electrical Conductivity (EC) were further adjusted to 6.5–7.0 and 1.4–1.8 dS m⁻¹, respectively, after salt addition. The average air ambient temperature was 25.97 ± 4.37 °C and the average water temperature was 24.03 ± 3.92 °C. The average relative humidity was 65.4% and the light intensity was 338.55 ± 40.06 W m⁻².

Measurements. Three plants sample were taken during the vegetative and flowering stages (four and seven weeks after transplanting, respectively) for growth measurement and chemical analysis. Plant height, root length and the fresh and dry weight of leaves, stems and roots were determined. After measuring fresh mass, the plants were oven dried at 65 °C until constant weight was reached¹⁹. Total content of macro elements was evaluated after being digested²⁰. Nitrogen was determined by Kjeldahl digestion methods²¹. Potassium, Calcium and magnesium were determined by Photofatometer (Model Jenway PFP7—Range 0—160 mmol L⁻¹, USA) and phosphorus (P) was determined colorimetrically method²². The content of oil was determined in different organs: leaves, stems and inflorescences according to²³.

Water samples were taken, at inlet and outlet of the culture units for measuring nitrogen (N), phosphorus (P), potassium (K), calcium (Ca) and magnesium (Mg) were measured every week at 10 am during the experimental period.

Total production cost. The cost calculation based on the following parameters was also performed:

Fixed costs (Fc). Depreciation costs (D_c)





Figure 1. (a) The experimental setup. (b) Images of system.

$$D_c = \frac{P_d - S_r}{L_d} \tag{1}$$

where D_c is the depreciation cost, EGP (Egyptian pound) year⁻¹. ($\$ = 15.63$ EGP). P_d is the system price, EGP. S_r is the salvage rate (0.1 P_d) EGP. L_d is the system life, year.

Interest costs (I_n):

$$I_n = \frac{P_d + S_r}{2} \times i_n \tag{2}$$

where I_n is the interest, EGP year⁻¹. i_n is the interest as compounded annually, decimal (12%). Shelter, taxes and insurance costs (S_i).

Shelter, taxes and insurance costs were assumed to be 3% of the purchase price of the automatic feeder (P_m). Then:

$$\text{Fixed cost} = D_c + I_n + 0.03 P_m / \text{hour of use per year} \tag{3}$$

Variable (operating) costs (Vc). Repair and maintenance costs (R_m):

$$R_m = 100\% \text{ depreciation cost/hour of use per year} \quad (4)$$

Energy costs (E):

$$E = EC \times EP \quad (5)$$

where E is the energy costs, EGP h⁻¹. EC is the electrical energy consumption, kWh. EP is the energy price, 0.57 EGP kW⁻¹.

Labor costs (L_a):

$$L_a = \text{Salary of one worker} \times \text{No. of workers} \quad (6)$$

where L_a is the Labor costs, EGP h⁻¹. Salary of one worker = 10 EGP h⁻¹. No. of workers = 1.

Then:

$$\text{Variable costs} = R_m + E + L_a \quad (7)$$

Total costs (T_c).

$$\text{Total costs} = \text{Fixed costs} + \text{Variable costs} \quad (8)$$

Table 1 shows the input parameters of calculate total production costs of basil plants grown in different soil-less systems.

Nutrients consumption rate. The Nutrients consumption rate were calculated as the differences between the nutrients at inlet and outlet of culture units by the following formula²⁴:

$$C_{Nc} = \frac{Nc_{in} - Nc_{out}}{\text{Number of plants}} \times Q \times 24 \quad (9)$$

where C_{Nc} is the nutrients consumption rate, mg day⁻¹ plant⁻¹. Nc_{in} is the nutrients at inlet of the hydroponic unit, mg L⁻¹. Nc_{out} is the nutrients at outlet of the hydroponic unit, mg L⁻¹. Q is the discharge, L h⁻¹.

Model development of nutrient consumption. Model assumptions:

- N, P, K, Ca and Mg are the nutrients used in study.
- The plants are uniformity distributed in the solution, so they work as a uniform sink for water and minerals with space at any time.
- The root systems are uniformly dispersed in the solution with uniform root length density at any time.
- The whole root system uptake characteristics are uniform.
- Water losses by evaporation are negligible.

The simplest nutrient consumption models relate the nutrient consumption to the concentration gradient using some sort of proportionality factor such as root permeability or conductivity^{25,26}. The nutrient consumption was determined by using the following equation:

$$NC = a_{NC} \cdot \Delta C \quad (10)$$

where NC is the nutrient consumption, mg plant⁻¹ day⁻¹. ΔC is the concentration gradient, mg plant⁻¹ day⁻¹. a_{NC} is the proportionality factor, dimensionless.

A similar model of nutrient consumption takes into consideration the differing effects caused by variations in root growth stage. Assuming that growth follows a first order differential equation and assuming that the root growth is exponential²⁷, then Eq. (11) can be derived. This equation is presented in similar form to Eq. (10) and use the following equation:

$$NC = \left(\frac{C_{plant} - C_{plant0}}{A_r - A_{r0}} \right) \cdot \left(\frac{\ln \left(\frac{A_r}{A_{r0}} \right)}{t - t_0} \right) \cdot A_r \quad (11)$$

where C_{plant0} is the concentration of the nutrients in the plant at time t₀, mg plant⁻¹. A_r is the root surface area at time t, cm² plant⁻¹. A_{r0} is the root surface area at time t₀, cm² plant⁻¹.

Root surface area was calculated from root length and mean root radius using the following equation:

$$A_r = 2\pi r_0 L_r \quad (12)$$

The root length increment using the following equation²⁸:

$$\Delta L_r = \Delta DW_{root} v \quad (13)$$

where ΔL_r is the root length increment, cm day⁻¹. ΔDW_{root} is the daily amount of root dry mass increment, g day⁻¹. v is the ratio of root length and mass of roots, cm g⁻¹.

The daily amount of dry weight of roots is calculated from the following equation²⁹:

Cost Item	Units	Production system		
		Aeroponics	Hydroponics	Peatmoss Slab
Fixed cost (EGP)				
Culture units	EGP kg ⁻¹	1.14	2.18	2.96
Pumps and fittings	EGP kg ⁻¹	0.19	0.36	0.59
Total fixed cost	EGP kg ⁻¹	1.33	2.54	3.55
Variable cost (EGP)				
Basil seedlings	EGP kg ⁻¹	0.57	1.09	1.33
Cups	EGP kg ⁻¹	0.11	0.22	0.27
Peat moss	EGP kg ⁻¹	0.07	0.13	0.16
Labor	EGP kg ⁻¹	0.10	0.10	0.10
Energy	EGP kg ⁻¹	0.23	0.43	0.27
Fertilizers	EGP kg ⁻¹	0.11	0.11	0.11
Chemicals	EGP kg ⁻¹	0.08	0.08	0.08
Total variable cost	EGP kg ⁻¹	1.24	2.16	2.32
Deprecation	EGP kg ⁻¹	0.26	0.47	0.56
Maintenance	EGP kg ⁻¹	0.05	0.05	0.05

Table 1. The input parameters of calculate total production costs of basil plants grown in different soilless systems.

$$\Delta DW_{root} = \begin{cases} 5LAI & \text{for } LAI \leq 0.5 \\ 2.5 + 23.9(LAI-0.5) & \text{for } LAI > 0.5 \end{cases} \quad (14)$$

where LAI is the leaf area index.

Leaf area index was changed in the same proportions as root length density to maintain a constant ratio between roots and shoots. The leaf area index is calculated from the following equation³⁰:

$$LAI = \frac{LAI_{max}}{1 + K_2 e^{-k_1 t}} \quad (15)$$

where LAI_{max} is the maximum leaf area index. K₂ and k₁ are the coefficients of the growth functions.

All computational procedures of the model were carried out using Excel spreadsheet. The computer program was devoted to mass balance for predicting the nutrients consumption. The differences between the predicted and measured values were evaluated using RMSE indicator (root means square error) which is calculated using the following equation:

$$RMSE = \sqrt{\frac{\sum (Predicted - Measured)^2}{n}} \quad (16)$$

The parameters used in the model that were obtained from the literature are listed in Table 2. Figure 2 shows flow chart of the model.

Statistical analysis. Three replicates of each treatment were allocated in a Randomize Complete Block Design (RCBD) in the system. Data were analyzed one-way ANOVA (analysis of variance) using statistical package for social sciences (spss v21). Means were separated using New Duncan Multiple Range Test (DMRT). Data presented are mean ± standard division (SD) of four replicates.

Results and discussion

Shoot length. Figure 3 shows the shoot length of basil plants grown in different soilless systems (Aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results indicate that the shoot in aeroponic was taller than those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the shoot length of basil plants were 62.00 ± 2.65, 57.83 ± 7.42 and 48.77 ± 2.89 cm for aeroponic, hydroponic and peatmoss slabs, respectively, after 4 weeks from transplanting. Meanwhile, they were 71.67 ± 2.89, 65.67 ± 1.15 and 62.33 ± 2.31 cm for aeroponic, hydroponic and peatmoss slabs, after 7 weeks from transplanting at the same previous. These results agreed with those obtained by³³ whose found that the plants grown aeroponically were twice as high as those in hydroponics and 4 times taller than those grown in sand. These previous results may be due that the roots of aeroponics systems are hanged in mid-air inside containers or chambers at 100% humidity and fed up a fine mist of nutrient solutions. This pervious system stimulates absorption of roots to much needed oxygen and nutrients, those increasing metabolism and rate of growth compared with soil³⁴.

Parameter	Units	Value	References
V	cm g ⁻¹	1.7	31
LAI _{max}	m ² m ⁻²	4.8	32
K ₂	-	500	30
k ₁	day ⁻¹	0.53	30

Table 2. The parameters used in the model.




Figure 2. Flow chart of nutrients consumption rate.




Figure 3. The shoot length of basil plants grown in different soilless systems.

The statistical analysis showed that the differences between the obtained data of shoot length due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant.




Figure 4. The root length of basil plants grown in different soilless systems.

Root length. Figure 4 shows the root length of basil plants grown in different soilless systems (Aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results of measurements of root of the plants grown in aeroponic system were taller than those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the highest value of root length of basil plants was 37.67 ± 6.66 cm for aeroponic system, while, the lowest value of root length of basil plants was 27.67 ± 0.58 cm was found with peatmoss slabs. The root length for basil plants grown in aeroponic system were 1.68 and 2.12 times taller than those grown in peatmoss slabs after 4 and 7 weeks from transplanting, respectively. These results agreed with those obtained by³⁵. Also, many studies showed that the aeroponic system enhance the rates of plants growth by promoting the root aeration because of the root system is grown totally suspended at the air, giving the plant stem and roots systems access to 100% of the available oxygen at the air³⁶. These results are in agreement with findings which were reported by³⁷ that they showed that plant root length, area, volume of aeroponic system were significantly exceeded the hydroponic and substrate systems.

The statistical analysis showed that the differences between the obtained data of root length due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant.

Fresh and dry mass of shoot. Figure 5a,b show the fresh and dry mass of shoot of basil plants grown in different soilless systems (Aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results indicate that the fresh and dry of shoot grown in aeroponic system were better than those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the fresh and dry mass of shoot of basil plants were 140.00 ± 13.76 , 139.02 ± 10.19 and 102.06 ± 35.54 g plant⁻¹ and 44.77 ± 0.97 , 32.36 ± 0.68 and 28.48 ± 0.91 g plant⁻¹ for Aeroponic, hydroponic and peatmoss slabs, respectively, after 4 weeks from transplanting. Meanwhile, the results also indicate that the fresh and dry mass of shoot of basil plants were 438.61 ± 42.61 , 229.33 ± 10.30 and 187.99 ± 24.84 g plant⁻¹ and 117.93 ± 1.40 , 77.85 ± 0.77 and 72.98 ± 0.83 g plant⁻¹ for aeroponic, hydroponic and peatmoss slabs, respectively, after 7 weeks from transplanting. We can explain those that the aeroponic system enhance the rates of plants growth by promoting the root aeration because of the root system is grown totally suspended at the air, giving the plant stem and roots systems access to 100% of the available oxygen at the air³⁸.

The statistical analysis showed that the differences between the obtained data of fresh mass of shoot due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant. Also, the statistical analysis showed that the differences between the obtained data of dry mass of shoot due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was non-significant.

Fresh and dry mass of root. Figure 6a,b show the fresh and dry mass of root of basil plants grown in different soilless systems (Aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results indicate that the fresh and dry of root grown in aeroponic system were better than those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the fresh and dry mass of root of basil plants were 150.52 ± 0.72 , 128.15 ± 2.32 and 49.17 ± 4.52 g plant⁻¹ and 39.11 ± 2.14 , 33.82 ± 1.57 and 24.73 ± 1.76 g plant⁻¹ for aeroponic, hydroponic and peatmoss slabs, respectively, after 4 weeks from transplanting. Meanwhile, the results also indicate that the fresh and dry mass of root of basil plants were 452.02 ± 8.94 , 337.97 ± 12.20 and 324.94 ± 5.48 g plant⁻¹ and 114.22 ± 5.05 , 97.16 ± 3.35 and 66.88 ± 2.36 g plant⁻¹ for aeroponic, hydroponic and peatmoss slabs, respectively, after 7 weeks from transplanting.

The statistical analysis showed that the differences between the obtained data of fresh mass of root due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant. Also, the statistical analysis showed that the differences between the obtained




Figure 5. Fresh and dry mass of shoot of basil plants, (a) at vegetative stage, and (b) at flowering stage.




Figure 6. Fresh and dry mass of root of basil plants, (a) at vegetative stage and (b) at flowering stage.

data of dry mass of root due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was non-significant.

Nutrients uptake. Table 3 shows the nitrogen, phosphorus, potassium, calcium and magnesium uptake of basil plants grown in different soilless systems (Aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results indicate that the uptake of nitrogen, phosphorus, potassium, calcium and magnesium by the basil plants were higher in aeroponic system compared those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the nitrogen uptake of basil plants values were 262.50 ± 6.84 , 145.01 ± 4.91 and 185.58 ± 4.22 mg plant^{-1} and 753.99 ± 5.65 , 409.10 ± 5.28 and 387.50 ± 5.29 mg plant^{-1} after 4 and 7 weeks from transplanting, respectively, for aeroponic, hydroponic and peatmoss slabs.

The results indicate that the phosphorus uptake by basil plants values were 74.34 ± 2.90 , 48.34 ± 2.05 and 46.40 ± 3.28 mg plant^{-1} and 224.88 ± 3.05 , 131.86 ± 2.77 and 128.13 ± 2.85 mg plant^{-1} after 4 and 7 weeks from transplanting, respectively, for aeroponic, hydroponic and peatmoss slabs. The potassium uptake by basil plants values were 195.13 ± 4.09 , 136.10 ± 5.51 and 135.06 ± 2.97 mg plant^{-1} and 449.75 ± 4.59 , 375.91 ± 4.34 and 371.00 ± 3.97 mg plant^{-1} after 4 and 7 weeks from transplanting, respectively, for aeroponic, hydroponic and

Soilless Systems	Nutrients uptake, mg plant ⁻¹				
	Nitrogen	Phosphorus	Potassium	Calcium	Magnesium
Vegetative stage (4 weeks after transplanting)					
Aeroponic	262.50 ± 6.84c	74.43 ± 2.90 ^b	195.13 ± 4.09 ^b	132.41 ± 1.54 ^c	41.81 ± 0.83 ^c
Hydroponic	185.58 ± 4.91 ^b	48.34 ± 2.05 ^a	136.10 ± 5.51 ^a	92.86 ± 0.87 ^b	30.53 ± 0.90 ^b
Peatmoss Slabs	145.01 ± 4.22 ^a	46.40 ± 3.28 ^a	135.06 ± 2.97 ^a	83.51 ± 1.32 ^a	24.74 ± 0.58 ^a
Flowering stage (7 weeks after transplanting)					
Aeroponic	753.99 ± 5.65f	224.90 ± 3.05 ^d	449.75 ± 4.59d	529.12 ± 6.63f	112.43 ± 1.67f
Hydroponic	409.10 ± 5.28 ^e	131.86 ± 2.77 ^c	375.91 ± 4.34 ^c	371.91 ± 3.97 ^e	84.53 ± 1.08 ^e
Peatmoss Slabs	387.50 ± 5.29 ^d	128.13 ± 2.85 ^c	371.00 ± 3.97 ^c	262.50 ± 3.20 ^d	71.88 ± 1.10 ^d

Table 3. The nutrients uptake of basil plants grown in different soilless systems. Means on the same column with different superscripts are significantly different ($p < 0.05$).

peatmoss slabs. The calcium uptake by basil plants values were 132.41 ± 1.54 , 92.86 ± 0.84 and 83.51 ± 1.32 mg plant⁻¹ and 529.12 ± 6.63 , 371.91 ± 3.97 and 262.50 ± 3.20 mg plant⁻¹ after 4 and 7 weeks from transplanting, respectively, for aeroponic, hydroponic and peatmoss slabs. The magnesium uptake by basil plants values were 41.81 ± 0.83 , 30.53 ± 0.90 and 24.74 ± 0.58 mg plant⁻¹ and 112.44 ± 1.67 , 84.53 ± 1.08 and 71.88 ± 1.10 mg plant⁻¹ after 4 and 7 weeks from transplanting, respectively, for aeroponic, hydroponic and peatmoss slabs.

The highest values of the N, P, K, Ca and Mg uptakes were 262.50 ± 6.84 , 74.34 ± 2.90 , 195.13 ± 4.09 , 132.41 ± 1.54 and 41.81 ± 0.83 mg plant⁻¹ and 753.99 ± 5.65 , 224.88 ± 3.05 , 449.75 ± 4.59 , 529.12 ± 6.63 and 112.44 ± 1.67 mg plant⁻¹ after 4 and 7 weeks from transplanting, respectively, were found with aeroponic system. While, the lowest values of the N, P, K, Ca and Mg uptakes were 185.58 ± 4.22 , 46.40 ± 3.28 , 135.06 ± 2.97 , 83.51 ± 1.32 and 24.74 ± 0.58 mg plant⁻¹ and 387.50 ± 5.29 , 128.125 ± 2.85 , 371.00 ± 3.97 , 262.50 ± 3.20 and 71.88 ± 1.10 mg plant⁻¹ after 4 and 7 weeks from transplanting, respectively, were found with peatmoss slabs. These results agreed with those obtained by³⁷ they reported that the nutrients uptake of both aeroponic and hydroponic were higher than that in substrate cultivated.

The statistical analysis showed that the differences between the obtained data of nutrients uptake due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant as shown in Table 3.

Content of oil. Figure 7 shows the basil oil content in different soilless systems (aeroponic, hydroponic and peatmoss slabs) at the vegetative stage (4 weeks after transplanting) compared to the flowering stage (7 weeks after transplanting). The results indicate that the basil oil content higher in aeroponic system compared to those of hydroponic system and peatmoss slabs at the vegetative and flowering stages. It could be seen that the basil oil content values were 2.520 (1.80%), 1.722 (1.24%) and 1.129 (1.11%) g plant⁻¹ for aeroponic, hydroponic and peatmoss slabs, respectively, after 4 weeks from transplanting. Meanwhile, the results also indicate that the basil oil content were 6.318 (1.44%), 4.359 (1.90%) and 2.664 (1.42%) g plant⁻¹ for aeroponic, hydroponic and peatmoss slabs, after 7 weeks from transplanting at the same previous. These results are in agreement with findings which were reported by³⁹. The statistical analysis showed that the differences between the obtained data of basil oil content due to the effect of culture system (A) and plant age (B) were significant. The analysis showed also that the interaction between both AB was significant.

Production costs. Table 4 shows the total production costs of basil plants grown in different soilless systems (aeroponic, hydroponic and peatmoss slabs) at the end growing period. It could be seen that the results indicate that the production costs of basil plant were 2.93, 5.27 and 6.24 EGP kg⁻¹ of plant. The total production costs of basil plants grown in hydroponic system were 1.8 times higher than those basil plants grown in aeroponic system, also the total production costs of basil plants grown in peatmoss slabs were 2.1 times higher than those basil plants grown in aeroponic system. Besides it is considered as an organic product which is safe for the human health.




Figure 7. The basil oil content grown in different soilless systems.

Cost item	Units	Production system		
		Aeroponics	Hydroponics	Peatmoss slab
Total direct cost	EGP kg ⁻¹	2.57	4.7	5.57
Total indirect cost	EGP kg ⁻¹	0.36	0.57	0.67
Total cost	EGP kg ⁻¹	2.93	5.27	6.24

Table 4. The total production costs of basil plants grown in different soilless systems.

Model results and validation: The model was validated using the experimental data. Figures 8 and 9 show the predicted and the measured nitrogen, phosphorus, potassium, calcium and magnesium consumption of basil plants during the whole growth period. It could be seen that the N, P, K, Ca and Mg consumption by basil plants increased gradually until it reached the peak after 6 week and then decreased. The results indicate also that, the average daily N, P, K, Ca and Mg consumption by the model was in a reasonable agreement with those measured, where, the nitrogen ranged 2.657 to 13.763 mg plant⁻¹ day⁻¹ theoretically while it was from 2.024 to 13.459 mg plant⁻¹ day⁻¹ experimentally during the whole period. The phosphorus ranged 0.417 to 3.593 mg plant⁻¹ day⁻¹ theoretically while it was from 0.292 to 3.739 mg plant⁻¹ day⁻¹ experimentally during the whole period. The potassium ranged 8.635 to 29.511 mg plant⁻¹ day⁻¹ theoretically while it was from 5.963 to 28.318 mg plant⁻¹ day⁻¹ experimentally during the whole period. The calcium ranged 3.076 to 14.442 mg plant⁻¹ day⁻¹ theoretically while it was from 3.495 to 13.853 mg plant⁻¹ day⁻¹ experimentally during the whole period. The magnesium ranged 0.471 to 1.376 mg plant⁻¹ day⁻¹ theoretically while it was from 0.427 to 1.344 mg plant⁻¹ day⁻¹ experimentally during the whole period. RMSE of N, P, K, Ca and Mg consumption were 0.73, 0.21, 1.5, 0.21 and 0.11, respectively, which means the predicted values were close to the measured values.

The best fit for the relationship between the predicted and the measured values of nutrients consumption was in the following form:

$$NC_p = aNC_M + b \quad (17)$$

where NC_p is the predicted nutrients consumption, mg plant⁻¹ day⁻¹. NC_M is the measured nutrients consumption, mg plant⁻¹ day⁻¹.

The constants of these equation and coefficient of determination are listed in Table 5.

Conclusions

An experiment study was conducted to investigate the possibility of growing basil under three soilless systems (aeroponic, hydroponic and peatmoss slabs). The vegetative parameters, nutrient uptake and oil content were studied. A mathematical model for mass balance of the system was developed successively for predicted the nutrients consumption by basil plant. It is concluded that the aeroponic system recorded the highest values of vegetation parameters (roots, shoots and leaves) and essential oil content. Meanwhile, it consumed the highest values of nutrients (N, P, K, Ca and Mg) and recorded the lowest costs (2.93 EGP kg⁻¹ of plant). The model results were in a reasonable agreement with the experimental ones.




Figure 8. The predicted and the measured nutrients consumption by basil plants during the whole growth period. (a) N, (b) P, (c) K, (d) Ca, (e) Mg.




Figure 9. The comparison between the predicted and the measured nutrients consumption by basil plants during the whole growth period. (a) N, (b) P, (c) K, (d) Ca, (e) Mg.

Items	a	b	R ²
N	0.962	0.717	0.97
P	0.917	0.078	0.98
K	0.903	2.962	0.99
Ca	1.102	-0.984	0.97
Mg	1.065	0.008	0.93

Table 5. The constants a, b and coefficient of determination for nutrients consumption.

Received: 14 December 2020; Accepted: 1 June 2021

Published online: 17 June 2021

References

- Bourgau, F., Gravot, A., Milesi, S. & Gontier, E. Production of plant secondary metabolites: a historical perspective. *Plant Sci.* **161**, 839–851 (2001).
- Pardossi, A., Malorgio, F., Incrocci, L. & Tognoni, F. Hydroponic technologies for greenhouse crops. In *Crops: Quality, Growth and Biotechnology* Vol. 23 (ed. Dris, R.) 360–378 (WFL Publisher, 2006).
- Prasad, A. *et al.* Growth and centelloside production in hydroponically established medicinal plant-*Centella asiatica* (L.). *Ind. Crop Prod.* **35**, 309–312 (2012).
- De Masi, L. *et al.* Assessment of agronomic, chemical and genetic variability in common basil (*Ocimum basilicum* L.). *Eur. Food Res. Technol.* **223**, 273–281 (2006).
- Makri, O. & Kintzios, S. *Ocimum* sp. (Basil): botany, cultivation, pharmaceutical properties, and biotechnology. *J. Herbs. Spices Med. Plants* **13**, 123–150 (2007).
- Saha, S., Monroe, A. & Day, M. R. Growth, yield, plant quality and nutrition of basil (*Ocimum basilicum* L.) under soilless agricultural systems. *Ann. Agric. Sci.* **61**, 181–186 (2016).
- Winsor, G.W. & Schwarz, M. Soilless culture for horticulture crop production. FAO plant production and protection paper, publication no. 101, Rome, Italy (1990).
- Medina, M., Jayachandran, K., Bhat, M. G. & Deoraj, A. Assessing plant growth, water quality and economic effects from application of a plant-based aquafeed in a recirculating aquaponic system. *Aquac. Int.* **24**, 415–427 (2016).
- Johanson, E. K. Aquaponics and hydroponics on a budget. *Tech. Dir.* **69**, 21–23 (2009).
- Rana, S. *et al.* Reclamation of municipal domestic wastewater by aquaponics of tomato plants. *Ecol. Eng.* **37**, 981–988 (2011).
- Hosseinzadeh, S. *et al.* Recirculating water treatment in closed hydroponic systems: assessment of granular activated carbon and soft templated mesoporous carbon for adsorptive removal of root exudates. *Environ. Processes* **6**, 1–23 (2019).
- Qi, L. J., Development of hydro-atomized spraying system for a controlled environment aeroponic farm. Fac. Electrical Engineering, Universiti Teknikal Malaysia Melaka (2012).
- Chalchat, J. C. & Ozcan, M. M. Comparative essential oil composition of flowers, leaves and stems of basil (*Ocimum basilicum* L.) used as herb. *Food Chem.* **110**, 501–503 (2008).
- Ahmed, E. A., Hassan, E. A., Tobgy, K. M. & Ramadan, E. M. Evaluation of rhizobacteria of some medicinal plants for plant growth promotion and biological control. *Ann. Agric. Sci.* **59**(2), 273–280 (2014).
- Nguyen, P. M., Kwee, E. M. & Niemeyer, E. D. Potassium rate alters the antioxidant capacity and phenolic concentration of basil (*Ocimum basilicum* L.) leaves. *Food Chem.* **123**, 1235–1241 (2010).
- Mangmang, J. S., Rosalind, D. & Rogers, G. Inoculation effect of *Azospirillum brasilense* on basil grown under aquaponics production system. *Org. Agric.* **6**(1), 65–74 (2016).
- ASAE Standards. *ASAE S353 DEC97. Moisture Measurement-Meat and Products*, 45th ed.; Adopted and published by ASAE 1998; ASAE: St. Joseph, MI, USA, p. 552 (1998).
- Khater, E.G. Aquaponics: the integration of fish and vegetable culture in recirculating systems. M.Sc. Thesis, in Agric. Eng., Fac. Agric., Moshtohor, Benha Univ., Egypt (2006).
- Parmar, M. R., Bhalodiya, V. B. & Kapdi, S. S. Temperature effect on drying and phytochemicals of basil leaves. *Int. J. Eng. Sci. Investig.* **7**(1), 34–44 (2018).
- Chapman, H.D. & Partt, F.P. Methods of analysis of soils, plant and water. Cal. Univ. 150–200 (1961).
- Bremner, J.M. & Mulvaney, C.S. Nitrogen-total. In *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties*, 2nd ed.; Agronomy series No. 9 ASA; Page, A.L., Miller, R.H., Keeney, D.R., Eds.; SSSA: Madison, WI, USA, pp. 595–624 (1982).
- Murphy, J. & Riley, J. P. A modified single solution method for determination of phosphate in natural waters. *Anal. Chem. Acta* **27**, 31–36 (1962).
- Kiferle, C. *et al.* Rosmarinic acid content in basil plants grown in vitro and in hydroponics. *Cent. Eur. J. Biol.* **6**, 946–957 (2011).
- Khater, E. G., Bahnasawy, A. H., Shams, A. S., Hassaan, M. S. & Hassan, Y. A. Utilization of effluent fish farms in tomato cultivation. *Ecol. Eng.* **83**, 199–207 (2015).
- Nye, P. H. & Tinker, P. B. The concept of a root demand coefficient. *J. Appl. Ecol.* **6**, 293–300 (1969).
- Akeson, M. & Munns, D. N. Uptake of aluminum into root cytoplasm: predicted rates for important solution complexes. *J. Plant Nutr.* **13**, 467–484 (1990).
- Silberbush, M. & Gbur, E. E. Using williams equation to evaluate nutrient uptake rate by intact plants. *Agron. J.* **86**, 107–110 (1994).
- Ritchie, J.T., Johnson, B.S. Soil and plant factors affecting evapotranspiration. In: Stewart, B.A., Nielsen, D.A. (Eds.), *Irrigation of Agricultural Crops*. Agronomy Monograph No. 30, ASA-CSSA-SSSA, Madison, WI, USA, 364–390 (1990).
- Coelho, M. B., Villalobos, F. J. & Mateos, L. Modeling root growth and the soil-plant-atmosphere continuum of cotton crops. *Agric. Water Manag.* **60**, 99–118 (2003).
- Silberbush, M. & Ben-Asher, J. B. Simulation study of nutrient uptake by plants from soilless cultures as affected by salinity buildup and transpiration. *Plant Soil* **233**, 59–69 (2001).
- Villalobos, F. J., Hall, A. J., Ritchie, J. T. & Orgaz, F. OILCROP-SUN: development, growth, and yield model of the sunflower crop. *Agron. J.* **88**, 403–415 (1996).
- Massa, D. *et al.* Simulation of crop water and mineral relations in greenhouse soilless culture. *Environ. Model. Softw.* **26**, 711–722 (2011).
- Martin-Laurent, F. *et al.* A new approach to enhance growth and nodulation of *Acacia mangium* through aeroponic culture. *Biol. Fertil. Soils* **25**, 7–12 (1997).

34. Lakkireddy, K. K. R., Kasturi, K. & Sambasiva Rao, K. R. S. Role of hydroponics and aeroponics in soilless culture in commercial food production. *s research & reviews. J. Agric. Sci. Technol.* **1**(1), 26–35 (2012).
35. Hale, C., Lamotte, F. & Iannetta P. The development of hydroponic and aeroponic culture systems for the high-throughout production of basil (2015). www.hutton.ac.uk/webfm_send/413.
36. Mehandru, P., Shekhawat, N. S., Rai, M. K., Kataria, V. & Gehlot, H. S. Evaluation of aeroponics for clonal propagation of *Caralluma edulis*, *Leptadenia reticulata* and *Tylophora indica*-Three threatened medicinal *Asclepiads*. *Physiol. Mol. Biol. Plants* **20**, 365–373 (2014).
37. Li, Q., Li, X., Tang, B. & Gu, M. Growth responses and root characteristics of lettuce grown in aeroponics, hydroponics, and substrate culture. *Horticulturae* **43**(5), 1–9 (2018).
38. Buckseth, T., Sharma, A. K., Pandey, K. K., Singh, B. P. & Muthuraj, R. Methods of pre-basic seed potato production with special reference to aeroponics—a review. *Sci. Hortic.* **204**, 79–87 (2016).
39. Cheliku, N. *et al.* Essential oil composition of five Basil cultivars (*Ocimum basilicum*) from Albania. *Maced. Pharm. bull.* **61**(2), 11–18 (2015).

Acknowledgements

This work is fully sponsored by the Support and Development of Scientific Research Center, Benha University. We would also like to show our gratitude to the Arab Academy for Science and Technology and Maritime Transport, Cairo, Egypt.

Author contributions

E.-S.K., A.B., W.A., O.M., H.E.-G., Y.S. and M. E.: Investigation, Resources, Writing—Original Draft Preparation, Writing—Review and Editing.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to E.-S.K.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021