



OPEN **Author Correction:** Enhancing
and quantifying spatial
homogeneity in monolayer WS_2

Yameng Cao, Sebastian Wood, Filipe Richheimer, J. Blakesley, Robert J. Young & Fernando A. Castro

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-021-94263-9>, published online 21 July 2021

The original version of this Article contained an error in Figure 5 where panel (e) was incorrectly captured. The original Figure 5 and accompanying legend appear below.

The original Article has been corrected.

Published online: 06 September 2021




Figure 5. Maps and statistics from large area WS_2 flake illuminated with high power ($I_L = 78k Wcm^{-2}$, 10 s) laser in raster-scan. PL maps are acquired before (a) and after (d) the laser illumination, S_{CX} and S_{X0} maps were extracted from modelling the PL maps both before (b,c) and after (e,f) the laser illumination. Histograms of the PL peak intensity (g), charged exciton (h) and exciton (i) spectral weightings before and after the laser illumination are then plotted.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021