OPEN **Publisher Correction:** Association between 9-month isoniazid prophylaxis of latent tuberculosis and severe hepatitis in patients treated with TNF inhibitors

Edward Chia-Cheng Lai, Hsun-Yin Liang, Ya-Chun Huang, Wei-I. Huang, Pi-Hui Chao, Wen-Wen Chen & Meng-Yu Weng

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-021-97444-8>, published online 09 September 2021.

In the original version of this Article, the image on the right in Table 2 and Table 4 did not display correctly. As a result, each bar of the forest plot did not align with the IRR (95% CI) in the adjacent columns.

Additionally, legends for Table 2 and Table 4, erroneously included the sentence: “The figure on right hand side is skew (did not match the IRR)” which has now been removed.

The original Table 2 and Table 4 and their accompanying legends appear below.

The original Article has been corrected.

Published online: 12 October 2021

	No of events (n)	Follow-up duration (years)	Incidence rate ^b (per 100 PYs)	IRR (95% CI) ^c	
Unexposed periods	25	373.23	6.70	1	
Pre-exposure (within 28 day prior to exposure)	1	3.15	3.15	3.81 (0.46–31.59)	
Exposed to INH	5	17.93	27.88	5.1 (1.57–16.55)	
1–28	< 3 ^a	–	32.71	4.00 (0.48–33.30)	
29–56	0	2.30	0	0	
57–84	< 3 ^a	–	91.14	17.29 (3.11–96.25)	
85–168	< 3 ^a	–	38.28	10.55 (1.90–58.51)	
169–252	0	3.65	0	0	
253–280	0	1.50	0	0	
Washout (within 28 days after INH discontinuation)	0	2.84	0	0	

Table 2. Risk of severe hepatitis before and after treatment with isoniazid (n = 31). The figure on right hand side is skew (did not match the IRR) ^aNumbers less than 3 considered as identifiable. ^bIncidence rate was reported as per 100 person-years. ^cIncidence rate ratios (IRR) were adjusted for covariates listed in Table 1.

	No of events (n)	Follow-up duration (years)	Incidence rate ^b (per 100 PYs)	IRR (95% CI) ^c	
Unexposed periods	22	330.65	6.65	1	
Pre-exposure (within 28 day prior to exposure)	0	2.84	0	0	
Exposed to INH	5	15.42	32.42	4.75 (1.43–15.86)	
1–28	< 3 ^a	–	35.71	3.15 (0.37–26.46)	
29–56	0	2.07	0	0	
57–84	< 3 ^a	–	101.81	14.64 (2.49–86.04)	
85–168	< 3 ^a	–	42.47	9.41 (1.68–52.62)	
169–252	0	3.27	0	0	
253–280	0	0.61	0	0	
Washout (within 28 after INH discontinuation)	0	2.53	0	0	

Table 4. Sensitivity analysis of risk of severe hepatitis before and after treatment with isoniazid (excluding death cases) (n = 27). The figure on right side is skew (did not match the results of IRR). ^aNumbers less than 3 considered as identifiable. ^bIncidence rate was reported as per 100 person-years. ^cIncidence rate ratios (IRR) were adjusted for covariates listed in Table 1.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021