



OPEN

# Author Correction: Cochlear SGN neurons elevate pain thresholds in response to music

R. I. M. Dunbar, Eiluned Pearce, Bronwyn Tarr, Adarsh Makdani, Joshua Bamford, Sharon Smith & Francis McGlone

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-021-93969-0>, published online 15 July 2021

The original version of this Article contained an error.

In the ‘General discussion’ section,

“The fact that SGNs are most dense in the lower part of the cochlea may explain why bass instruments are so commonly used to provide the rhythmic accompaniment to music.”

now reads:

“The fact that SGNs are most dense in the apical part of the cochlea may explain why bass instruments are so commonly used to provide the rhythmic accompaniment to music.”

The original Article has been corrected.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022