



OPEN

Author Correction: Fishmeal replacement by periphyton reduces the fish in fish out ratio and alimentation cost in gilthead sea bream *Sparus aurata*

Gilda Savonitto, Roy Barkan, Sheenan Harpaz, Amir Neori, Helena Chernova, Antonio Terlizzi & Lior Guttman

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-021-00466-5>, published online 25 October 2021

The original version of this Article omitted an affiliation for Amir Neori. The correct affiliations are listed below.

Morris Kahn Marine Research Station, Marine Biology Department, The Leon H. Charney School of Marine Sciences, University of Haifa, 3498838 Haifa, Israel

The Interuniversity Institute for Marine Sciences in Eilat, 8810302, Eilat, Israel

The original Article has been corrected.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022