



OPEN

# Association of Toll-like receptors polymorphisms with the risk of acute lymphoblastic leukemia in the Brazilian Amazon

Lilyane Amorim Xabregas<sup>1,2</sup>, Fabíola Silva Alves Hanna<sup>1,2,3</sup>, Fábio Magalhães-Gama<sup>3,4</sup>, Gláucia Lima Souza<sup>1,2</sup>, Daniele Sá Pereira<sup>1,2</sup>, Amanda Barros de Lima<sup>2,3</sup>, Diana Mota Toro<sup>3</sup>, Mirian Rodrigues Ribeiro Santiago<sup>1,2</sup>, Leny Nascimento da Motta Passos<sup>1,2</sup>, Andréa Monteiro Tarragô<sup>1,2,5</sup>, Adriana Malheiro<sup>1,2,3,5</sup> & Allyson Guimarães Costa<sup>1,2,3,4,5,6,7</sup>✉

Acute lymphoblastic leukemia (ALL) is the most common hematologic malignancy in children in childhood. Single-nucleotide polymorphism (SNPs) in key molecules of the immune system, such as Toll-like receptors (TLRs) and CD14 molecules, are associated with the development of several diseases. However, their role in ALL is unknown. A case–control study was performed with 152 ALL patients and 187 healthy individuals to investigate the role of SNPs in TLRs and the CD14 gene in ALL. In this study, *TLR6 C > T rs5743810* [OR: 3.20, 95% CI: 1.11–9.17,  $p = 0.003$ ] and *TLR9 C > T rs187084* (OR: 2.29, 95% CI: 1.23–4.26,  $p = 0.000$ ) seems to be a risk for development of ALL. In addition, the *TLR1 T > G rs5743618* and *TLR6 C > T rs5743810* polymorphisms with protection against death (OR: 0.17, 95% IC: 0.04–0.79,  $p = 0.008$ ; OR: 0.48, 95% IC: 0.24–0.94,  $p = 0.031$ , respectively). Our results show that SNPs in TLRs genes may be involved in the pathogenesis of ALL and may influence clinical prognosis; however, further studies are necessary to elucidate the role of *TLR1*, *TLR4*, *TLR5*, *TLR6*, *TLR9* and *CD14* polymorphisms in this disease.

Leukemia (acute and chronic) represents the 10th most frequent cause of cancer worldwide<sup>1</sup>. In Brazil, for the triennium 2020–2022, approximately 10,810 new cases in women and men are expected and approximately 300 new cases in North region of Brazil, where leukemia (acute and chronic) is the fifth most frequent cancer<sup>2,3</sup>. The most common type of leukemia in childhood is acute lymphoblastic leukemia (ALL) with a prevalence up to 25% of cancers in children who are under the age of 15 years<sup>4</sup>.

The neoplastic process results from genetic errors that contribute to blocking cell maturation and accumulation of leukemic clones (blasts) in the bone marrow microenvironment. Its etiology is still unknown; however, some risk factors are associated, including environmental, genetic and infectious factors<sup>5,6</sup>. Evidence from previous studies suggests that ALL is related to a deficit in immune system regulation in early childhood<sup>7–12</sup>. Furthermore, it is suggested that polymorphisms or genetic variations in the genes of molecules that are important in the development and progression of diseases may be important factors in the increase of intrinsic biological differences, influencing clinically distinct results and conferring genetic susceptibility to cancer<sup>13,14</sup>.

Toll-like receptors make up the main family of pattern recognition receptors (PRRs) of the innate immune system, and are involved in fighting pathogens and inflammation, and recognizing pathogen-associated molecular patterns (PAMPs) and damage-associated molecular patterns (DAMPs), which thus modulates the immune response via the activation of cells that mediate the immune response. In addition, TLRs are vital molecules

<sup>1</sup>Programa de Pós-Graduação em Ciências Aplicadas à Hematologia, Universidade do Estado do Amazonas (UEA), Manaus, AM, Brazil. <sup>2</sup>Diretoria de Ensino e Pesquisa, Fundação Hospitalar de Hematologia e Hemoterapia do Amazonas (HEMOAM), Av. Constantino Nery, 4397, Chapada, Manaus, AM 69050-001, Brazil. <sup>3</sup>Programa de Pós-Graduação em Imunologia Básica e Aplicada, Instituto de Ciências Biológicas, Universidade Federal do Amazonas (UFAM), Manaus, AM, Brazil. <sup>4</sup>Programa de Pós-Graduação em Ciências da Saúde, Instituto René Rachou – Fundação Oswaldo Cruz (FIOCRUZ) Minas, Belo Horizonte, Brazil. <sup>5</sup>Rede Genômica de Vigilância em Saúde do Amazonas (REGESAM), Manaus, AM, Brazil. <sup>6</sup>Programa de Pós-Graduação em Medicina Tropical, Universidade do Estado do Amazonas (UEA), Manaus, AM, Brazil. <sup>7</sup>Escola de Enfermagem de Manaus, Universidade Federal do Amazonas, Manaus, AM, Brazil. ✉email: allyson.gui.costa@gmail.com

Variables	Control group	ALL cases
	(n = 187)	(n = 152)
Age, years (median [IQR])	30 [23–39]	12 [4–18]
<b>Gender</b>		
Male, n (%)	133 (71%)	94 (62%)
Female, n (%)	54 (29%)	58 (38%)
<b>Ethnicity</b>		
Non-hispanic white	39 (21%)	19 (12%)
Multiracial	130 (69%)	130 (86%)
Black	7 (4%)	–
Indigenous	9 (5%)	–
East Asian	2 (1%)	3 (2%)
<b>Comorbidities</b>		
Yes, n (%)	–	67 (44%)
Infectious diseases, n (%)	–	64 (42%)
Others, n (%)	–	3 (2%)
No, n (%)	–	85 (56%)
<b>Immunophenotype</b>		
B-ALL	–	127 (84%)
T-ALL	–	25 (16%)
<b>Residence</b>		
Manaus	187 (100%)	87 (57%)
Interior of Amazonas state	–	53 (35%)
Other state	–	12 (8%)
<b>Relapse</b>		
Yes, n (%)	–	121 (80%)
No, n (%)	–	31 (20%)
<b>Death</b>		
Yes, n (%)	–	70 (46%)
No, n (%)	–	82 (54%)
Hemoglobin, g/dL (median [IQR])	–	8.6 [6.5–10.1]
Hematocrit, % (median [IQR])	–	25.1 [20.2–30.7]
Leukocyte, unit $\times 10^3/\text{mm}^3$ (median [IQR])	–	9435 [3,388–57,850]
Platelets, unit $\times 10^6/\text{mm}^3$ (median [IQR])	–	58,000 [0–152, 800]

**Table 1.** Epidemiological and clinical characteristics of the study population.

in regulating the activation of adaptive immunity and are essential in preventing and curing cancer<sup>15–18</sup>. CD14 normally acts as a receptor for lipopolysaccharide (LPS) and LPS-binding proteins, which are responsible for IL-1, IL-6, and TNF- $\alpha$  inflammatory cytokine production. Studies have reported that CD14 may be involved in development of cancers such as acute lymphoblastic leukemia<sup>19</sup>.

Single-nucleotide polymorphism (SNPs) in TLRs and CD14 genes have been identified in different types of tumors, both in solid tumors and hematologic malignancies<sup>13,20–23</sup>. However, the role of immunogenic genes that play a crucial role in ALL is still poorly understood. In this study, we demonstrated the role of *TLR1*, *TLR4*, *TLR5*, *TLR6*, *TLR9*, and CD14 co-receptor polymorphisms in patients diagnosed with ALL.

## Results

**General characteristics of the study population.** As shown in Table 1, the male gender was predominant in both groups with the median age 30 years for the control group and 12 years for case study group, 133 (71%) and 95 (62%), respectively. Most of the ALL patients had B-ALL (84%) and lived in Manaus (87%). A total of 64 (42%) had infectious comorbidities on diagnosis, 121 (80%) relapsed after induction therapy and 70 (46%) died. These patients also showed a median hemoglobin concentration of 8.6 g/dL, hematocrit of 25.1 g/dL, WBC  $9.435/\text{mm}^3$  and platelets  $58.000/\text{mm}^3$ .

***TLR6 C > T rs5743810* and *TLR9 C > T rs5743836* polymorphisms predict risk of acute lymphoblastic leukemia.** In our analyses, only *TLR5 R > S rs5744105* (OR: 1.39, 95% CI: 1.03–1.89,  $p=0.000$ ) and *TLR6 C > T rs5743810* (OR: 0.57, 95% CI: 0.34–0.95,  $p=0.033$ ) in case group deviated from Hardy–Weinberg equilibrium. In Table 2, it can be observed that the *TLR6 C > T rs5743810* polymorphism seems to be a risk for developing acute lymphoblastic leukemia [OR: 3.20, 95% CI: 1.11–9.17,  $p=0.003$  [Codominant model]], as well as *TLR9 C > T rs187084* (OR: 2.29, 95% CI: 1.23–4.26,  $p=0.000$ , OR<sub>adj</sub>: 2.07, 95% CI<sub>adj</sub>: 0.86–4.98,  $p=0.008$

Genetic models	Control group n = 187 (%)	ALL patients n = 152 (%)	OR <sup>a</sup> (95% CI) <sup>b</sup>	p value <sup>c</sup>	AIC <sup>d</sup>	OR (95% CI) adj <sup>e</sup>	p value adj	AIC
<b>TLR6 C &gt; T rs5743810</b>								
Codominant								
TT	121 (65%)	107 (70%)						
CT	61 (32%)	31 (21%)	0.58 (0.35–0.96)	<b>0.003</b>	459.2	0.61 (0.32–1.17)	0.056	284.8
CC	5 (3%)	14 (9%)	3.20 (1.11–9.17)			3.35 (0.74–15.14)		
Dominant								
TT	121 (65%)	107 (70%)	0.78 (0.49–1.23)	0.284	467.6	0.77 (0.42–1.42)	0.408	287.9
CT-CC	66 (35%)	45 (30%)						
Recessive								
TT-CT	182 (97%)	138 (91%)						
CC	5 (3%)	14 (9%)	3.72 (1.31–10.58)	<b>0.008</b>	461.8	3.90 (0.87–17.48)	0.060	285.0
Overdominant								
TT-CC	126 (67%)	121 (80%)	0.53 (0.32–0.88)	<b>0.012</b>	462.5	0.56 (0.29–1.08)	0.080	285.5
CT	61 (33%)	31 (20%)						
Log-additive 0,1,2	187 (55%)	152 (45%)	1.03 (0.72–1.48)	0.864	468.7	1.00 (0.61–1.64)	0.990	288.5
<b>TLR9 C &gt; T rs187084</b>								
Codominant								
TT	61 (33%)	50 (33%)						
CT	102 (54%)	57 (37%)	0.68 (0.42–1.12)	<b>0.000</b>	455.5	0.58 (0.30–1.14)	<b>0.008</b>	281.1
CC	24 (13%)	45 (30%)	2.29 (1.23–4.26)			2.07 (0.86–4.98)		
Dominant								
TT	61 (33%)	50 (33%)	0.99 (0.63–1.56)	0.957	470.3	0.82 (0.44–1.54)	0.544	288.3
CT-CC	126 (67%)	102 (67%)						
Recessive								
TT-CT	163 (87%)	107 (70%)						
CC	24 (13%)	45 (30%)	2.86 (1.64–4.96)	<b>0.000</b>	455.8	2.85 (1.30–6.24)	<b>0.007</b>	281.6
Overdominant								
TT-CC	85 (45%)	95 (62%)		<b>0.001</b>	460.5		<b>0.008</b>	281.8
CT	102 (55%)	57 (38%)	0.50 (0.32–0.77)			0.45 (0.25–0.82)		
Log-additive 0,1,2	187 (55%)	152 (45%)	1.38 (1.02–1.87)	0.035	465.9	1.26 (0.82–1.91)	0.287	287.6

**Table 2.** Analysis of the association *TLR6 C > T rs5743810* and *TLR9 C > T rs187084* with the risk of acute lymphoblastic leukemia. <sup>a</sup>OR: Odds Ratio; <sup>b</sup>95% confidence interval; <sup>c</sup>p value: <0.05; <sup>d</sup>AIC: Akaike information criterion value; <sup>e</sup>Adj: Adjusted for sex and age.

[Codominant model]) after Bonferroni correction. The other SNPs and allele associations can be found in Supplementary Table 2. We performed a regression model by ancestrally but no difference was observed.

**Association of *TLR1 T > G rs5743618* and *TLR6 C > T rs5743810* polymorphisms with death in acute lymphoblastic leukemia.** In Table 3, the *TLR1 T > G rs5743618* polymorphism was associated with protection from death (OR: 0.48, 95% IC: 0.24–0.94),  $p = 0.031$  [Overdominant model], 95% IC<sub>adj</sub>: 0.21–0.87,  $p = 0.017$  [Overdominant model]), as well as *TLR6 C > T rs5743810* (OR: 0.17, 95% IC: 0.04–0.79,  $p = 0.008$  [Recessive model], OR<sub>adj</sub>: 0.14, 95% IC<sub>adj</sub>: 0.03–0.69,  $p = 0.004$  [Recessive model] after Bonferroni correction. No association was found for infectious comorbidities (Supplementary Table 4). The allelic frequencies from all SNPs are shown in Supplementary Table 5.

## Discussion

Genetic variations in the genes of molecules that are important in the immune response that is involved with the progression of diseases may prove to be important factors in the amplification of intrinsic biological differences, thus influencing clinically distinct results and conferring genetic susceptibility to cancer<sup>13,14</sup>. Single-variant polymorphisms in Toll-like receptors generally altered the TLRs ability to recognize pathogens in two forms: first, by increasing the response against pathogens that contribute to persistent inflammation and cancer and, second, by decreasing their response that contributes to infection susceptibility, as we see in chronic lymphoblastic leukemia (CLL)<sup>13,18,24–26</sup>.

The state of Amazonas, located in the Amazonian region of Brazil, is known for its tropical climate and a highly heterogeneous population that is exposed to several pathogens. In addition, it has different endemic areas for infectious diseases, which could contribute to the modulation of the immune response and consequently

Genetic models	Death		OR <sup>a</sup> (95% CI) <sup>b</sup>	p value <sup>c</sup>	AIC <sup>d</sup>	OR (95% CI) adj <sup>e</sup>	p value adj	AIC
	No n = 82 (%)	Yes n = 70 (%)						
<b>TLR1 T &gt; G rs5743618</b>								
Codominant								
TT	42 (51%)	44 (63%)						
TG	36 (44%)	19 (27%)	0.50 (0.25–1.01)	0.071	210.5	0.45 (0.21–0.93)	<i>0.046</i>	203.4
GG	4 (5%)	7 (10%)	1.67 (0.46–6.12)			1.60 (0.42–6.11)		
Dominant								
TT	42 (51%)	44 (63%)	0.62 (0.32–1.19)	0.148	211.7	0.56 (0.28–1.11)	0.093	204.7
TG-GG	40 (49%)	26 (37%)						
Recessive								
TT-TG	78 (95%)	63 (90%)						
GG	4 (5%)	7 (10%)	2.17 (0.61–7.73)	0.223	212.3	2.16 (0.58–8.00)	0.240	206.1
Overdominant								
TT-GG	46 (56%)	51 (73%)	0.48 (0.24–0.94)	<i>0.031</i>	209.1	0.42 (0.21–0.87)	<b>0.017</b>	201.8
TG	36 (44%)	19 (27%)						
Log-additive 0,1,2	82 (54%)	70 (46%)	0.85 (0.51–1.41)	0.523	213.4	0.80 (0.47–1.36)	0.407	206.8
<b>TLR6 C &gt; T rs5743810</b>								
Codominant								
TT	55 (67%)	52 (74%)						
CT	15 (18%)	16 (23%)	1.13 (0.51–2.51)	<i>0.028</i>	208.7	1.03 (0.44–2.41)	<b>0.017</b>	201.4
CC	12 (15%)	2 (3%)	0.18 (0.04–0.83)			0.14 (0.03–0.70)		
Dominant								
TT	55 (67%)	52 (74%)						
CT-CC	27 (33%)	18 (26%)	0.71 (0.35–1.43)	0.330	212.8	0.61 (0.29–1.29)	0.195	205.8
Recessive								
TT-CT	70 (85%)	68 (97%)						
CC	12 (15%)	2 (3%)	0.17 (0.04–0.79)	<b>0.008</b>	206.8	0.14 (0.03–0.69)	<b>0.004</b>	199.4
Overdominant								
TT-CC	67 (82%)	54 (77%)						
CT	15 (18%)	16 (23%)	1.32 (0.60–2.92)	0.486	213.3	1.23 (0.53–2.83)	0.629	207.3
Log-additive 0,1,2	82 (54%)	70 (46%)	0.62 (0.37–1.05)	0.068	210.5	0.56 (0.32–0.97)	<b>0.032</b>	203.0

**Table 3.** Analysis of the association of single nucleotide polymorphisms (SNPs) in study with the risk of death in acute lymphoblastic leukemia patients. <sup>a</sup>OR: Odds Ratio; <sup>b</sup>95% confidence interval; <sup>c</sup>p value: < 0.05; <sup>d</sup>AIC: Akaike information criterion value; <sup>e</sup>Adj: Adjusted for sex and age. Significant values are in italics.

the triggering of physiological, genetic, and hematological changes<sup>27</sup>. The biological profile of ALL is still little known, and has different patterns according to the geographic and ethnic regions of the world<sup>27–30</sup>.

It is important to note that the population in study is exclusively composed by mixed population. Variations in Brazilian population have been clearly described in literature with a high degree of admixture from Americans, African and/ or European ancestors<sup>31</sup>. Besides, the Amazon Region population has a high degree of inter-ethnic admixture due to the intense miscegenation process and the strong indigenous influence. In this study, it was found that the allelic frequency of *TLR1 T > G rs5743618*, *CD14 C > T rs2569191*, *TLR4 A > G rs4986790*, *TLR4 C > T rs4986791* was similar to the American population, *TLR6 C > T rs5743810* and *TLR9 C > T rs187084* to European population, *TLR9 C > T rs5743836* to African population and *TLR5 R > S rs5744105* to East Asian population<sup>32</sup>. Previous studies demonstrated that children with admixed ancestry have a high risk of developing ALL because of the Native American ancestry<sup>33</sup>, which is highly concentrated in the north of the country, within the Amazon region<sup>34</sup> and might be one of the reasons to explain the number of cases in the region. However, more genetic studies in this population are required to confirm this hypothesis.

To our knowledge, this is the first report to investigate the possible association between polymorphisms in *TLR* and *CD14* genes and the susceptibility to acute lymphoblastic leukemia in the Brazilian Amazon region. In our study, *TLR6 C > T rs5743810* and *TLR9 C > T rs5743836* polymorphisms seem to be a risk for developing acute lymphoblastic leukemia. TLR-6 is a partner of TLR-2, which is responsible for PAMP recognition of invading pathogens and induces inflammation through myeloid differentiation primary response protein 88 (MyD88) and TRAF6 mediated activation of NF-kappa-B (NF-kB). In addition, in most studies, *TLR6 C > T rs5743810* is associated with infectious diseases like tuberculosis in the African population and the induction of resistance to

asthma in children<sup>35</sup>. In CLL, high *TLR6* expression was observed in cells of the patients<sup>36</sup>, but it is still not clear what the role of this polymorphism is in leukemia.

*TLR-9* is found in the endoplasmic reticulum membrane and functions via the MyD88-dependent pathway, which leads to NF- $\kappa$ B activation, cytokine secretion, and inflammatory response. A meta-analysis showed that *TLR9* polymorphisms are associated with increased risk for cancer and some hematological neoplasms, such as non-Hodgkin's lymphoma and acute myeloid leukemia (AML)<sup>37–39</sup>. In AML, the [C] allele of *TLR9 C > T rs187084* and [T] *TLR9 C > T rs5743836* are associated with disease development, as is the TT genotype of the *TLR9 C > T rs5743836* polymorphism with relapse episodes<sup>40</sup>. Additionally, the *TLR9* gene is significantly expressed in chronic lymphocytic leukemia cells<sup>36</sup>.

Since the exposure of hematopoietic stem cells (HSCs) to TLR ligands influences the cycling, differentiation, and their functions, chronic TLR stimulation occurs, which leads to the impairment of normal HSC repopulating activity. Furthermore, high TLR expression and signaling are associated with myelodysplastic syndromes (MDS), which are a group of hematological neoplasms with a high risk of transformation to acute leukemias<sup>13,41</sup>. Therefore, these studies suggested that TLRs seem to be involved in the chronic neoplastic process, but further studies are necessary to confirm this.

A study involving the Turkish population showed no association between *TLR4 A > G rs4986790* and *TLR4 C > T rs4986791* and the risk for ALL<sup>42</sup>. Interestingly, in a cohort of AML patients, both these SNPs were independent risk factors for the development of sepsis and pneumonia<sup>43</sup>. In our study, we did not observe association between SNPs in TLRs with comorbidities infectious and, unfortunately, it was not possible to determine the association between the SNPs with cause of death because of lack information. Polymorphisms in *TLR4* are responsible for decreased recognition of ligands (e.g., LPS). Since homozygotes and heterozygotes are hyporesponsive to LPS stimulation, this polymorphism may be associated with episodes of sepsis<sup>44</sup>. In addition, in B-chronic lymphoblastic leukemia (B-CLL), the reduced *TLR4* expression was associated with increased risk of disease progression and a poor outcome, infectious episodes, and evolution to autoimmune diseases, which suggests an impaired innate immunity<sup>45</sup>. In the literature, this SNP is associated with the risk of cervical<sup>46</sup>, rectal and neck cancer<sup>47</sup>.

It is important to note that our study has some limitations: (i) the sample size is a factor that influences the non-significance of the variants under study, however, studies with a larger sample size are required to increase the statistical power; (ii) the cytokines quantification and gene expression would allow us to better understand the influence of these variants in acute lymphoblastic leukemia patients but we understand that their absence not necessary weak our conclusions; (iii) our control group was composed by blood donors (> 18 years). Although we have included this topic as a limitation of the study, we understand that when using children as a control group, results can be obtained that do not reflect reality, since they could develop the disease after the study occurs; (iv) we did not have access to cause of the death in patients during chemotherapy (disease progression, infection, toxicity and other reason); (v) our clinical data collection was retrospective, so it is possible, for example, that not all patients with infections were identified, since we considered as infected only those who had requests for the described pathogens. Another point is that this was not always requested or the diagnosis was not available. Moreover, the patient could have an infection with some untested pathogen. This leads us to think that the number of patients with infectious comorbidities may be higher.

## Material and methods

**Patients and controls.** A total of 152 pediatric ALL patients diagnosed according to the criteria of the World Health Organization (WHO)<sup>48</sup>, who received treatment at Fundação Hospitalar de Hematologia e Hemoterapia do Amazonas (HEMOAM), 0 to 18 years old, either gender and unrelated were included in the case study group. In the control group, 187 blood donor candidates of either gender and age 18 to 65 years old and who were considered healthy according to the Brazilian Ministry of Health technical standards were included<sup>49</sup>.

**Genomic DNA extraction.** Genomic DNA was extracted from 4 mL of peripheral whole blood at diagnosis using the illustra triplePrep Genomicprep DNA Extraction kits® (GE Healthcare Life Sciences) and BIOPUR Mini Spin plus® extraction kit (Mobius Life Science) for the case group and the QIAmp DNA kit (QIAGEN, Chatsworth, CA, USA) for the control group, following standard laboratory protocols.

**SNP selection and genotyping.** Eight SNPs [*TLR1 I602S (rs5743618)*, *TLR4 A299G (rs4986790)*, *TLR4 T399I (rs4986791)*, *TLR5 R392StopCodon (rs5744105)*, *TLR6 S249P (rs5743810)*, *TLR9-1237C/T (rs5743836)*, *TLR9-1486 C/T (rs187084)* and *CD14-159 (rs2569191)*] were selected according to previously reported hematological disease associations (Non-Hodgkin's lymphoma and AML) and frequency in the Brazilian Amazon population<sup>37,50–52</sup>. Genotyping was performed using the polymerase chain reaction restriction fragment length polymorphism (PCR–RFLP) technique according to the protocol described by COSTA et al. (2017)<sup>52</sup>. The PCR reaction consisted of 2  $\mu$ L of genomic DNA (20 ng) and 23  $\mu$ L of amplification mixture containing 0.1  $\mu$ L of Platinum Taq DNA Polymerase (2 UI), 2.5  $\mu$ L of 10 $\times$  buffer (containing 100 mmol/L Tris–HCl [pH 8.3] and 500 mmol/L KCL), 2  $\mu$ L of MgCl<sub>2</sub> (1.5 mmol/L), 1  $\mu$ L of dNTPs (40 mmol/L), 0.25  $\mu$ L for each forward and reverse primer (0.25 pmol/L) and 16.9  $\mu$ L of ultrapure water. A total of 10  $\mu$ L of PCR product was digested with 5 U of the respective restriction endonuclease (New England Biolabs) and 10 $\times$  enzyme buffer according to the manufacturer's instructions. Primers, PCR cycling conditions, and restriction endonucleases are described in Supplementary Table 1. PCR–RFLP generated fragments were separated using 3–4% agarose gel electrophoresis stained with GelRed™ nucleic acid gel stain (Biotium, Hayward, CA, USA) and visualized on a UV light gel + DocXR system transilluminator (Bio-Rad Corporation, Hercules, CA, USA) with a photographic documentation system.



**Data collection.** The collection of clinical and demographic data was obtained at diagnosis via a search in physical records of the Medical and Statistical Care System (SAME) at the HEMOAM Foundation. In this study, we used the Updated Guidance on the Reporting of Race and Ethnicity in Medical and Science Journals<sup>53</sup>. Infectious comorbidities were considered infections that serologically tested as IgG+ and IgM+ (cytomegalovirus, toxoplasmosis, rubella, varicella and parasitic diseases, among others) according to ALVES et al. (2021)<sup>54</sup>, and pancreatitis, hemorrhage, and sepsis were considered as “Others”. As the relapse criterion, patients who relapsed after induction therapy (35th day of treatment from Grupo Brasileiro de Tratamento das Leucemias Infantis [GBTLI] 2009 protocol)<sup>55</sup> were included and, for death, patients who died within 5 years were included.

**Descriptive and statistical analysis.** Comparison between groups was performed using Fisher’s exact test using GraphPad Prism v.5.0, with a significance level of 5%. Allele analysis was performed using the website <https://ihg.helmholtz-muenchen.de/ihg/snps.html>. A logistic regression analysis that was adjusted for sex and age was performed to find associations between the genotype frequencies and ALL, infectious comorbidities, and death using the package “SNPassoc” version 1.9-2 (<https://cran.r-project.org/web/packages/SNPpassoc/index.html>) for R software version 3.4.3 ([www.r-project.org](http://www.r-project.org)). The best genetic model was performed using the Akaike information criterion (AIC). Hardy–Weinberg equilibrium was evaluated for all SNPs. Bonferroni correction for multiple tests was also performed and the *p*-value adjusted was reported in bold italic characters. The results were shown as the odds ratio (OR) and 95% confidence intervals (95% CI) from multivariate logistic regression analyses.

**Ethics approval and consent to participate.** All protocols and consent forms were approved by the Research Ethics Committee at the HEMOAM Foundation (CEP/HEMOAM process 3.335.123/2019). The informed consent form was obtained from all patients and parents/legal guardians for minors in study. This study was carried out following the guidelines of the Declaration of Helsinki and Resolution 466/12 of the Brazilian National Health Council for research involving human beings.

## Conclusion

To our knowledge, this is the first study to describe the frequency of SNPs in the *TLR1*, *TLR4*, *TLR5*, *TLR6*, *TLR9* and *CD14* genes in patients with ALL in the Brazilian Amazon region. Our study demonstrated that *TLR6 C > T rs5743810* and *TLR9 C > T rs5743836* polymorphisms are associated with the risk of acute lymphoblastic leukemia and *TLR1 T > G rs5743618* and *TLR9 C > T rs5743810* is involved with death. Further studies are necessary to elucidate the role of the *TLR1*, *TLR4*, *TLR5*, *TLR6*, *TLR9*, and *CD14* polymorphisms in the pathogenesis of leukemia.

## Data availability

All data generated or analysed during this study are included in this published article.

Received: 23 February 2022; Accepted: 24 August 2022

Published online: 07 September 2022

## References

1. Ferlay, J. et al. Cancer incidence and mortality worldwide: Sources, methods and major patterns in GLOBOCAN 2012. *Int. J. Cancer* **136**, E359–E386 (2015).
2. Instituto Nacional do Câncer. Estimativa de câncer para o triênio 2020–2022. in *Inca* 122 (2020).
3. Alves, S. et al. Acute lymphoid and myeloid leukemia in a Brazilian Amazon population: Epidemiology and predictors of comorbidity and deaths. *PLoS ONE* **14**, 1–16 (2019).
4. Bethesda. *Childhood Acute Lymphoblastic Leukemia Treatment (PDQ(R)): Health Professional Version*, in *PDQ Cancer Information Summaries*. (2020).
5. Zuckerman, T. & Rowe, J. Pathogenesis and prognostication in acute lymphoblastic leukemia. *F1000Prime Rep.* **6**, 3–7 (2014).
6. Ladines-Castro, W. et al. Morphology of leukaemias. *Rev. Méd. Hosp. Gen. Méx.* **79**, 107–113 (2016).
7. Greaves, M. F. Differentiation-linked leukemogenesis origins of phenotypic diversity in. *Science* **234**, 697–704 (1986).
8. Greaves, M. F. Aetiology of acute leukaemia. *Lancet* **349**, 344–349 (1997).
9. Greaves, M. The ‘delayed infection’ (aka ‘hygiene’) hypothesis for childhood leukaemia. In *The Hygiene Hypothesis and Darwinian Medicine* (ed. Rook, G. A. W.) 239–255 (Birkhäuser Basel, 2009). [https://doi.org/10.1007/978-3-7643-8903-1\\_13](https://doi.org/10.1007/978-3-7643-8903-1_13).
10. Greaves, M. A causal mechanism for childhood acute lymphoblastic leukaemia. *Nat. Rev. Cancer* **18**, 471–484 (2018).
11. Kinlen, L. Childhood leukaemia, nuclear sites, and population mixing. *Br. J. Cancer* **104**, 12–18 (2011).
12. Kinlen, L. J. Childhood leukemia and population mixing. *Cancer Causes Control* **27**, 1499 (2016).
13. Monlish, D. A., Bhatt, S. T. & Schuettelpelz, L. G. The role of Toll-like receptors in hematopoietic malignancies. *Front. Immunol.* **7**, 390 (2016).
14. Saborit-Villarroya, I. et al. E2A is a transcriptional regulator of CD38 expression in chronic lymphocytic leukemia. *Leukemia* **25**, 479–488 (2011).
15. Porcelli, S. A. Effector mechanisms in autoimmunity and inflammation chapter 17 innate immunity. in *Kelley and Firestein’s Textbook of Rheumatology, 2-Volume Set* (Elsevier Inc., 2017). <https://doi.org/10.1016/B978-0-323-31696-5.00017-6>.
16. McDonald, D. R. & Levy, O. 3—Innate immunity. in *Clinical Immunology* (Elsevier Ltd). <https://doi.org/10.1016/B978-0-7020-6896-6.00003-X>.
17. Ishii, K. J. & Akira, S. Cap 3—Innate immunity. in *Clinical Immunology: Principles and Practice* (Elsevier, 2008). <https://doi.org/10.1016/B978-0-323-04404-2.10003-X>.
18. Kaisho, T. & Akira, S. Toll-like receptor function and signaling. *J. Allergy Clin. Immunol.* **117**, 979–987 (2006).
19. Yu, X. et al. Genetic variations in CD14 promoter and acute lymphoblastic leukemia susceptibility in a Chinese population. *DNA Cell Biol.* **30**, 777–782 (2011).
20. Harsini, S., Beigy, M., Akhavan-Sabbagh, M. & Rezaei, N. Toll-like receptors in lymphoid malignancies: Double-edged sword. *Crit. Rev. Oncol. Hematol.* **89**, 262–283 (2014).

21. Messaritakis, I. *et al.* Evaluation of the detection of Toll-like receptors (TLRs) in cancer development and progression in patients with colorectal cancer. *PLoS ONE* **13**, 1–14 (2018).
22. Zhao, S., Zhang, Y., Zhang, Q., Wang, F. & Zhang, D. Toll-like receptors and prostate cancer. *Front. Immunol.* **5**, 1–6 (2014).
23. Rahman, H. A. A., Khorshid, M. M., Khorshid, O. M. R. & Mahgoub, S. M. Toll-like receptor 2 and 9 genetic polymorphisms and the susceptibility to B cell non-Hodgkin lymphoma in Egypt. *Ann. Hematol.* **93**, 1859–1865 (2014).
24. Takeda, K. & Akira, S. Toll-like receptors in innate immunity. *Int. Immunol.* **17**, 1–14 (2005).
25. Wolska, A., Lech-Marañda, E. & Robak, T. Toll-like receptors and their role in carcinogenesis and anti-tumor treatment. *Cell. Mol. Biol. Lett.* **14**, 248–272 (2009).
26. Trejo-De La O, A., Hernández-Sancén, P. & Maldonado-Bernal, C. Relevance of single-nucleotide polymorphisms in human TLR genes to infectious and inflammatory diseases and cancer. *Genes Immun.* **15**, 199–209 (2014).
27. Azevedo-Silva, F., De Camargo, B. & Pombo-de-Oliveira, M. S. Implications of infectious diseases and the adrenal hypothesis for the etiology of childhood acute lymphoblastic leukemia. *Braz. J. Med. Biol. Res.* **43**, 226–229 (2010).
28. Reis, R. D. S. *et al.* Childhood leukemia incidence in Brazil according to different geographical regions. *Pediatr. Blood Cancer* **56**, 58–64 (2011).
29. Chan, L. C. *et al.* Is the timing of exposure to infection a major determinant of acute lymphoblastic leukaemia in Hong Kong?. *Paediatr. Perinat. Epidemiol.* **16**, 154–165 (2002).
30. Pui, C.-H., Robison, L. L. & Look, A. T. Acute lymphoblastic leukaemia. *Lancet* **371**, 1030–1043 (2008).
31. Giolo, S. R. *et al.* Brazilian urban population genetic structure reveals a high degree of admixture. *Eur. J. Hum. Genet.* **20**, 111–116 (2012).
32. 1000 genomes. A deep catalog of human genetic variation. [internationalgenome.org](http://internationalgenome.org) (2020).
33. Carvalho, D. C. *et al.* Amerindian genetic ancestry and INDEL polymorphisms associated with susceptibility of childhood B-cell Leukemia in an admixed population from the Brazilian Amazon. *Leuk. Res.* **39**, 1239–1245 (2015).
34. de Nazaré Cohen-Paes, A. *et al.* Characterization of PCLO gene in Amazonian native American populations. *Genes (Basel)* **13**, 499 (2022).
35. Mukherjee, S., Huda, S. & Sinha Babu, S. P. Toll-like receptor polymorphism in host immune response to infectious diseases. A review. *Scand. J. Immunol.* **90**, 0–2 (2019).
36. Morsi, M. G. *et al.* Quantitative expression of Toll-like receptors TLR-7 and TLR-9 on peripheral blood mononuclear cells in leukemias. *J. Hematol.* **5**, 17–24 (2016).
37. Carvalho, A. *et al.* The rs5743836 polymorphism in TLR9 confers a population-based increased risk of non-Hodgkin lymphoma. *Genes Immun.* **13**, 1–9 (2012).
38. Zhang, L. S., Qin, H. J., Guan, X., Zhang, K. & Liu, Z. R. The TLR9 gene polymorphisms and the risk of cancer: Evidence from a meta-analysis. *PLoS ONE* **8**, e71785 (2013).
39. Nieters, A., Beckmann, L., Deeg, E. & Becker, N. Gene polymorphisms in Toll-like receptors, interleukin-10, and interleukin-10 receptor alpha and lymphoma risk. *Genes Immun.* **7**, 615–624 (2006).
40. Rybka, J. *et al.* Variations in genes involved in regulation of the nuclear factor- $\kappa$ B pathway and the risk of acute myeloid leukaemia. *Int. J. Immunogenet.* **43**, 101–106 (2016).
41. Paracatu, L. C. & Schuettel, L. G. Contribution of aberrant toll like receptor signaling to the pathogenesis of myelodysplastic syndromes. *Front. Immunol.* **11**, 1–10 (2020).
42. Batar, B., Mutlu, T., Özdemir, N., Celkan, T. & Güven, M. TLR4 ve NOD2 polimorfizmlerinin çocukluk çağı akut lenfoblastik lösemi ile ilişkiyi/association of the TLR4 and NOD2 polymorphisms with childhood. *Bezmialem Sci.* **6**, 119–125 (2018).
43. Schnetzke, U. *et al.* Polymorphisms of Toll-like receptors (TLR2 and TLR4) are associated with the risk of infectious complications in acute myeloid leukemia. *Genes Immun.* **16**, 83–88 (2015).
44. Schröder, N. W. J. & Schumann, R. R. Single nucleotide polymorphisms of Toll-like receptors and susceptibility to infectious disease. *Lancet Infect. Dis.* **5**, 156–164 (2005).
45. Barcellini, W. *et al.* Toll-like receptor 4 and 9 expression in B-chronic lymphocytic leukemia: Relationship with infections, autoimmunity and disease progression. *Leuk. Lymphoma* **55**, 1768–1773 (2014).
46. Pandey, N. *et al.* Association of TLR4 and TLR9 gene polymorphisms and haplotypes with cervicitis susceptibility. *PLoS One* **14**, 1–11 (2019).
47. Sheng, W. Y., Yong, Z., Yun, Z., Hong, H. & Hai, L. L. Systematic review/meta-analysis Toll-like receptor 4 gene polymorphisms and susceptibility to colorectal cancer: A meta-analysis and review. *Arch. Med. Sci.* **2015**(11), 699–707 (2015).
48. Arber, D. A. *et al.* The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood* **127**, 2391–2406 (2016).
49. da Saúde, M. *Investigação da transmissão de doenças pelo sangue. Investigação da transmissão de doenças pelo sangue* vol. 1 (2004).
50. Aref, S., Abd Elmaksoud, A. S., Elaziz, S. A., Mabed, M. & Ayed, M. Clinical implication of Toll-like receptors (TLR2 and TLR4) in acute myeloid leukemia patients. *Asian Pac. J. Cancer Prev.* **21**, 3177–3183 (2020).
51. Paolo, S. & Marta, G. Toll-like receptors signaling: A complex network for NF- $\kappa$ B activation in B-cell lymphoid malignancies. *Semin. Cancer Biol.* <https://doi.org/10.1016/j.semcancer.2016.07.001> (2016).
52. Costa, A. G. *et al.* Association of TLR variants with susceptibility to *Plasmodium vivax* malaria and parasitemia in the Amazon region of Brazil. *PLoS ONE* **12**, 1–14 (2017).
53. Flanagan, A., Frey, T. & Christiansen, S. L. Updated guidance on the reporting of race and ethnicity in medical and science journals. *JAMA J. Am. Med. Assoc.* **326**, 621–627 (2021).
54. Alves, F. S. *et al.* Genetic polymorphisms of inflammasome genes associated with pediatric acute lymphoblastic leukemia and clinical prognosis in the Brazilian Amazon. *Sci. Rep.* **11**, 1–10 (2021).
55. Sociedade Brasileira de Oncologia Pediátrica. *Protocolo brasileiro de tratamento da leucemia linfóide aguda na infância GBTLI LLA-2009.* (2011).

## Acknowledgements

We thank the study participants from the HEMOAM Foundation for their important support in this research. Furthermore, we would like to thank all the authors, researchers at LABGEN/HEMOAM and UFAM for their critical discussions and insightful and encouraging ideas.

## Author contributions

L.A.X., F.S.A. and AGC contributed to the conception of the study, laboratory works, data collection and analysis, and writing of the manuscript. A.G.C. was involved in the conception of the study, data analysis, and revision of the manuscript. F.M.G., G.L.S., D.S.P., and A.B.L. contributed to the study design, data acquisition and data analysis. D.M.T., M.R.R.S. and L.N.M.P. were involved in collecting the samples and carrying out the protocol. A.M.T, A.M. and A.G.C. contributed to the study concept and provided advice on genetic aspects. All authors have read and approved the final manuscript.

## Funding

This work was funded by Fundação de Amparo à Pesquisa do Estado do Amazonas (FAPEAM) (Pró-Estado Program [#002/2008, #007/2018 and #005/2019], and POSGRAD Program [#006/2020 and #008/2021]), Rede de Genômica de Vigilância em Saúde do Estado do Amazonas (REGESAM), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). LAX, GLS, DSP and ABL have fellowships from CAPES and FAPEAM (Masters and SI students). FSAH, FMG and DMT have fellowships from CNPq and CAPES (PhD students). AM is a level 2 research fellow from CNPq. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Competing interests

The authors declare no competing interests.

## Additional information

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1038/s41598-022-19130-7>.

**Correspondence** and requests for materials should be addressed to A.G.C.

**Reprints and permissions information** is available at [www.nature.com/reprints](http://www.nature.com/reprints).

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022