



OPEN

Conservation laws, solitary wave solutions, and lie analysis for the nonlinear chains of atoms

Muhammad Junaid-U-Rehman , Grzegorz Kudra & Jan Awrejcewicz

Nonlinear chains of atoms (NCA) are complex systems with rich dynamics, that influence various scientific disciplines. The lie symmetry approach is considered to analyze the NCA. The Lie symmetry method is a powerful mathematical tool for analyzing and solving differential equations with symmetries, facilitating the reduction of complexity and obtaining solutions. After getting the entire vector field by using the Lie scheme, we find the optimal system of symmetries. We have converted assumed PDE into nonlinear ODE by using the optimal system. The new auxiliary scheme is used to find the Travelling wave solutions, while graphical behaviour visually represents relationships and patterns in data or mathematical models. The multiplier method enables the identification of conservation laws, and fundamental principles in physics that assert certain quantities remain constant over time.

The Lie symmetry analysis approach^{1–12} has many applications in different fields, including physics, engineering, and mathematical modelling. It can be used to study a wide range of nonlinear PDEs, including those that are difficult to solve using other methods. Additionally, this approach provides a powerful tool for developing new theories and models that improve our understanding of complex physical systems. Overall, the Lie symmetry analysis approach is a valuable tool for studying nonlinear PDEs and has many significant applications in different branches of science and engineering.

The Lie symmetry analysis approach^{13–23} is a powerful method used in the study of nonlinear PDEs. It is based on the concept of Lie groups and Lie algebras, which are mathematical structures that describe the symmetries of a system. The Lie symmetry analysis approach involves transforming a given PDE into an equivalent system of ODEs using a Lie group transformation. This transformation is constructed from a set of symmetry generators that preserve the form of the original PDE. Once the PDE is transformed into an equivalent system of ODEs, it is possible to use various analytical and numerical methods to solve the system and obtain the solution to the original PDE. Additionally, the Lie symmetry analysis approach can be used to identify the conservation laws that govern the physical behaviour of the system under study. These conservation laws provide important insights into the underlying physical mechanisms responsible for the observed behaviour of the system.

Nonlinear PDEs^{24–27} play a critical role in mechanical engineering by modelling complex phenomena such as stress and deformation in materials, fluid flow, and heat transfer. Unlike linear PDEs, which can be solved analytically in many cases, nonlinear PDEs require numerical or approximate methods to solve due to their complex nature. The use of nonlinear PDEs is essential in the design and optimization of mechanical systems such as turbines, engines, and aircraft. They also provide a framework for predicting the behaviour of materials under different conditions, such as high temperatures, high pressure, and rapid deformation. By incorporating nonlinear PDEs into mechanical engineering models, engineers can improve the accuracy of their designs and ensure that their systems are safe, reliable, and efficient.

Nonlinear chains of atoms^{28–34} have a wide range of applications in mechanical engineering, particularly in the study of materials science and solid mechanics. These models provide valuable insights into the behaviour of materials at the atomic level, enabling the design of high-performance materials for various applications. Applications of nonlinear chains of atoms include the study of thermal conductivity in materials, investigating the deformation and fracture mechanisms of materials under various loading conditions, and studying the dynamics of crystals, such as the propagation of waves and the formation of defects. Nonlinear chains of atoms models are crucial in understanding the behaviour of materials under extreme conditions and can inform the design of materials for high-performance applications, leading to the development of new materials with enhanced mechanical and thermal properties.

Department of Automation, Biomechanics, and Mechatronics, Lodz University of Technology, 1/15 Stefanowski St. (Building A22), 90-924 Lodz, Poland. ✉email: muhammad-junaid.u-rehman@dokt.p.lodz.pl

The analysis of nonlinear chains of atoms using Lie symmetry analysis and conservation laws is motivated by the desire to uncover the fundamental principles that govern the behaviour of these complex systems and gain a deeper understanding of their dynamics and properties. While there has been extensive research on nonlinear chains of atoms, there exist research gaps that can be addressed through the application of Lie symmetry analysis and conservation laws. These gaps include a need for comprehensive studies on the symmetries and conservation laws specific to nonlinear atomic chains, exploration of multiscale behaviour, and bridging the gap between theoretical analysis and practical applications. By addressing these research gaps, researchers can contribute to a more comprehensive understanding of nonlinear chains of atoms and pave the way for advancements in fields such as materials design, nanotechnology, and engineering applications.

The new auxiliary method^{35,36} is a recently proposed method for solving challenging nonlinear PDEs. This method involves introducing an auxiliary variable and constructing a system of coupled equations involving both the original variables and the auxiliary variable. The resulting system of equations can be solved using numerical methods to obtain the solution to the original PDE. The new auxiliary method can handle highly nonlinear PDEs that are difficult to solve using other numerical methods, such as the finite difference approach or the finite element scheme. Additionally, this method can be used to obtain exact solutions to certain types of nonlinear PDEs, reducing the computational cost required to solve some types of nonlinear PDEs. Overall, the new auxiliary method is a promising tool for solving challenging nonlinear PDEs in various fields, including physics, engineering, and mathematical modelling.

Conservation laws of nonlinear PDEs^{37–40} are essential concepts that relate to the principle of conservation of physical quantities like mass, energy, and momentum. These laws are expressed in terms of PDEs and have crucial importance in various fields, including engineering, physics, and mathematical modelling. They provide a mathematical framework to predict the behaviour of complex physical systems accurately and develop new theories and models to improve our understanding of the underlying physical mechanisms. Furthermore, conservation laws play a vital role in the design and analysis of physical systems and the development of numerical methods for solving challenging nonlinear PDEs, making them fundamental concepts in the study of nonlinear PDEs.

The layout of this research is presented as. Formation of supposed model is described in section 2. NAM and multiplier method are explained in section 3. technique and multiplier scheme are presented in section 2. Lie group analysis method is applied on supposed model and entire vector field is described in section 4. Optimal system, similarity reduction, wave solutions, and graphics are represented in section 5. Discussion of graphs and conservation laws of assumed model are described in section 6 and 7 respectively. The conclusions are stated in Sect. 8.

Formation of model

The Hamiltonian of the system is Foroutan et al.⁴¹:

$$H = \sum_n \left\{ \sum_{l \neq n} \mathcal{V}(|\mathcal{U}_n - \mathcal{U}_l|) + \frac{1}{2} m \dot{\mathcal{U}}_n^2 \right\}, \quad (1)$$

where m is the mass of the atom, $\mathcal{V}(|\mathcal{U}_n - \mathcal{U}_l|)$ stands for nonlinear potential and dot indicates for derivative w.r.t time. We consider $l = 1 \pm n$ and the subsequent potential:

$$\mathcal{V}(h_{nl}) = \frac{1}{4} \beta_i h_{nl}^4 + \frac{1}{3} \alpha_i h_{nl}^3 + \frac{1}{2} \gamma_i h_{nl}^2, \quad (2)$$

where h_{nl} is relative displacement among l – th atom and n – th . The index i shows the distinct interactions via the particles. We omit our focus on the first and second neighbours. From Eqs. (1) and (2) through Hamiltonian equations which are

$$\frac{\partial H}{\partial \mathcal{U}_n} = -P_n, \quad \frac{\partial H}{\partial P_n} = \frac{\partial \mathcal{U}_n}{\partial \tau} = \dot{\mathcal{U}}_n,$$

which gives us the equation of motion;

$$\begin{aligned} \frac{d^2 \mathcal{U}_n}{d\tau^2} = & \gamma_1 (\mathcal{U}_{n+1} - 2\mathcal{U}_n + \mathcal{U}_{n-1}) \\ & + \gamma_2 (\mathcal{U}_{n+2} - 2\mathcal{U}_n + \mathcal{U}_{n-2}) \\ & + \alpha_1 \left\{ (\mathcal{U}_{n+1} - \mathcal{U}_n)^2 - (\mathcal{U}_n - \mathcal{U}_{n-1})^2 \right\} \\ & + \alpha_2 \left\{ (\mathcal{U}_{n+2} - \mathcal{U}_n)^2 - (\mathcal{U}_n - \mathcal{U}_{n-2})^2 \right\} \\ & + \beta_1 \left\{ (\mathcal{U}_{n+1} - \mathcal{U}_n)^2 - (\mathcal{U}_n - \mathcal{U}_{n-1})^2 \right\} \\ & + \beta_2 \left\{ (\mathcal{U}_{n+2} - \mathcal{U}_n)^2 - (\mathcal{U}_n - \mathcal{U}_{n-2})^2 \right\}. \end{aligned} \quad (3)$$

In Hamiltonian's equations, P_n stands for generalized momentum. Assuming that the δ (inter-atom spacing) is small enough so that the continuum limit is reached, we substitute $\delta_n \rightarrow \chi$. Then

$$U_{n\pm 1} = U \pm \delta U_x + \frac{1}{2} \delta^2 U_{xx} \pm \frac{1}{6} \delta^3 U_{xxx} + \frac{1}{24} \delta^4 U_{xxxx} + \dots, \tag{4}$$

and

$$U_{n\pm 2} = U \pm 2\delta U_x + \frac{4}{2} \delta^2 U_{xx} \pm \frac{8}{6} \delta^3 U_{xxx} + \frac{16}{24} \delta^4 U_{xxxx} + \dots, \tag{5}$$

hence, Eq. (3) can be supposed as Foroutan et al.⁴²

$$\frac{\partial^2 U}{\partial \tau^2} = \delta_o^2 \frac{\partial^2 U}{\partial \chi^2} + p_o \frac{\partial U}{\partial \chi} \frac{\partial^2 U}{\partial \chi^2} + q_o \left(\frac{\partial U}{\partial \chi} \right)^2 \frac{\partial^2 U}{\partial \chi^2} + r \frac{\partial^4 U}{\partial \chi^4}, \tag{6}$$

with the subsequent constants;

$$\delta_o^2 = \frac{\delta^2}{m} (\gamma_1 + 4\gamma_2), p_o = \frac{2\delta^3}{m} (\gamma_1 + 8\alpha_2), q_o = \frac{3\delta^4}{m} (\beta_1 + 16\beta_2), r = \frac{\delta^4}{12m} (\gamma_1 + 16\gamma_2). \tag{7}$$

here in this paper, we will find out wave solutions and conservation laws for nonlinear Eq. (6) with the use of an appropriate transformation method.

Preliminaries

New auxiliary approach. Assuming the general form of partial PDE is of the form:

$$\mathcal{F}(U, U_\tau, U_\chi, U_{\chi\chi}, \dots) = 0, \tag{8}$$

where τ is the time part and χ is the spatial part and $U = U(\chi, \tau)$ is the dependent variable. We will follow the following steps.

Step 1 Suppose the new similarity variables or transformation is of the form

$$U(\chi, \tau) = \mathcal{H}(\varrho), \quad \text{where } \varrho = k(\chi + c\tau), \tag{9}$$

where k and c both are actual parameters for Eq. (8). Putting the Eq. (9) into Eq. (8) and we get the new ODE below.

$$\mathcal{P}(\mathcal{H}, \mathcal{H}', \mathcal{H}'', \dots) = 0. \tag{10}$$

Step 2 Assume the general solution for Eq. (10) is of the form

$$\mathcal{H}(\varrho) = \sum_{i=0}^N C_i \mathfrak{F}^{iq(\varrho)}, \tag{11}$$

in the above solution, the C_i 's are constants and we will find later and the 1st ODE satisfied $q(\varrho)$.

$$q'(\varrho) = \frac{1}{\ln(\mathfrak{F})} \{ \mathfrak{B}_2 \mathfrak{F}^{-q(\varrho)} + \mathfrak{B}_1 + \mathfrak{B}_3 \mathfrak{F}^{q(\varrho)} \}, \quad \mathfrak{F} > 0, \quad \mathfrak{F} \neq 1. \tag{12}$$

Step 3 In this step, we will use the balancing scheme to execute the value of N . For this, we have to compare the highest-order linear and nonlinear terms to find the value of N .

Step 4 Getting the coefficients of the powers of $\mathfrak{F}^{q(\varrho)}$ ($i = 0, 1, 2, 3, \dots$) by Eqs. (8), (11), and (12). Then collecting the terms of the same power and put them equal to zero which gives us a system of algebraic equations. After solving this system of equations by *Maple*.

Step 5 Finally we will get the different family of solutions for Eq. (12) of the form:

Case 1 When $\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3 < 0$ and $\mathfrak{B}_3 \neq 0$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right), \tag{13}$$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right). \tag{14}$$

Case 2 When $\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3 > 0$ and $\mathfrak{B}_3 \neq 0$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right), \tag{15}$$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} - \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right). \quad (16)$$

Case 3 When $\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3 > 0$ and $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{2} \varrho \right), \quad (17)$$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{2} \varrho \right). \quad (18)$$

Case 4 When $\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3 < 0$, $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{2} \varrho \right), \quad (19)$$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2^2)}}{2} \varrho \right). \quad (20)$$

Case 5 When $\mathfrak{B}_1^2 - \mathfrak{B}_2^2 < 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right), \quad (21)$$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right). \quad (22)$$

Case 6 When $\mathfrak{B}_1^2 - \mathfrak{B}_2^2 > 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right), \quad (23)$$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right). \quad (24)$$

Case 7 When $\mathfrak{B}_2\mathfrak{B}_3 > 0$, $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_1 = 0$

$$\mathfrak{F}^{q(\varrho)} = \sqrt{\frac{-\mathfrak{B}_2}{\mathfrak{B}_3}} \tanh \left(\frac{\sqrt{-\mathfrak{B}_2\mathfrak{B}_3}}{2} \varrho \right), \quad (25)$$

$$\mathfrak{F}^{q(\varrho)} = \sqrt{\frac{-\mathfrak{B}_2}{\mathfrak{B}_3}} \coth \left(\frac{\sqrt{-\mathfrak{B}_2\mathfrak{B}_3}}{2} \varrho \right). \quad (26)$$

Case 8 When $\mathfrak{B}_1 = 0$ and $\mathfrak{B}_2 = -\mathfrak{B}_3$

$$\mathfrak{F}^{q(\varrho)} = \frac{-(1 + e^{2\mathfrak{B}_2\varrho}) \pm \sqrt{2(1 + e^{2\mathfrak{B}_2\varrho})}}{e^{2\mathfrak{B}_2\varrho} - 1}. \quad (27)$$

Case 9 When $\mathfrak{B}_1^2 = \mathfrak{B}_2\mathfrak{B}_3$

$$\mathfrak{F}^{q(\varrho)} = \frac{-\mathfrak{B}_2(\mathfrak{B}_1\varrho + 2)}{\mathfrak{B}_1^2\varrho}. \quad (28)$$

Case 10 When $\mathfrak{B}_1 = k$, $\mathfrak{B}_2 = 2k$ and $\mathfrak{B}_3 = 0$

$$\mathfrak{F}^{q(\varrho)} = e^\varrho - 1. \quad (29)$$

Case 11 When $\mathfrak{B}_1 = k$, $\mathfrak{B}_3 = 2k$ and $\mathfrak{B}_2 = 0$

$$\mathfrak{F}^{q(\varrho)} = \frac{e^\varrho}{1 - e^\varrho}. \quad (30)$$

Case 12 When $2\mathfrak{B}_1 = \mathfrak{B}_2 + \mathfrak{B}_3$

$$\mathfrak{F}^{q(\varrho)} = \frac{1 + \mathfrak{B}_2 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}{1 + \mathfrak{B}_3 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}. \quad (31)$$

Case 13 When $-\mathfrak{B}_1 = \mathfrak{B}_2 + \mathfrak{B}_3$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_2 + \mathfrak{B}_2 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}{\mathfrak{B}_3 + \mathfrak{B}_3 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}. \quad (32)$$

Case 14 When $\mathfrak{B}_2 = 0$

$$\mathfrak{F}^{q(\varrho)} = \frac{\mathfrak{B}_1 e^{\mathfrak{B}_1 \varrho}}{1 + \frac{\mathfrak{B}_3}{2} e^{\mathfrak{B}_1 \varrho}}. \quad (33)$$

Case 15 When $\mathfrak{B}_2 = \mathfrak{B}_1 = \mathfrak{B}_3 \neq 0$

$$\mathfrak{F}^{q(\varrho)} = \frac{-(\mathfrak{B}_2 \varrho + 2)}{\mathfrak{B}_2 \varrho}. \quad (34)$$

Case 16 When $\mathfrak{B}_2 = \mathfrak{B}_3$, $\mathfrak{B}_1 = 0$

$$\mathfrak{F}^{q(\varrho)} = \tan\left(\frac{\mathfrak{B}_2 \varrho + c}{2}\right). \quad (35)$$

Case 17 When $\mathfrak{B}_3 = 0$

$$\mathfrak{F}^{q(\varrho)} = e^{\mathfrak{B}_1 \varrho} - \frac{\mathfrak{B}_2}{2\mathfrak{B}_1}. \quad (36)$$

Multiplier approach. Supposing the Eq. (8) and applying the following steps below:

- (1) Defining the total differential as:

$$D_i = \frac{\partial}{\partial \chi^i} + \mathcal{U}_i \frac{\partial}{\partial \mathcal{U}} + \mathcal{U}_{ij} \frac{\partial}{\partial \mathcal{U}_j} + \dots, \quad i = 1, 2, 3 \dots m. \quad (37)$$

- (2) Defining the Euler operator as below:

$$\frac{\delta}{\delta \mathcal{U}} = \frac{\partial}{\partial \mathcal{U}} - D_i \frac{\partial}{\partial \mathcal{U}_i} + D_{ij} \frac{\partial}{\partial \mathcal{U}_{ij}} - D_{ijk} \frac{\partial}{\partial \mathcal{U}_{ijk}} + \dots. \quad (38)$$

- (3) Let us define n-tuple $f = (f^1, f^2, f^3, \dots, f^m)$, $i = 1, 2, \dots, m$,

$$D_i f^i = 0, \quad (39)$$

Equation (39) is said to be the conservation laws and it fulfils all results of Eq. (8).

- (4) The purpose of $\Lambda(\chi, \tau, \mathcal{U})$ of the Eq. (8):

$$D_i f^i = \Lambda(\chi, \tau, \mathcal{U})H, \quad (40)$$

for some function $\mathcal{U}(\mu^1, \mu^2, \dots, \mu^m)$.

- (5) We will obtain the determining equations for $\Lambda(\chi, \tau, \mathcal{U})$ after calculating the derivative of $\Lambda(\chi, \tau, \mathcal{U})$ in Eq. (40):

$$\frac{\delta}{\delta \mathcal{U}} (\Lambda(\chi, \tau, \mathcal{U})H) = 0. \quad (41)$$

Equation (41) depends for some function $\mathcal{U}(\mu^1, \mu^2, \dots, \mu^m)$. Finally, when we calculate the $\Lambda(\chi, \tau, \mathcal{U})$ with use of Eq. (41), the conservation laws can be acquired by Eq. (40).

Lie group analysis of Eq. (6)

Here, we are supposing the Lie approach for assumed Eq. (6). Now, suppose the one-parameter Lie group of infinitesimal transformations on $(\tau, \chi, \mathcal{U})$ given by

$$\begin{aligned} \bar{\tau} &= \tau + \varepsilon \zeta^1(\tau, \chi, \mathcal{U}) + O(\varepsilon^2), \\ \bar{\chi} &= \chi + \varepsilon \zeta^2(\tau, \chi, \mathcal{U}) + O(\varepsilon^2), \\ \bar{\mathcal{U}} &= u + \varepsilon \eta(\tau, \chi, \mathcal{U}) + O(\varepsilon^2), \end{aligned}$$

and $\varepsilon \ll 1$ is a Small parameter. The associated Lie algebra of infinitesimal symmetries is generated by vector fields

$$\mathfrak{X} = \zeta^1(\tau, \chi, \mathcal{U})\partial_\tau + \zeta^2(\tau, \chi, \mathcal{U})\partial_\chi + \eta(\tau, \chi, \mathcal{U})\partial_{\mathcal{U}}. \tag{42}$$

Equation (42) creates a symmetry of Eq. (6), and \mathfrak{X} satisfy the Lie group conditions

$$Pr^{(4)}\mathfrak{X} \left(\frac{\partial^2 \mathcal{U}}{\partial \tau^2} = \delta_0^2 \frac{\partial^2 \mathcal{U}}{\partial \chi^2} + p_0 \frac{\partial \mathcal{U}}{\partial \chi} \frac{\partial^2 \mathcal{U}}{\partial \chi^2} + q_0 \left(\frac{\partial \mathcal{U}}{\partial \chi} \right)^2 \frac{\partial^2 \mathcal{U}}{\partial \chi^2} + r \frac{\partial^4 \mathcal{U}}{\partial \chi^4} \right) |_{Eq.(6)=0} = 0.$$

The fourth prolongation $Pr^{(4)}$ for \mathfrak{X} can be written as:

$$Pr^{(4)} = \mathfrak{X} + \eta^\chi \frac{\partial}{\partial \mathcal{U}_\chi} + \eta^{\tau\tau} \frac{\partial}{\partial \mathcal{U}_{\tau\tau}} + \eta^{\chi\chi} \frac{\partial}{\partial \mathcal{U}_{\chi\chi}} + \eta^{\chi\chi\chi\chi} \frac{\partial}{\partial \mathcal{U}_{\chi\chi\chi\chi}}, \tag{43}$$

furthermore, we have

$$\begin{cases} \eta^\chi = D_\chi(\eta) - \mathcal{U}_\chi D_\chi(\zeta^1) - \mathcal{U}_\tau D_\chi(\zeta^2), \\ \eta^{\chi\chi} = D_\chi(\eta^\chi) - \mathcal{U}_{\chi\chi} D_\chi(\zeta^1) - \mathcal{U}_{\tau\chi} D_\chi(\zeta^2), \\ \eta^{\tau\tau} = D_\tau(\eta) - \mathcal{U}_\chi D_\tau(\zeta^1) - \mathcal{U}_\tau D_\tau(\zeta^2), \\ \eta^{\tau\chi} = D_\tau(\eta^\chi) - \mathcal{U}_{\tau\tau} D_\tau(\zeta^1) - \mathcal{U}_{\chi\tau} D_\tau(\zeta^2), \\ \eta^{\chi\chi\chi\chi} = D_\chi(\eta^{\chi\chi}) - \mathcal{U}_{\chi\chi\chi} D_\chi(\zeta^1) - \mathcal{U}_{\tau\chi\chi} D_\chi(\zeta^2). \end{cases} \tag{44}$$

Let $(x^1, x^2) = (\chi, \tau)$, where D_i can be written as:

$$D_i = \frac{\partial}{\partial \chi^i} + \mathcal{U}_i \frac{\partial}{\partial \mathcal{U}} + \mathcal{U}_{ij} \frac{\partial}{\partial \mathcal{U}_j} + \dots, \quad i = 1, 2.$$

Substituting the values of η^i which gives us the following vectors:

$$\mathfrak{X}_1 = \frac{\partial}{\partial \chi}, \quad \mathfrak{X}_2 = \frac{\partial}{\partial \tau}, \quad \mathfrak{X}_3 = \frac{\partial}{\partial \mathcal{U}}, \quad \mathfrak{X}_4 = \tau \frac{\partial}{\partial \mathcal{U}}. \tag{45}$$

We see that

$$[\mathfrak{X}_i, \mathfrak{X}_j] = 0, \quad \text{where } i, j = 1, 2, 3.$$

Optimal system

In this section, we observe that from the obtained vector field Eq. (45), the $\mathfrak{X} = \{\mathfrak{X}_1, \mathfrak{X}_2\}$ forms an abelian algebra. So we can use the (42) and get:

$$\begin{aligned} \mathfrak{E}_1 &= \langle \mathfrak{X}_1 \rangle, \\ \mathfrak{E}_2 &= \langle \mathfrak{X}_1 + k_1 \mathfrak{X}_2 \rangle. \end{aligned} \tag{46}$$

Similarity reduction of Eq. (6). Here, we will find the similarity variables and analytical results for Eq. (6).

$\mathfrak{E}_1 = \langle \mathfrak{X}_1 \rangle$. Using the vector \mathfrak{E}_1 , we get the new variable

$$u(\tau, \chi) = \mathcal{H}(\varrho), \quad \text{where } \varrho = \chi, \tag{47}$$

putting the (47) into Eq. (6), which gives us

$$\mathcal{U}(\tau, \chi) = m_1 \tau + m_2, \tag{48}$$

where m_1 and m_2 are integration constants.

$\mathfrak{E}_2 = \langle \mathfrak{X}_1 + k_1 \mathfrak{X}_2 \rangle$. Using the vector \mathfrak{E}_2 , we get the new variable

$$u(\tau, \chi) = \mathcal{H}(\varrho), \quad \text{where } \varrho = \chi + k_1 \tau, \tag{49}$$

putting the (49) into Eq. (6), which gives us

$$6(k_1^2 - \delta_0^2)\mathcal{H}' - 3p_0(\mathcal{H}')^2 - 2q_0(\mathcal{H}')^3 - 3r\mathcal{H}''' = 0. \tag{50}$$

Application of new auxiliary method. Here, we aim to construct the wave patterns for Eq. (6) from Eq. (50) with the use of the proposed technique. Using the balancing method and we obtain $N = 1$. Using the value of $N = 1$ in (11) and we have

$$\mathcal{H}(\varrho) = C_0 + C_1 \mathfrak{F}^{q(\varrho)}. \quad (51)$$

We have to put Eq. (51) into Eq. (50) and we get the system of the equation after comparing the coefficients of $\mathfrak{F}^{q(\varrho)}$. With the use of Maple, we solved the obtained system of equations and got the following results.

$$C_0 = V_1, \quad C_1 = -\frac{12r\mathfrak{B}_3}{p_0}, \quad k_1 = \pm \sqrt{r\mathfrak{B}_1^2 - 4r\mathfrak{B}_2\mathfrak{B}_3 + \delta_0^2}. \quad (52)$$

Using Eq. (52) into Eq. (51) gives us the following set of solutions.

$$\mathcal{U}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \mathfrak{F}^{q(\varrho)}, \quad \text{where } \varrho = \chi \pm \sqrt{r\mathfrak{B}_1^2 - 4r\mathfrak{B}_2\mathfrak{B}_3 + \delta_0^2} \tau. \quad (53)$$

Where V_1 is an arbitrary constant.

Case 1 When $\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3 < 0$ and $\mathfrak{B}_3 \neq 0$

$$\mathcal{U}_1(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}, \quad (54)$$

$$\mathcal{U}_2(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}. \quad (55)$$

Case 2 When $\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3 > 0$ and $\mathfrak{B}_3 \neq 0$

$$\mathcal{U}_3(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}, \quad (56)$$

$$\mathcal{U}_4(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} - \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}. \quad (57)$$

Case 3 When $\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3 > 0$ and $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathcal{U}_5(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}, \quad (58)$$

$$\mathcal{U}_6(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}. \quad (59)$$

Case 4 When $\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3 < 0$, $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathcal{U}_7(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}, \quad (60)$$

$$\mathcal{U}_8(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 + \mathfrak{B}_2\mathfrak{B}_3)}}{2} \varrho \right) \right\}. \quad (61)$$

Case 5 When $\mathfrak{B}_1^2 - \mathfrak{B}_2^2 < 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathcal{U}_9(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_0} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tan \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right) \right\}, \quad (62)$$

$$\mathcal{U}_{10}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \cot \left(\frac{\sqrt{-(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right) \right\}. \quad (63)$$

Case 6 When $\mathfrak{B}_1^2 - \mathfrak{B}_2^2 > 0$ and $\mathfrak{B}_3 \neq -\mathfrak{B}_2$

$$\mathcal{U}_{11}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \tanh \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right) \right\}, \quad (64)$$

$$\mathcal{U}_{12}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-\mathfrak{B}_1}{\mathfrak{B}_3} + \frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{\mathfrak{B}_3} \coth \left(\frac{\sqrt{(\mathfrak{B}_1^2 - \mathfrak{B}_2^2)}}{2} \varrho \right) \right\}. \quad (65)$$

Case 7 When $\mathfrak{B}_2\mathfrak{B}_3 > 0$, $\mathfrak{B}_3 \neq 0$ and $\mathfrak{B}_1 = 0$

$$\mathcal{U}_{13}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \sqrt{\frac{-\mathfrak{B}_2}{\mathfrak{B}_3}} \tanh \left(\frac{\sqrt{-\mathfrak{B}_2\mathfrak{B}_3}}{2} \varrho \right) \right\}, \quad (66)$$

$$\mathcal{U}_{14}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \sqrt{\frac{-\mathfrak{B}_2}{\mathfrak{B}_3}} \coth \left(\frac{\sqrt{-\mathfrak{B}_2\mathfrak{B}_3}}{2} \varrho \right) \right\}. \quad (67)$$

Case 8 When $\mathfrak{B}_1 = 0$ and $\mathfrak{B}_2 = -\mathfrak{B}_3$

$$\mathcal{U}_{15}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-(1 + e^{2\mathfrak{B}_2\varrho}) \pm \sqrt{2(1 + e^{2\mathfrak{B}_2\varrho})}}{e^{2\mathfrak{B}_2\varrho} - 1} \right\}. \quad (68)$$

Case 9 When $\mathfrak{B}_1^2 = \mathfrak{B}_2\mathfrak{B}_3$

$$\mathcal{U}_{16}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-\mathfrak{B}_2(\mathfrak{B}_1\varrho + 2)}{\mathfrak{B}_1^2\varrho} \right\}. \quad (69)$$

Case 10 When $\mathfrak{B}_1 = k$, $\mathfrak{B}_2 = 2k$ and $\mathfrak{B}_3 = 0$

$$\mathcal{U}_{17}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ e^\varrho - 1 \right\}. \quad (70)$$

Case 11 When $\mathfrak{B}_1 = k$, $\mathfrak{B}_3 = 2k$ and $\mathfrak{B}_2 = 0$

$$\mathcal{U}_{18}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{e^\varrho}{1 - e^\varrho} \right\}. \quad (71)$$

Case 12 When $2\mathfrak{B}_1 = \mathfrak{B}_2 + \mathfrak{B}_3$

$$\mathcal{U}_{19}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{1 + \mathfrak{B}_2 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}{1 + \mathfrak{B}_3 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}} \right\}. \quad (72)$$

Case 13 When $-2\mathfrak{B}_1 = \mathfrak{B}_2 + \mathfrak{B}_3$

$$\mathcal{U}_{20}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{\mathfrak{B}_2 + \mathfrak{B}_2 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}}{\mathfrak{B}_3 + \mathfrak{B}_3 e^{\frac{1}{2}(\mathfrak{B}_2 - \mathfrak{B}_3)\varrho}} \right\}. \quad (73)$$

Case 14 When $\mathfrak{B}_2 = 0$

$$\mathcal{U}_{21}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{\mathfrak{B}_1 e^{\mathfrak{B}_1\varrho}}{1 + \frac{\mathfrak{B}_3}{2} e^{\mathfrak{B}_1\varrho}} \right\}. \quad (74)$$

Case 15 When $\mathfrak{B}_2 = \mathfrak{B}_1 = \mathfrak{B}_3 \neq 0$

$$\mathcal{U}_{22}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \frac{-(\mathfrak{B}_2\varrho + 2)}{\mathfrak{B}_2\varrho} \right\}. \quad (75)$$

Case 16 When $\mathfrak{B}_2 = \mathfrak{B}_3$, $\mathfrak{B}_1 = 0$

$$\mathcal{U}_{23}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ \tan \left(\frac{\mathfrak{B}_2\varrho + c}{2} \right) \right\}. \quad (76)$$

Case 17 When $\mathfrak{B}_3 = 0$

$$U_{24}(\chi, \tau) = V_1 - \frac{12r\mathfrak{B}_3}{p_o} \left\{ e^{\mathfrak{B}_1 \varrho} - \frac{\mathfrak{B}_2}{2\mathfrak{B}_1} \right\} \tag{77}$$

where $\varrho = \chi \pm \sqrt{r\mathfrak{B}_1^2 - 4r\mathfrak{B}_2\mathfrak{B}_3 + \delta_o^2} \tau$ is given according to.

Graphics and discussion

A graphical representation of obtained solutions is discussed here in this section. By using the new auxiliary method we have constructed the analytical behaviour of the considered model in the form of trigonometric functions, hyperbolic trigonometric functions, and exponential, and algebraic-type results. The graph of the tangent function is periodic with a period of Π and has vertical asymptotes at odd multiples of $\frac{\Pi}{2}$. As χ approaches these vertical asymptotes, the tangent function approaches positive or negative infinity depending on the direction of the approach. The graph of the cotangent function is also periodic with a period of Π and has horizontal asymptotes at even multiples of Π . As χ approaches these horizontal asymptotes, the cotangent function approaches zero. We have plotted the behaviour of some obtained results. Figure 1 shows the graphical behavior of $u_1(\chi, \tau)$ for the choice of parameters $\delta_o = 1, r = 2, p_o = 3, \mathfrak{B}_1 = 2, \mathfrak{B}_2 = 1, \mathfrak{B}_3 = 2, \tau = 1$. Figure 2 represent the behaviour of $u_2(\chi, \tau)$ for the choice of parameters $\delta_o = 2, r = 1, p_o = 1, \mathfrak{B}_1 = 3, \mathfrak{B}_2 = 2, \mathfrak{B}_3 = 1, \tau = 2$. Figure 3 shows the Graphics of $u_3(\chi, \tau)$ for the choice of parameters $\delta_o = 1.5, r = 1, p_o = 4, \mathfrak{B}_1 = 4, \mathfrak{B}_2 = 3, \mathfrak{B}_3 = 5, \tau = 2$. Figure 4 shows the Graphics of $u_4(\chi, \tau)$ for the choice of parameters $\delta_o = 5, r = 0.5, p_o = 3.5, \mathfrak{B}_1 = 1.5, \mathfrak{B}_2 = 3.5, \mathfrak{B}_3 = 2.5, \tau = 3$. Figure 5 represent the behaviour of $u_{19}(\chi, \tau)$ for the choice of parameters $\delta_o = 6, r = 3, p_o = 5, \mathfrak{B}_1 = 1, \mathfrak{B}_2 = 3, \mathfrak{B}_3 = 5, \tau = 5$. The behaviour of U_{19} shows Singularities exist in exponential functions due to their intrinsic nature of rapid and unbounded growth or decay. The exponential function, typically represented as $f(\chi) = e^\chi$, where e is Euler's number approximately equal to 2.71828, captures the exponential growth or decay phenomenon. However, certain values of χ result in problematic behaviour or undefined outcomes. When χ approaches positive or negative infinity, the exponential function exhibits an asymptotic behaviour, approaching infinity or zero, respectively, without reaching a definite value. These instances represent singularities where the exponential function becomes indeterminate or encounters difficulties in providing a well-defined result. These singularities in the exponential function highlight the inherent limitations and special characteristics associated with exponential growth or decay processes.




Figure 1. Graphics of (a) $U_1(\chi, \tau)$ versus time component τ for the choice of parameters $\delta_o = 1, r = 2, p_o = 3, \mathfrak{B}_1 = 2, \mathfrak{B}_2 = 1, \mathfrak{B}_3 = 2$ and (b) considering at $\tau = 1$.




Figure 2. Graphics of (a) $U_2(\chi, \tau)$ versus time component τ for the choice of parameters $\delta_o = 2, r = 1, p_o = 1, \mathfrak{B}_1 = 3, \mathfrak{B}_2 = 2, \mathfrak{B}_3 = 1$, and (b) considering at $\tau = 2$.




Figure 3. Graphics of (a) $\mathcal{U}_3(\chi, \tau)$ versus time component τ for the choice of parameters $\delta_o = 1.5, r = 1, p_o = 4, \mathfrak{B}_1 = 4, \mathfrak{B}_2 = 3, \mathfrak{B}_3 = 5$, and (b) considering at $\tau = 2$.




Figure 4. Graphics of (a) $\mathcal{U}_4(\chi, \tau)$ versus time component τ for the choice of parameters $\delta_o = 5, r = 0.5, p_o = 3.5, \mathfrak{B}_1 = 1.5, \mathfrak{B}_2 = 3.5, \mathfrak{B}_3 = 2.5$, and (b) considering at $\tau = 3$.




Figure 5. Graphics of (a) $\mathcal{U}_{19}(\chi, \tau)$ versus time component τ for the choice of parameters $\delta_o = 6, r = 3, p_o = 5, \mathfrak{B}_1 = 1, \mathfrak{B}_2 = 3, \mathfrak{B}_3 = 5$, and (b) considering at $\tau = 5$.

Conservation laws

In this portion, we will construct the conservation laws by multiplier approach for Eq. (6). We obtain the determinant equation for $\Lambda(X, t, u)$ by Eq. (41).

$$\frac{\delta}{\delta \mathcal{U}} \left[\Lambda \left(\frac{\partial^2 \mathcal{U}}{\partial \tau^2} - \delta_o^2 \frac{\partial^2 \mathcal{U}}{\partial \chi^2} - p_o \frac{\partial \mathcal{U}}{\partial \chi} \frac{\partial^2 \mathcal{U}}{\partial \chi^2} - q_o \left(\frac{\partial \mathcal{U}}{\partial \chi} \right)^2 \frac{\partial^2 \mathcal{U}}{\partial \chi^2} - r \frac{\partial^4 \mathcal{U}}{\partial \chi^4} \right) \right] = 0. \tag{78}$$

Using Eq. (38), we can write the Euler operator is of the form

$$\frac{\delta}{\delta \mathcal{U}} = \frac{\partial}{\partial \mathcal{U}} - D_\tau \frac{\partial}{\partial \mathcal{U}_\tau} - D_\chi \frac{\partial}{\partial \mathcal{U}_\chi} + D_\tau^2 \frac{\partial}{\partial \mathcal{U}_{\tau\tau}} + D_\chi^2 \frac{\partial}{\partial \mathcal{U}_{\chi\chi}} + D_\chi D_\tau \frac{\partial}{\partial \mathcal{U}_{\tau\chi}} - \dots, \tag{79}$$

defining the total derivative operators D_τ and D_χ from Eq. (37).

$$\begin{aligned}
 D_\tau &= \frac{\partial}{\partial \tau} + \mathcal{U}_\tau \frac{\partial}{\partial \mathcal{U}} + \mathcal{U}_{\tau\tau} \frac{\partial}{\partial \mathcal{U}_\tau} + \mathcal{U}_{\tau\chi} \frac{\partial}{\partial \mathcal{U}_\chi} \cdots, \\
 D_\chi &= \frac{\partial}{\partial \chi} + \mathcal{U}_\chi \frac{\partial}{\partial \mathcal{U}} + \mathcal{U}_{\chi\chi} \frac{\partial}{\partial \mathcal{U}_\chi} + \mathcal{U}_{\tau\chi} \frac{\partial}{\partial \mathcal{U}_\tau} \cdots,
 \end{aligned}
 \tag{80}$$

computing Eq. (78) and we obtain the following multipliers and conservation laws:

$$\Lambda = C_1\tau + C_2, \tag{81}$$

by Eqs. (40) and (81), The following conservation laws are found

$$\begin{aligned}
 \mathcal{T}^\tau &= \mathcal{U}_\tau(C_1\tau + C_2) - C_1\mathcal{U}, \\
 \mathcal{T}^\chi &= (C_1\tau + C_2) \left(\frac{-1}{3}\mathcal{U}_\chi^3 q_0 - \frac{1}{2}\mathcal{U}_\chi^2 p_0 - \mathcal{U}_\chi \delta_0^2 - r\mathcal{U}_{\chi\chi\chi} \right).
 \end{aligned}
 \tag{82}$$

Using Eq. (82) and we get the following two cases of conservation laws.

Case 1 For $C_1 = 1$, $C_2 = 0$, then $\Lambda_1 = \tau$, we get the following fluxes:

$$\begin{aligned}
 \mathcal{T}_1^\tau &= \tau\mathcal{U}_\tau - \mathcal{U}, \\
 \mathcal{T}_1^\chi &= \frac{-1}{3}\mathcal{U}_\chi^3 \tau q_0 - \frac{1}{2}\mathcal{U}_\chi^2 \tau p_0 - \mathcal{U}_\chi \tau \delta_0^2 - r\mathcal{U}_{\chi\chi\chi} \tau.
 \end{aligned}
 \tag{83}$$

Case 2 For $C_1 = 0$, $C_2 = 1$, then $\Lambda_1 = 1$, we get the following fluxes:

$$\begin{aligned}
 \mathcal{T}_2^\tau &= \mathcal{U}_\tau, \\
 \mathcal{T}_2^\chi &= \frac{-1}{3}\mathcal{U}_\chi^3 q_0 - \frac{1}{2}\mathcal{U}_\chi^2 p_0 - \mathcal{U}_\chi \delta_0^2 - r\mathcal{U}_{\chi\chi\chi}.
 \end{aligned}
 \tag{84}$$

Conclusion

In this research, nonlinear chains of atoms (NCA) are studied. NCA play a crucial role in materials design and fabrication, enabling the development of advanced materials with tailored properties. The unique structural arrangements and interactions of nonlinear chains of atoms influence the mechanical, electrical, and optical properties of materials, which are essential for designing innovative technologies in fields such as nanotechnology, optoelectronics, and energy systems. Also, NCA are of great interest in the study of dynamical systems and nonlinear phenomena. They provide valuable insights into the behaviour of complex systems, allowing for the analysis and modelling of intricate dynamics observed in areas like physics, biology, and control theory. The Lie symmetry method and NAM enhances existing techniques, offering additional insights, improved accuracy, or simplified computations. Travelling wave solutions describe wave-like behaviour propagating through systems, while graphical behaviour provides visual representations of relationships and patterns in data or mathematical models. The multiplier method allows for the identification of conservation laws, which are fundamental principles in physics that state certain quantities remain constant over time. Understanding conservation laws and utilizing mathematical techniques such as the Lie symmetry method, travelling wave solutions, and graphical analysis contributes to a deeper understanding of nonlinear chains of atoms and their dynamics.

Data availability

All data that support the findings of this study are included within the article.

Received: 5 June 2023; Accepted: 12 July 2023

Published online: 17 July 2023

References

- Riaz, M. B., Atangana, A., Jhangeer, A. & Junaid-U-Rehman, M. Some exact explicit solutions and conservation laws of Chaffee-Infante equation by Lie symmetry analysis. *Phys. Scr.* **96**(8), 084008 (2021).
- Kumar, S., Jadaun, V. & Ma, W. X. Application of the Lie symmetry approach to an extended Jimbo-Miwa equation in (3+ 1) dimensions. *Eur. Phys. J. Plus* **136**(8), 1–30 (2021).
- Jadaun, V. Soliton solutions of a (3+ 1)-dimensional nonlinear evolution equation for modelling the dynamics of ocean waves. *Phys. Scr.* **96**(9), 095204 (2021).
- Jadaun, V. Lie symmetry approach to study the impact of a bulge on hemodynamics in an elastic-compliant artery. *Math. Methods Appl. Sci.* (2023).
- Jhangeer, A. *et al.* Lie analysis, conservation laws, and travelling wave structures of nonlinear Bogoyavlenskii-Kadomtsev-Petviashvili equation. *Results Phys.* **19**, 103492 (2020).
- Noether, E. Invariante variations probleme. *Nachr. König. Ges. Wiss. Gött. Math.-Phys. Kl. Heft.* **2**, 235–257 (1918).
- Kara, A. H. & Mahomed, F. M. Noether-type symmetries and conservation laws via partial Lagrangians. *Nonlinear Dyn.* **45**, 367–383 (2006).
- Freire, I. L. New conservation laws for inviscid Burgers equation. *Comput. Appl. Math.* **31**, 559–567 (2012).
- Torrisi, M. & Tracina, R. Quasi self-adjointness of a class of third-order nonlinear dispersive equations. *Nonlinear Anal. Real World Appl.* **14**, 1496–1502 (2013).
- Gandarias, M. L. & Bruzon, M. S. Some conservation laws for a forced KdV equation. *Nonlinear Anal. Real World Appl.* **13**, 2692–2700 (2012).

11. Riaz, M. B., Baleanu, D., Jhangeer, A. & Abbas, N. Nonlinear self-adjointness, conserved vectors, and travelling wave structures for the kinetics of phase separation dependent on ternary alloys in iron (Fe-Cr-Y (Y= Mo, Cu)). *Results Phys.* **25**, 104151 (2021).
12. Riaz, M. B., Awrejcewicz, J., Jhangeer, A. & Junaid-U-Rehman, M. A variety of new traveling wave packets and conservation laws to the nonlinear low-pass electrical transmission lines via lie analysis. *Fractal Fract.* **5**, 170 (2021).
13. Rogers, C. & Schief, W. K. *Bäcklund and Darboux Transformation Geometry and Modern Applications in Soliton Theory* (Cambridge University Press, 2002).
14. Bluman, G. W., Cheviakov, A. F. & Anco, S. C. *Applications of Symmetry Methods to Partial Differential Equations* (Springer, 2000).
15. Ibragimov, N. H. A new conservation theorem. *J. Math. Anal. Appl.* **333**, 311–28 (2007).
16. Olver, P. J. *Application of Lie Groups to Differential Equations* (Springer-Verlag, 1993).
17. Ovsiannikov, L. V. *Group Analysis of Differential Equations* (Academic, 1982).
18. Hirota, R. *Direct Methods in Soliton Theory* (Springer, 2004).
19. Bluman, G. W. & Kumei, S. *Symmetries and Differential Equations* (Springer-Verlag, 1989).
20. Riaz, M. B., Jhangeer, A., Abualnaja, K. M. & Junaid-U-Rehman, M. Conserved quantities and travelling wave profiles to the nonlinear transmission line via Lie group analysis. *Phys. Scr.* **96**(10), 104013 (2021).
21. Hussain, A., Junaid-U-Rehman, M., Jabeen, F. & Khan, I. Optical solitons of NLS-type differential equations by the extended direct algebraic method. *Int. J. Geom. Methods Modern Phys.* **19**(05), 2250075 (2022).
22. Hussain, A., Anjum, A., Junaid-U-Rehman, M., Khan, I., Sameh, M. A. & Al-Johani, A. S. Symmetries, optimal system, exact and soliton solutions of (3 + 1)-dimensional Gardner-KP equation. *J. Ocean Eng. Sci.* (2022).
23. Riaz, M. B. *et al.* Some new wave profiles and conservation laws in a Pre-compressed one-dimensional granular crystal by Lie group analysis. *Eur. Phys. J. Plus* **137**(3), 401 (2022).
24. Seadawy, A. R. *et al.* Conservation laws, optical molecules, modulation instability, and Painlevé analysis for the Chen-Lee-Liu model. *Opt. Quantum Electron.* **53**(4), 1–15 (2021).
25. Seadawy, A. R., Rizvi, S. T. R., Ahmad, S., Younis, M. & Baleanu, D. Lump, lump-one stripe, multi-wave and breather solutions for the Hunter-Saxton equation. *Open Phys.* **19**(1), 1–10 (2021).
26. Seadawy, A. R., Ahmed, H. M., Rabie, W. B. & Biswas, A. Chirp-free optical solitons in fibre Bragg gratings with dispersive reflectivity having polynomial law of nonlinearity. *Optik* **225**, 165681 (2021).
27. Jhangeer, A., Hussain, A., Junaid-U-Rehman, M., Baleanu, D. & Riaz, M. B. Quasi-periodic, chaotic, and travelling wave structures of modified Gardner equation. *Chaos Solitons Fractals* **143**, 110578 (2021).
28. Peyrard, M. & Remoissenet, M. Solitonlike excitations in a one-dimensional atomic chain with a nonlinear deformable substrate potential. *Phys. Rev. B* **26**(6), 2886 (1982).
29. Fang, X., Wen, J., Yin, J. & Yu, D. Wave propagation in nonlinear metamaterial multi-atomic chains based on homotopy method. *AIP Adv.* **6**(12), 121706 (2016).
30. Braun, O. M. & Kivshar, Y. S. *The Frenkel-Kontorova Model: Concepts, Methods, and Applications* 475 (Springer, 2004).
31. Kaur, L. & Wazwaz, A. M. Einstein's vacuum field equation: Painlevé analysis and Lie symmetries. *Waves Rand. Compl. Media* **31**(2), 199–206 (2021).
32. Khan, M. H. & Wazwaz, A. M. Lump, multi-lump, cross kinky-lump and manifold periodic-soliton solutions for the (2 + 1)-D Calogero-Bogoyavlenskii-Schif equation. *Heliyon* **6**(4), 03701 (2020).
33. Khater, A. H., Callebaut, D. K. & Seadawy, A. R. General soliton solutions for nonlinear dispersive waves in convective type instabilities. *Phys. Sci.* **74**, 384–393 (2006).
34. Kumar, S., Kumar, D. & Kumar, A. Lie symmetry analysis for obtaining the abundant exact solutions, optimal system and dynamics of solitons for a higher-dimensional Fokas equation. *Chaos Solitons Fractals* **142**, 110507 (2021).
35. Jarad, F., Jhangeer, A., Awrejcewicz, J., Riaz, M. B. & Junaid-U-Rehman, M. Investigation of wave solutions and conservation laws of generalized Calogero-Bogoyavlenskii-Schiff equation by group theoretic method. *Results Phys.* **37**, 105479 (2022).
36. Khan, M. A., Akbar, M. A. & binti Abd Hamid, N. N. Travelling wave solutions for space-time fractional cahn Hilliard equation and space-time fractional symmetric regularized long-wave equation. *Alex. Eng. J.* **60**, 1317–24 (2021).
37. Ablowitz, M. J. & Clarkson, P. A. Solitons, nonlinear evolution equations, and inverse scattering, in *Vol. 149 of London Mathematical Society Lecture Note Series* (Cambridge University Press, 1991).
38. Naz, R., Mahomed, F. M. & Mason, D. P. Comparison of different approaches to conservation laws for some partial differential equations in fluid mechanics. *Appl. Math. Comput.* **205**, 212–30 (2008).
39. Bruzón, M. S., Gandarias, M. L. & Ibragimov, N. H. Self-adjoint sub-classes of generalized thin film equations. *J. Math. Anal. Appl.* **357**, 307–13 (2009).
40. Ibragimov, N. H. Nonlinear self-adjointness and conservation laws. *J. Phys. A* **44**, 432002 (2011).
41. Foroutan, M., Zamanpour, I. & Manafan, J. Applications of IBSOM and ETEM for solving the nonlinear chains of atoms with long-range interactions. *Eur. Phys. J. Plus* **132**(10), 1–18 (2017).
42. Ahmed, S., Seadawy, A. R. & Rizvi, S. T. Study of breathers, rogue waves, and lump solutions for the nonlinear chains of atoms. *Opt. Quantum Electron.* **54**(5), 320 (2022).

Acknowledgements

The first author, Muhammad Junaid U Rehman, is a doctoral candidate in the Interdisciplinary Doctoral School at the Lodz University of Technology, Poland. This work has been supported by the Polish National Science Centre, Poland under the Grant OPUS 18 No.2019/35/B/ST8/00980.

Author contributions

M.J.: Conceptualization, Methodology, Investigation, Software, Writing—original draft. G.K.: Supervision, Software, Validation. J.A.: Visualization, Investigation, Reviewing the original draft.

Competing Interest

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.J.-U.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023