



OPEN

Effects of climate warming and human activities on the distribution patterns of *Fritillaria unibracteata* in eastern Qinghai-Tibetan Plateau

Dan Zhao¹, Jun Wang¹✉, Wei Dai¹, Kunhao Ye¹, Jie Chen¹, Qianglong Lai¹, Haiying Li², Binglian Zhong² & Xiaoli Yu²

Fritillaria unibracteata is an endangered medicinal material species endemic to the Qinghai Tibet Plateau, and belongs to the national Class III endangered plant. In addition to expelling wind and removing dampness, it also warms menstruation and relieves pain in clinic use of traditional Chinese medicine. In recent years, affected by the destruction of shrubs and climate change, the habitat of *F. unibracteata* wild resources has been seriously damaged, indicating of great significance to predict its potential suitable habitat using MaxEnt model. The AUC values without human activities were 0.983 ± 0.013 – 0.988 ± 0.001 , while it is 0.982 ± 0.015 – 0.989 ± 0.000 with human activities, justifying their applications for predicting the potential areas of *F. unibracteata*. Without human activities, there were 8.47×10^4 km² of highly suitable habitats in northern Sichuan, southern Gansu and southeastern Qinghai. But the poorly, moderately and highly suitable areas of *F. unibracteata* have decreased to 33.8×10^4 km², 9.66×10^4 km² and 6.64×10^4 km² due to human activities. Environmental variables affecting *F. unibracteata* distribution included the minimum temperature in the coldest month (-16.89 – -4.96 °C), annual precipitation (416.64–866.96 mm), temperature annual range (24.83–31.97 °C), elevation (2879.69–3981.82 m), human footprint (2.58–23.66) and mean UV-B of highest month (7381.92–8574.27 kJ/m²). In the 2050s and 2090s, human activities would significantly reduce the highly suitable habitats of *F. unibracteata*. Under SSP1-2.6, SSP2-4.5 and SSP5-8.5 scenarios, the centroid would move to the low latitude area from the current position first, and then to a high latitude area. Wild resources of *F. unibracteata* in China can be effectively conserved based on our results.

In the past century, due to the impact of changes in natural conditions and human activities, the global climate is developing towards a warming trend¹. In the past century, the global average temperature has increased by 0.74 °C, and extreme weather events have occurred frequently². Climate change will promote the migration of species in suitable areas, and even put some species at risk of extinction^{3,4}. China is extremely vulnerable to climate change with its location in the East Asian monsoon region. In recent years, researchers in China have been increasingly interested in how will climate change affects species survival and the suitable areas^{4–7}.

With the intensification, generalization, and diversification of the impact of human activities on ecosystems, the spatiotemporal distribution of species has become increasingly intense and complex. The focus of ecosystem dynamics research has rapidly shifted to the patterns and mechanisms of ecosystem changes under human activity interference. Studies have confirmed that human activities have a serious impact on plant diversity⁸. Sayit et al. used MaxEnt to develop the models of predicting suitable habitats of *Calligonum mongolicum* under different climatic conditions and human activities so as to quantitatively demonstrate the different climate change scenarios⁵. In the study of invasive alien plants, Sayit et al. also chose to introduce human activities as an environmental variable, applied MaxEnt and ArcGIS to construct a suitable habitat prediction model for *Solanum*

¹Crop Characteristic Resources Creation and Utilization Key Laboratory of Sichuan Province, Mianyang Academy of Agricultural Sciences, Mianyang 621023, People's Republic of China. ²College of Life Science and Biotechnology, Mianyang Normal University, Mianyang 621000, People's Republic of China. ✉email: 374327731@qq.com

*rostratum*⁹. Different human activity ranges, ways, and intensities can lead to different types of land use (arable land, forest land, shrubland land, and residential land), which directly affects the spatial distribution and diversity of plants^{10,11}. Therefore, taking human activities as prediction variables can more accurately show the potential distribution range and spatial pattern of species in the future.

MaxEnt (the maximum entropy model) invented by Jaynes in 1957, can infer incomplete information, and was used to predict species distributions in 2004. In today's world, MaxEnt has been widely applied in a wide range of fields, including ecology, evolution, and resource management^{12,13}. MaxEnt was more accurate than other niche models, including CLIMEX, GARP, Bioclim and Domain. In spite of incomplete data on species distribution and environmental variables, it can accurately predict distribution areas of species¹⁴. Additionally, MaxEnt has high stability and is applied in the field of animal and plant conservation, including predicting potential habitats for medicinal plants and endangered species^{15,16}. She et al. predicted the distribution hot spots of *Notopterygium incisum*, an important medicinal resource and endangered species in the Three Rivers Source Area under the future climate change scenarios using MaxEnt¹⁷. Ji et al.¹⁸ applied MaxEnt and combined with statistical methods to simulate the current and future distribution of *Paris verticillata*, explored the relationship between its geographical distribution, and understood the development trend of its geographical distribution in the future.

Fritillaria unibracteata P.K.Hsiao & K.C.Hsia (Liliaceae: *Fritillaria*) is a perennial herb (Fig. 1). The dried bulbs of *F. unibracteata* with important medicinal value are widely used to treat lung heat and dry cough, dry cough with less phlegm, yin deficiency and labor cough, expectoration with blood¹⁹. In addition, it can also reduce the permeability of blood vessels and effectively achieve the anti-inflammatory effect^{19,20}. The total saponins and total alkaloids in bulbs have a certain antihypertensive effect on the cardiovascular system^{19,21}. There is a growing market demand for *F. unibracteata* since it has better efficacy than other basic species of Chuanbei, such as *Fritillaria przewalskii*, *Fritillaria cirrhosa* and *Fritillaria delavayi*. As a medicinal plant, the previous research on *F. unibracteata* not only involves its biological characteristics, chemical components and pharmacological effects, but also involves its community ecology. Huang et al.²² found that the distribution of *F. unibracteata* is in the transition from China-Japan forest plant subregion to China-Himalayan plant subregion, which is the most abundant area of alpine flora in the world. Due to the narrow niche and long growth cycle, as well as the deterioration of the ecological environment caused by climate change and people's excessive excavation, the wild resources of *F. unibracteata* have been greatly reduced in recent years and have been listed as a national third-class endangered medicinal material species²². Currently, studies using models to simulate the habitat of *F. unibracteata* have not included human activities as an important variable. Wang et al. analyzed the suitable habitat of *F. unibracteata* in China by using TCMGIS-I system²³. Two reports have applied MaxEnt model to simulate the distribution of *F. unibracteata* with the impact of current climate condition^{24,25}. On the basis of



Figure 1. Habitat of wild *F. unibracteata*.

previous studies, we introduced human activities as a comparative variable, and our purpose was to solve the following issue: (1) analyze the key environmental variables affecting the potential distribution of *F. unibracteata*. (2) simulate the distribution pattern of *F. unibracteata* in eastern Qinghai-Tibetan Plateau. (3) clarify the role of human activities in the geographical distribution of *F. unibracteata*. The results can provide data support for the wild cultivation and introduction of *F. unibracteata*, and provide theoretical basis for the protection and scientific research of the *Fritillaria* genus.

Results

Importance of environmental variables

Without human activities, the percent contribution rate (44.54%) and the regularized training gain (1.96) of elevation (El) were the highest (Fig. 2A and Table 1), indicating that it was the most important variable for the distribution of *F. unibracteata*. When modelling with only variable, the gain (1.63) of min temperature of cold-month (Bio6) ranked second (Fig. 2A), while modelling without Bio6, the score of the model decreased the




Figure 2. Import of environmental variables for prediction based on jackknife test.

Variable	Percent contribution (%)	
	Without human activities	With human activities
Elevation	44.54	45.63
Annual precipitation	28.46	25.82
Mean UV-B of highest month	18.17	15.04
Temperature annual range	3.69	4.10
Potential of hydrogen	1.25	0.94
Min temperature of coldest month	1.04	1.01
Aspect	0.82	0.81
Topsoil sand fraction	0.78	1.65
Isothermality	0.42	0.40
Mean diurnal range	0.26	0.16
Topsoil sand fraction	0.20	0.21
Precipitation of warmest quarter	0.20	0.19
Reference soil depth	0.17	0.39
Mean temperature of warmest quarter	0.02	0.03
Human footprint index	/	3.61

Table 1. Percent contribution of 15 environmental variables.

most, indicating the influence of this variable on simulation. The training gain of mean UV-B of highest month (UV-B3) and annual precipitation (Bio12) exceed 1.3 (Fig. 2A), and their percent contribution rates reached 18.17% and 28.46% respectively (Table 1).

With human activities, the percent contribution rate (45.63%) and the regularized training gain (1.95) of elevation (El) were the highest (Fig. 2B and Table 1), indicating its importance. The regularized training gain (1.58) and the percent contribution rate (15.04%) of mean UV-B of highest month (UV-B3) ranked second and third respectively (Fig. 2B and Table 1). The training gain value for "without variables" was most affected by mean UV-B of the highest month (UV-B3) (Fig. 2B), indicating that its unique information and great impact to the distribution of *F. unibracteata*. The regularized training gain of human footprint index was 1.48 (Fig. 2B), which showed the necessity of introducing it.

By the above comparison method, min temperature of coldest month (Bio6), elevation (El), annual precipitation (Bio12), human footprint index (Hf), mean UV-B of highest month (UV-B3) and temperature annual range (Bio7) were identified to be the dominant environmental variables.

Suitable value range of main environmental variables

When the elevation (El) < 3568.52 m, it had a positive impact on the presence probability of *F. unibracteata*, that was, the presence probability increased with elevation, while when the elevation > 3568.52 m, it had a negative impact on the presence probability (Fig. 3). When the min temperature of coldest month (Bio6) was -10.5°C , the presence probability of *F. unibracteata* decreased steadily after reaching the peak (Fig. 3). The presence probability of *F. unibracteata* increased rapidly to the highest value 10.52 ($P=0.7$) when the human footprint value reached 2.47, then followed by a decrease in probability. According to Fig. 3, the suitable ranges of min temperature of coldest month, annual precipitation, temperature annual range, elevation, human footprint and mean UV-B of highest month were $-16.89--4.96^{\circ}\text{C}$ (Fig. 3A), 416.64–866.96 mm (Fig. 3B), $24.83--31.97^{\circ}\text{C}$ (Fig. 3C), 2879.69–3981.82 m (Fig. 3D), 2.58–23.66 (Fig. 3E) and 7381.92–8574.27 kJ/m^2 (Fig. 3F), respectively.

Geographical distribution and GAP analysis of *F. unibracteata* under current situation

Without human activities, the suitable range of *F. unibracteata* was centered on the highly suitable habitat, and moderately and poorly suitable habitats expanded outward (Fig. 4A, Table 2). The highly suitable habitats ($8.47 \times 10^4 \text{ km}^2$) were distributed in northern Sichuan ($6.45 \times 10^4 \text{ km}^2$), southern Gansu ($1.35 \times 10^4 \text{ km}^2$) and southeastern Qinghai ($0.67 \times 10^4 \text{ km}^2$) in the eastern Qinghai Tibet Plateau. The moderately suitable habitats ($10.43 \times 10^4 \text{ km}^2$) extended to the periphery along the highly suitable habitats ($4.26 \times 10^4 \text{ km}^2$), mainly located in eastern Qinghai ($3.58 \times 10^4 \text{ km}^2$) and southeastern Gansu ($2.24 \times 10^4 \text{ km}^2$) (Fig. 4A and Table 2). The distribution of the poorly suitable habitats were the widest and scattered, including central and southern Gansu ($5.53 \times 10^4 \text{ km}^2$), eastern Qinghai ($11.22 \times 10^4 \text{ km}^2$), western and southern Sichuan ($11.48 \times 10^4 \text{ km}^2$), and eastern Tibet ($4.49 \times 10^4 \text{ km}^2$) (Fig. 4A and Table 2).

Under the influence of human activities, highly, moderately and poorly suitability areas of *F. unibracteata* were $6.64 \times 10^4 \text{ km}^2$, $9.66 \times 10^4 \text{ km}^2$ and $33.8 \times 10^4 \text{ km}^2$, respectively (Fig. 4B and Table 2). The fragmentation of suitable habitats of *F. unibracteata* were more apparent under human activities. The areas of highly, moderately, and poorly suitable habitats decreased by 21.63%, 48.91%, and 30.84%, suggesting a negative correlation between the *F. unibracteata*'s distribution and human activities. GAP analysis showed that the highly suitable habitats of *F. unibracteata* overlapping with the nature reserve was $2.34 \times 10^4 \text{ km}^2$ without human activities (Fig. 4C). With human activities (Fig. 4D), the overlapping area was $1.84 \times 10^4 \text{ km}^2$, accounting for 27.71% of the highly suitable habitats.




Figure 3. Response curves of variables.

Future potential distribution of *F. unibracteata*

Figure 5 showed the future suitable habitats of *F. unibracteata* without human activities. By 2050s, the highly suitable habitats area would decrease from 8.47×10^4 km² to 6.71×10^4 km² (SSP1-2.6), 6.96×10^4 km² (SSP2-4.5) and 7.24×10^4 km² (SSP5-8.5) (Table 2). The moderately suitable habitats would decrease from 10.43×10^4 km² to 9.17×10^4 km² (SSP1-2.6) and 8.91×10^4 km² (SSP5-8.5), while would increase to 10.61×10^4 km² under SSP2-4.5 (Table 2). The poorly suitable habitats would increase from 38.44×10^4 km² to 40.94×10^4 km² (SSP2-4.5) and 44.79×10^4 km² (SSP5-8.5), while would decrease to 36.86×10^4 km² under SSP1-2.6 (Table 2). By 2090s, the highly suitable habitats would decrease to 8.17×10^4 km² (SSP1-2.6) and 7.69×10^4 km² (SSP5-8.5), while would increase to 9.15×10^4 km² under SSP2-4.5 (Table 2). The moderately suitable habitats would increase to 11.87×10^4 km² (SSP1-2.6), 14.82×10^4 km² (SSP2-4.5) and 12.06×10^4 km² (SSP5-8.5) (Table 2). The poorly suitable habitats would increase to 42.01×10^4 km² (SSP1-2.6), 48.57×10^4 km² (SSP2-4.5) and 43.3×10^4 km² (SSP5-8.5) (Table 2).

Figure 6 showed the future suitable habitats of *F. unibracteata* with human activities. By 2050s, the highly suitable habitats area would decrease from 6.64×10^4 km² (Current) to 4.47×10^4 km² (SSP1-2.6), 4.45×10^4 km² (SSP2-4.5) and 4.51×10^4 km² (SSP5-8.5) (Table 2). The moderately suitable habitats would decrease from 9.66×10^4 km² (Current) to 6.84×10^4 km² (SSP1-2.6) and 8.66×10^4 km² (SSP5-8.5), while would increase to




Figure 4. Potential distribution and GAP analysis of *F. unibracteata* under current condition. (A): without human activities; (B): with human activities; (C): without human activities; (D): with human activities. MaxEnt v3.4.4: https://biodiversityinformatics.amnh.org/open_source/maxent/, ArcGIS v10.0: <https://www.arcgis.com/>.

$11.58 \times 10^4 \text{ km}^2$ under SSP2-4.5 (Table 2). The poorly suitable habitats would decrease from $33.8 \times 10^4 \text{ km}^2$ (Current) to $26.6 \times 10^4 \text{ km}^2$ (SSP1-2.6) and $24.51 \times 10^4 \text{ km}^2$ (SSP2-4.5), while would increase to $35.33 \times 10^4 \text{ km}^2$ under SSP5-8.5 (Table 2). By 2090s, the highly suitable habitats area would decrease to $5.28 \times 10^4 \text{ km}^2$ (SSP1-2.6), $5.74 \times 10^4 \text{ km}^2$ (SSP2-4.5) and $5.22 \times 10^4 \text{ km}^2$ (SSP5-8.5) (Table 2). The moderately suitable habitats would

	Scenario	Area of suitable habitats ($\times 10^4 \text{km}^2$)				Changed area compared to current ($\times 10^4 \text{km}^2$)		
		Poorly	Moderately	Highly	Total	Stable	Shrink	Expand
Without human activities	Current	38.44	10.43	8.47	57.34	/	/	/
	2050s, SSP1-2.6	36.86	9.17	6.71	52.74	14.45	4.40	1.24
	2050s, SSP2-4.5	40.94	10.61	6.96	58.51	15.59	3.27	1.72
	2050s, SSP5-8.5	44.79	8.91	7.24	60.94	14.11	4.75	1.83
	2090s, SSP1-2.6	42.01	11.87	8.17	62.05	13.92	4.95	5.72
	2090s, SSP2-4.5	48.57	14.82	9.15	72.54	15.05	3.82	8.55
	2090s, SSP5-8.5	43.30	12.06	7.69	63.05	13.86	5.00	5.55
With human activities	Current	33.80	9.66	6.64	50.1	10.51	5.62	0.62
	2050s, SSP1-2.6	26.61	6.84	4.47	37.92	10.82	5.31	0.68
	2050s, SSP2-4.5	24.51	11.58	4.45	40.54	11.43	4.70	1.41
	2050s, SSP5-8.5	35.33	8.66	4.51	48.5	11.49	4.65	3.74
	2090s, SSP1-2.6	37.70	10.45	5.28	53.43	11.85	4.29	4.91
	2090s, SSP2-4.5	38.53	11.59	5.74	55.86	10.87	5.26	2.96
	2090s, SSP5-8.5	36.14	9.00	5.22	50.36	10.51	5.62	0.62

Table 2. The areas of suitable habitats and their changes under future climate change scenarios.

increase to $10.45 \times 10^4 \text{ km}^2$ (SSP1-2.6) and $11.59 \times 10^4 \text{ km}^2$ (SSP2-4.5), while decrease to $9 \times 10^4 \text{ km}^2$ under SSP5-8.5 (Table 2). The poorly suitable habitats would increase to $37.7 \times 10^4 \text{ km}^2$ (SSP1-2.6), $38.53 \times 10^4 \text{ km}^2$ (SSP2-4.5) and $36.14 \times 10^4 \text{ km}^2$ (SSP5-8.5) (Table 2).

Changes in distribution of suitable habitats in the future

Figure 7 showed the stable, expand and shrink area of the suitable habitats in 2050s and 2090s without human activities compared with current situation. In 2050s, the proportion of the stable habitats to the current was 76.65% (SSP1-2.6), 82.65% (SSP2-4.5) and 74.8% (SSP5-8.5) respectively (Fig. 7A,B,C), while in 2090s it was 73.78% (SSP1-2.6), 79.75% (SSP2-4.5) and 73.49% (SSP5-8.5) respectively (Fig. 7D,E,F). The proportion of the shrink habitats to the current was 23.35% (SSP1-2.6), 17.35% (SSP2-4.5) and 25.2% (SSP5-8.5) respectively in 2050s (Fig. 7A,B,C), and 26.22% (SSP1-2.6), 20.25% (SSP2-4.5) and 26.51% (SSP5-8.5) in 2090s (Fig. 7D,E,F). The proportion of the expand habitats to the future was 7.91% (SSP1-2.6), 9.95% (SSP2-4.5) and 11.48% (SSP5-8.5) respectively in 2050s (Fig. 7A,B,C), and 29.14% (SSP1-2.6), 36.23% (SSP2-4.5) and 28.6% (SSP5-8.5) in 2090s respectively (Fig. 7D,E,F). By overlaying layers in ArcGIS, the suitability changes of the suitable habitats in 2050s and 2090s were obtained (Fig. 7G,H). The results showed that Aba, Ruergai, Hongyuan, Songpan, Markang, Heishui, Jinchuan, Lixian, Maoxian, Xiaojin, Danba, Wenchuan, Baoxing, Kangding and Luding in the western Sichuan Plateau, Maqu, Luqu, Xiahe and Hezuo in the western Gannan Plateau, Henan, Zeku, Tongren and Jiuzhi in the eastern Qinghai were identified as stable habitats for the growth and distribution of *F. unibracteata* (Fig. 7G,H). The expand habitats were distributed in the western Sichuan Plateau, the east of Tibet, and the northeast of Qinghai Province in strip or point form (Fig. 7G). The shrink habitat was scattered around the stable area (Fig. 7G,H).

Figure 8 showed the stable, expand and shrink area of the suitable habitats in 2050s and 2090s with human activities compared with current situation. In 2050s, the proportion of the stable habitats to the current was 65.15% (SSP1-2.6), 67.06% (SSP2-4.5) and 70.86% (SSP5-8.5) respectively (Fig. 8A,B,C), while in 2090s it was 71.18% (SSP1-2.6), 73.43% (SSP2-4.5) and 67.38% (SSP5-8.5) respectively (Fig. 8D,E,F). The proportion of the shrink habitats to the current was 34.85% (SSP1-2.6), 34.84% (SSP2-4.5) and 32.94% (SSP5-8.5) respectively in 2050s (Fig. 8A,B,C), while in 2090s it was 28.82% (SSP1-2.6), 26.57% (SSP2-4.5) and 32.62% (SSP5-8.5) (Fig. 8D,E,F). The proportion of the expand habitats to the future was 5.58% (SSP1-2.6), 5.88% (SSP2-4.5) and 10.95% (SSP5-8.5) respectively in 2050s respectively (Fig. 8A,B,C), while in 2090s it was 24.56% (SSP1-2.6), 29.3% (SSP2-4.5) and 21.4% (SSP5-8.5) respectively (Fig. 8D,E,F). By comparison, the geographical scope of the stable habitat was somewhat smaller than that without human activities, but the administrative coverage was basically the same. The proportion of the expand habitat tended to decrease with human activities, while the proportion of shrink habitat tended to increase. By overlaying layers in ArcGIS, the suitability change of the suitable habitat in 2050s and 2090s was obtained (Fig. 8G,H,I).

Centroid variations of the suitable habitats under climate change scenarios

Without human activities (Fig. 9A), under SSP1-2.6, the centroid of the highly suitable habitats would move 11.62 km from Hongyuan (Current/32.62°N, 102.51°E) to southeast to Hongyuan (2050s/32.52°N, 102.56°E), then 56.4 km to northwest to Aba (2090s/32.65°N, 102.01°E). By 2090s, the centroid would be displaced 50.48 km to the northwest. Under SSP2-4.5, the centroid would move 7.58 km from Hongyuan (Current/32.62°N, 102.51°E) to Hongyuan (2050s/32.59°N, 102.58°E), then 43.76 km to northwest to Aba (2090s/32.84°N, 102.23°E). By 2090s, the centroid would generally displace 36.28 km to the northwest. Under SSP5-8.5, the centroid of the highly suitable habitats would move 2.47 km from Hongyuan (Current/32.62°N, 102.51°E) to southwest to Hongyuan




Figure 5. Potential distribution of *F. unibracteata* in the future without human activities. (A): 2050s, SSP1-2.6; (B): 2050s, SSP2-4.5; (C): 2050s, SSP5-8.5; (D): 2090s, SSP1-2.6; (E): 2090s, SSP2-4.5; (F): 2090s, SSP5-8.5. MaxEnt v3.4.4: https://biodiversityinformatics.amnh.org/open_source/maxent/, ArcGIS v10.0: <https://www.arcgis.com/>.

(2050s/32.6°N, 102.5°E), then 16.07 km to northwest to Hongyuan (2090s/32.74°N, 102.43°E). By 2090s, the centroid would generally displace 14.4 km to the northwest (Fig. 9A).

With human activities (Fig. 9B), under SSP1-2.6, the centroid of the highly suitable habitats would displace 23.59 km from Hongyuan (Current/32.61°N, 102.61°E) to southeast to Hongyuan (2050s/32.37°N, 102.62°E), then 35.97 km to northwest to Hongyuan (2090s/32.59°N, 102.33°E). By 2090s, the centroid would displace 27.84 km to the southwest. Under SSP2-4.5, the centroid would move 23.82 km from Hongyuan (Current/32.61°N, 102.61°E) to southeast to Songpan (2050s/32.38°N, 102.68°E), then 52.91 km to northwest to Hongyuan (2090s/32.82°N, 102.38°E). By 2090s, the centroid may displace 30.98 km to the northwest. The centroid of the highly suitable habitats would move 4.69 km from Hongyuan (Current/32.61°N, 102.61°E) to southwest to Hongyuan (2050s/32.56°N, 102.63°E) under SSP5-8.5, then 12.18 km to northwest to Hongyuan (2090s/32.69°N, 102.62°E). By 2090s, the centroid may generally displace 7.84 km to the northeast (Fig. 9B).

Evaluation of models

Without human activities, based on the training data and test data of the MaxEnt under the current situation, the AUC values were 0.987 ± 0.001 and 0.983 ± 0.013 respectively. Under future climate scenarios, the AUC values for training data were 0.986 ± 0.001 – 0.988 ± 0.001 , while for the test data were 0.979 ± 0.021 – 0.984 ± 0.006 (Table 3).

With human activities, the AUC values of the training data and test data of the MaxEnt model under current conditions were 0.987 ± 0.001 and 0.982 ± 0.015 respectively. Under climate change scenarios, the AUC values of the training data were 0.986 ± 0.001 – 0.989 ± 0.000 , and that of the test data were 0.980 ± 0.015 – 0.984 ± 0.006 (Table 3).




Figure 6. Potential distribution of *F. unibracteata* in the future with human activities. (A): 2050s, SSP1-2.6; (B): 2050s, SSP2-4.5; (C): 2050s, SSP5-8.5; (D): 2090s, SSP1-2.6; (E): 2090s, SSP2-4.5; (F): 2090s, SSP5-8.5. MaxEnt v3.4.4: https://biodiversityinformatics.amnh.org/open_source/maxent/, ArcGIS v10.0: <https://www.arcgis.com/>.

Discussion

As a comprehensive environmental factor that affects plant growth and distribution, altitude largely controls the changes in other environmental factors, such as temperature, precipitation, and light intensity, which exhibit regular changes with altitude^{26–29}. Altitude is an important factor affecting *Fritillaria* plants. Jiang et al. found that the dominant factors related to the growth of wild plants of *Fritillaria cirrhosa* are altitude, precipitation in September, precipitation in November, and vegetation type. Zhao et al. demonstrated through simulation that although the dominant factors affecting the habitat suitability of *F. cirrhosa*, *F. unibracteata*, *Fritillaria przewalskii*, and *Fritillaria delavayi* are different, but altitude plays an important role in all species²⁴. *Fritillaria unibracteata* is a kind of alpine plant, which mainly grows in the high altitude area of 2800–4400 m on the eastern edge of Qinghai Tibet Plateau. In this area, altitude is an important factor controlling the combination form and variation degree of other environmental and biological factors^{30,31}. The growth status and survival strategy of plants growing in extreme alpine environment can show adaptability to altitude³². By comparing, the percent contribution rate of elevation was 44.54% without human activities and 45.63% with human activities, which were the most important variables under both scenarios. *F. unibracteata* grows and reproduces poorly at low altitudes, while the bulb biomass and total plant biomass decreased as altitude increases in high altitudes³³. Xu et al. demonstrated that the effects of altitude gradient and life history stage on single leaf area, plant height, and specific leaf area of *F. unibracteata* were very significant, and there was obvious interaction³⁴. Chen et al. believed that the reproductive organs of *F. unibracteata* will be affected by altitude, which is mainly related to snow melting time, temperature, ultraviolet intensity, flower visiting insects³⁵. Therefore, altitude has a great restrictive effect on the growth of *F. unibracteata*. A response curve revealed that the suitable elevation range was 2879.7–3981.82 m, confirming that altitude plays an important role in *F. unibracteata* distribution. Previous studies have shown




Figure 7. Changes in distribution of suitable habitats in the future without human activities. (A): 2050s, SSP1-2.6; (B): 2050s, SSP2-4.5; (C): 2050s, SSP5-8.5; (D): 2090s, SSP1-2.6; (E): 2090s, SSP2-4.5; (F): 2090s, SSP5-8.5. MaxEnt v3.4.4: https://biodiversityinformatics.amnh.org/open_source/maxent/, ArcGIS v10.0: <https://www.arcgis.com/>.

that within the altitude range of 2371–3076 m, the growth season length of *F. unibracteata* is relatively longer, which is conducive to its accumulation of sufficient nutrients to meet the needs of growth and development²⁶.




Figure 8. Changes in distribution of suitable habitats in the future with human activities. (A): 2050s, SSP1-2.6; (B): 2050s, SSP2-4.5; (C): 2050s, SSP5-8.5; (D): 2090s, SSP1-2.6; (E): 2090s, SSP2-4.5; (F): 2090s, SSP5-8.5. MaxEnt v3.4.4: https://biodiversityinformatics.amnh.org/open_source/maxent/, ArcGIS v10.0: <https://www.arcgis.com/>.

According to the statistical data, in the high-altitude areas of western Sichuan, southeast Qinghai and southern Gansu, the annual precipitation was about 400–1400 mm, and the temperature annual range was large, which was conducive to the accumulation of nutrients and promote the growth of bulb of *F. unibracteata*. Based on




Figure 9. Variations of the centroids of the suitable habitats under climate change scenarios without human activities (A) and with human activities (B). ArcGIS v10.0: <https://www.arcgis.com/>.

the response curve, the suitable range for the min temperature of coldest month was $-16.84-4.95^{\circ}\text{C}$, while the average temperature during the coldest month was about -2°C , which could ensure its safe wintering. For the northwest regions with relatively less rainfall (Xinjiang, Inner Mongolia and western Gansu), or the southern coastal areas with more rainfall, it is not conducive to the accumulation of nutrients of *F. unibracteata*, so its distribution in these areas is limited. In this study, the suitable range of main variables calculated by MaxEnt was basically consistent with the ecological environment distributed in the field, that was, high altitude, low temperature, thin air, strong sunshine, clear dry and rainy seasons and large temperature diurnal range^{34,35}.

The distribution of species can be limited by climate factors at large scales. However, many studies have proved that the use of climate factors separately will lead to a wider predicted niche. Now the most viewpoints are that even in the suitable area, the high fragmentation of natural habitat caused by human activities will hinder the expansion of species. A study by Xu et al. assessed whether and how human activities have changed the extent of 9701 vascular plants' climatic potential ranges in China, and they found narrow-ranged species showed negative

Scenario	With human activities		Without human activities	
	Training data	Test data	Training data	Test data
Current	0.987 ± 0.001	0.982 ± 0.015	0.987 ± 0.001	0.983 ± 0.013
2050s, SSP1-2.6	0.987 ± 0.001	0.980 ± 0.015	0.987 ± 0.001	0.980 ± 0.016
2050s, SSP2-4.5	0.988 ± 0.001	0.982 ± 0.012	0.986 ± 0.001	0.979 ± 0.021
2050s, SSP5-8.5	0.986 ± 0.001	0.980 ± 0.013	0.987 ± 0.001	0.982 ± 0.014
2090s, SSP1-2.6	0.988 ± 0.001	0.982 ± 0.009	0.988 ± 0.000	0.984 ± 0.006
2090s, SSP2-4.5	0.989 ± 0.000	0.984 ± 0.004	0.986 ± 0.001	0.983 ± 0.007
2090s, SSP5-8.5	0.989 ± 0.000	0.984 ± 0.006	0.988 ± 0.001	0.984 ± 0.007

Table 3. AUC values of models.

range-filling relationships to these human indicators³⁶. Zhang et al.'s simulation showed that the modern suitable area of *Rosa persica* was significantly reduced after the addition of human footprint data³⁷. Cao et al. compared the impact of human footprint on the suitability of *Swertia przewalskii*, a unique plant in the Qinghai Tibet Plateau, and revealed that after the introduction of this variable, the potential habitat of *S. przewalskii* species showed a fragmentation trend and the area decreased by 32%⁶. In view of this, in order to more comprehensively reflect the ecological environment of *F. unibracteata* and obtain accurate results, we not only selected the commonly used bioclimatic variables, but also supplemented altitude, soil factors and human activities. According to our results, human activities had a high percentage impact on the growth suitability of *F. unibracteata* (3.61%). It can be concluded that *F. unibracteata*'s growth, habitat and reproduction may be affected by a series of human activities such as population density, land use, roads and railways as well as construction.

Under current condition, the highly suitable habitats of *F. unibracteata* were mostly located in northern Sichuan, southern Gansu and southeastern Qinghai and the total area was 6.64×10^4 km². Field investigation and literature review showed that there were cultivation bases of *F. unibracteata* in Western Sichuan Plateau and southern Qinghai located in the predicted highly suitable habitats^{38,39}. Liu et al.'s prediction suggested that the most suitable growth areas of *F. unibracteata* with an area of 5.62×10^4 km² were dominantly distributed in western and northern Sichuan, south Gansu and south Qinghai, which is consistent with our result²⁵. Wang et al. predicted that the best suitable areas for *F. unibracteata* were Aba, Ganzi in Sichuan, and southwest Tibet, while Qinghai and Gansu were unsuitable areas, which was different from our results²³. Our simulation showed that under the impact of human activities, the area of highly, moderately and poorly suitable habitats decreased by 21.63%, 48.91% and 30.84% respectively. Most people believed that even in areas with suitable climate, the high fragmentation of natural habitats caused by human over use of land would hinder the expansion of species habitats^{5,40}. Combined with the prediction results and field observation, we speculated that an important reason for the area reduction of *F. unibracteata* was the destruction of land and habitat change caused by local urban expansion and road construction. In addition, animal husbandry was one of the important industries in Qinghai Tibet Plateau, therefore the impact of grazing could not be ignored.

Recently, it was found that the suitable areas of several medicinal plants were decreasing under future climate change scenarios. Ji et al. showed that under RCP 2.6, RCP 4.5, RCP 6.0 and RCP 8.5 scenarios, the total suitable area of *Paris verticillata* in China showed a decreasing trend in 2050s and 2070s¹⁸. A study by Liu et al. calculated the change in the highly suitable area of *Fritillariae Cirrhosae Bulbus* under SSP1-2.6, SSP2-4.5 and SSP5-8.5, and found that the highly suitable area decreased in size, and the northwest area would be the geometric center of the total suitable area⁴¹. Wei et al. confirmed that the most suitable area of *Fritillaria walujewii* will decrease in the middle and south of Yili River Valley and Tacheng area⁴². However, some scholars have reached different conclusions. Guan et al. believed that climate warming can promote the growth of some plants, and the suitable area of *Quercus acutissima* will increase in general under future climate conditions⁴³. This study found that under human influences, *F. unibracteata* habitats are likely to decrease significantly in 2050s and 2090s. The size of species distribution areas is an important characteristic of species, and narrow range species often have specific environmental needs and may be more sensitive to habitat changes. Under climate change and intense human activities, the distribution area of narrow range species is likely to shrink, while the distribution area of widely distributed species is likely to expand⁴⁴⁻⁴⁷. *F. unibracteata* is a typical narrow niche plant, this may explain the significant reduction in its habitat after adding human activities index as environmental variable.

Under the three scenarios, the centroid of *F. unibracteata* would move from the current position to the low latitude area first, and then to the high latitude area. As a typical alpine plant, *F. unibracteata* has the characteristics of cold tolerance, humidity preference, fear of high temperature and shade preference^{19,48}. We speculated that the change of suitable habitat and the moving trend of centroid in the future were the result of the dual influence of temperature and precipitation. Studies have shown that the increase of regional temperature and precipitation will destroy the habitat of some alpine plants^{49,50}. By the 2090s, the increasing trend of temperature and precipitation in the Qinghai Tibet Plateau will be more obvious, and the suitability of the original growth area of *F. unibracteata* will be further reduced.

With the development of green and healthy industries, medicinal plant resources have attracted increasing attention. However, the contradiction between the demand for traditional Chinese medicine and the increasing scarcity of resources is becoming increasingly prominent^{51,52}. Due to strict habitat requirements and indiscriminate mining, the wild resources of *F. unibracteata* have been depleted. Therefore, determining the optimal area

for artificial introduction and cultivation of medicinal plants through habitat suitability research is one of the effective methods to fundamentally alleviate the pressure on wild medicinal plant resources. The protection of rare and endangered plant resources is not only the key component of biodiversity protection and sustainable utilization, but also one of the important tasks of nature protection. In recent years, people have gradually realized that the loss of species habitat has become the biggest threat to biodiversity. The destruction of the original habitat by reservoirs, ponds and farmland has resulted in the disappearance of many wild populations of *F. unibracteata* in Qinghai and Sichuan^{24,53,54}. *F. unibracteata* is urgently in need of better protection. Our GAP analysis showed that there is a large protection GAP of *F. unibracteata* in northern Sichuan, southeastern Qinghai and southwestern Gansu. GAP analysis also showed that Taohe, Siguniangshan, Wolong, Sanjiangyuan, Zoige and other more than ten national nature reserves were located within the highly suitable habitats. Therefore, combined with the distribution characteristics in National Nature Reserves and the current national reserve management policy, we put forward the following suggestions: (1) In the areas where the highly suitable habitats overlaps with the National Nature Reserve, the core protection area of *F. unibracteata* will be established to protect its habitat completely, and the core protection area will also play a role in the conservation and radiation of its germplasm resources. (2) Studies revealed that it was feasible to transplant *F. unibracteata* artificially to a suitable habitat^{48,55}. Therefore, for the moderately and poorly suitable habitats, in the areas with dense population, frequent human activities, which have caused serious harm to the groups and germplasm resources of *F. unibracteata*. Depending on the environmental conditions such as altitude, soil, temperature and precipitation, they can be transplanted into reservoirs, farms, and parks suitable for their survival and free of human disturbance so as to achieve the purpose of ex-situ genetic resources protection. (3) In the general control area of the National Nature Reserve, the field distribution survey of *F. unibracteata* should be further carried out, and the community survey should be carried out in relevant villages and towns to understand the use of local residents' natural resources in the community. On this basis, local residents are encouraged to understand, support and participate in the protection of *F. unibracteata* by carrying out environmental education, science dissemination and seeking alternative livelihoods.

Methods

Study area and occurrence records of *F. unibracteata* in China

The study area is predominantly in the eastern edge of the Qinghai Tibet Plateau (Fig. 10), with a large space span, belonging to the plateau marine climate and continental cold dry climate. The air in this area is thin, the annual average temperature is mostly 7 ~ 10 °C, the highest temperature can reach 30 °C, and the lowest temperature is as low as -20 ~ -30 °C. Seasonal frozen layer is distributed in the area above 4000 m, with the maximum thickness of 50 cm. The vegetation is mainly alpine meadow and the overall vegetation coverage is generally low. There is almost no vegetation in the area above 4500 m, with obvious vertical zoning⁵⁶.

Occurrence data of *F. unibracteata* were acquired from the National Specimen Information Infrastructure (NSII, <http://www.nsi.org.cn/>), the Chinese Virtual Herbarium (CBV, <https://www.cvh.ac.cn/>), the Global Biodiversity Information Facility (GBIF, <https://www.gbif.org/>), field survey (Our field survey only recorded the




Figure 10. Study area and locations of occurrence records of *F. unibracteata* in China. ArcGIS v10.0: <https://www.arcgis.com/>.

Code	Variable	Resolution	Unit
Bio2	Mean diurnal range	2.5'	°C
Bio3	Isothermality	2.5'	/
Bio6	Min temperature of coldest month	2.5'	°C
Bio7	Temperature annual range	2.5'	°C
Bio10	Mean temperature of warmest quarter	2.5'	°C
Bio12	Annual precipitation	2.5'	mm
Bio14	Precipitation of warmest quarter	2.5'	mm
El	Elevation	2.5'	m
Aspect	Aspect	2.5'	°
PH	Potential of hydrogen	2.5'	/
T-C	Organic carbon pool topsoil	2.5'	%
T-sand	Topsoil sand fraction	2.5'	%
Depth	Reference soil depth	2.5'	m
UV-B3	Mean UV-B of Highest Month	2.5'	kJ/m ²
Hf	Human footprint index	2.5'	/

Table 4. Environmental variables used to predict the potential geographic distribution of *F. unibracteata*.

coordinates of occurrence data and did not involve field sampling), and literature. *F. unibracteata* distribution records were processed according to a literature review (Wang et al. 2019; Liu et al. 2021). The first step included using Baidu's coordinate picking system (<https://api.map.baidu.com/lbsapi/getpoint>) to determine longitudes and latitudes accurate to the town level. The second step involved using Microsoft Excel (2010) to remove duplicate records. Third, each point was measured with respect to its cell grid center, and the nearest point was selected. To establish MaxEnt, 123 distribution points of *F. unibracteata* were retained after the above procedures (Fig. 10).

Environmental variables

Five independent environmental variables were included in this study (Table S1). Eliminating the influence of multicollinearity was important for simulation, so Pearson correlation coefficient was adopted^{57,58}. We first calculated the percent contribution rate of 19 variables using MaxEnt, and we retained the variables with higher percent contribution rates (Table S2). Following that, SPSS was used to analyze Pearson's coefficients between two variables with percent contributions greater than 0 for 123 *F. unibracteata* occurrences. Thirdly, the higher coefficient was retained if the percentage contribution of the variables with the absolute value of the coefficient exceeded 0.85 (Table S3). The prediction model for *F. unibracteata* included 15 variables in addition to elevation (Table 4).

Modelling process

MaxEnt software operation procedure are presented as: (1) Occurrence data of *F. unibracteata* were imported into the "sample" and "environmental layer" data boxes of MaxEnt software (V3.4.4) in CSV and ASC formats (including bioclimatic variables, soil data, elevation and human footprint). (2) "Create response curves" and "Do jackknife to measure variable importance" were selected respectively to analyze the relationship between variables and presence probability of *F. unibracteata* and measure the importance of variables. (3) In the initial model, "Random test percentage" was set to 25%, while in the reconstructed model, "random seed" was selected, and the "replicates" was set to 10^{12,59}. An area value under the ROC curve, such as AUC value ranges from 0.5 to 1, is a highly recognized diagnostic test evaluation index^{60,61}.

Data availability

The data that support the findings of this study are openly available in the Science Data Bank at <https://www.doi.org/10.57760/sciencedb.06988>.

Received: 21 April 2023; Accepted: 17 September 2023

Published online: 22 September 2023

References

- Zou, J., Teng, F. & Fu, S. The latest progress in socioeconomic assessment of the mitigation of climate change-review of the Ipcc fifth assessment Wg iii report. *Adv. Clim. Chang. Res.* **10**, 313–322 (2015).
- Frame, D. J. *et al.* Climate change attribution and the economic costs of extreme weather events: a study on damages from extreme rainfall and drought. *Clim. Change.* **162**, 781–797 (2020).
- Kumar, S. & Stohlgren, T. J. Maxent modeling for predicting suitable habitat for threatened and endangered tree *Canacomyrica Monticola* in New Caledonia. *Afr. J. Ecol. Ecosyst.* **6**, 1–5 (2019).
- Liu, L. *et al.* Simulation of potential suitable distribution of alnus cremastogyne burk. In China under climate change scenarios. *Ecol. Indic.* **133**, 108396 (2021).
- Sayit, H., Nurbay, A., Li, X. P., Shao, H. & Arman, J. Effects of climate change and human activities on the distribution pattern of *Calligonum Mongolicum* Turcz. *Arid Zone Res.* **35**, 1450–1458 (2018).
- Cao, Q. *et al.* Impacts of human activities and environmental factors on potential distribution of *Swertia Przewalskii* Pissjauk. An endemic plant in qing-tibetan plateau, using Maxent. *Plant Sci. J.* **39**, 22–31 (2021).

7. Wang, R. L., Yang, H., Wang, M. T., Zhang, Z. & Li, Q. Predictions of potential geographical distribution of *Diaphorina Citri* (Kuwayama) in China under climate change scenarios. *Sci. Rep.* **10**, 1–9 (2020).
8. Cao, W., Huang, L., Xiao, T. & Wu, D. Effects of human activities on the ecosystems of China's national nature reserves. *Acta Ecol. Sin.* **39**, 1338–1350 (2019).
9. Sayit, H., Nurbay, A., Arman, J., Shao, H. & Vinra, Y. Impact of human activities on potential distribution of *solanum rostratum* dunal in Xinjiang. *Acta Ecol. Sin.* **39**, 629–636 (2019).
10. Liu, A., Yang, T., Xu, W., Shangguan, Z. & He, J. S. Status, issues and prospects of belowground biodiversity on the tibetan alpine grassland. *Biodivers. Sci.* **26**, 972–987 (2018).
11. Gao, R. R. *et al.* Chinese herbal medicine resources: where we stand. *Chin. Herb. Med.* **12**, 3–13 (2020).
12. Elith, J. *et al.* A statistical explanation of maxent for ecologists. *Divers. Distrib.* **17**, 43–57 (2015).
13. Phillips, S. J., Anderson, R. P. & Schapire, R. E. Maximum entropy modeling of species geographic distributions. *Ecol. Model.* **190**, 231–259 (2006).
14. Petitpierre, B., Kuffer, C., Broennimann, O. & Randin, C. Climatic niche shifts are rare among terrestrial plant invaders. *Science* **335**, 1344–1348 (2012).
15. Mbatudde, M., Mwanjololo, M., Kakudidi, E. K. & Dalitz, H. Modelling the potential distribution of endangered *prunus Africana* (Hook. F.) Kalkm. In East Africa. *Afr. J. Ecol.* **50**, 393–403 (2012).
16. Chen, L. D., Hu, W., Li, D. Q., Cheng, D. M. & Zhong, A. W. Prediction of suitable distribution areas of the endangered plant wild *nelumbo nucifera gaertn.* In China. *Plant Sci. J.* **37**, 731–740 (2019).
17. She, Y. D. *et al.* Suitable distribution of *Notopterygium Incisum* in the three rivers headwater region under climate change. *Ecol. Environ. Sci.* **30**, 2033–2041 (2021).
18. Ji, L. T., Zheng, T. Y., Chen, Q., Zhong, J. J. & Kang, B. Responses of potential suitable area of *Paris Verticillata* to climate change and its dominant climate factors. *Chin. J. Appl. Ecol.* **31**, 89–96 (2020).
19. Zhou, Q. M. *et al.* Chemical constituents from *Fritillaria Unibracteata*. *J. Chin. Med. Mater.* **39**, 2237–2239 (2016).
20. Jing, C., Ta, X., Hong, Z., Li, Z. Y. & Liu, S. Q. Screening, identification and antimicrobial activity of alkaloid produced by endophytic actinomycetes from *Fritillaria Unibracteata* in Western Sichuan Plateau. *China J. Chin. Materia Med.* **42**, 4582–4587 (2017).
21. Zhou, N., Guo, D. Q., Li, S., Chen, Q. Y. & Qin, Y. Comparative contents of four alkaloids in bulbs of *Fritillaria Taipaiensis* and *Fritillaria Unibracteata*. *Food Science.* **35**, 133–136 (2014).
22. Huang, L. Q., Xiao, P. G. & Wang, Y. Y. *Endangered Medicinal Plant Resources Investigation of China*, Shanghai Science and Technology Press (Shanghai, 2012).
23. Wang, Y. *et al.* Ecological evaluation of suitable producing areas of *Fritillaria Unibracteata* based on Tcmgis- I. *World Sci. Technol.-Modern. Trad. Chin. Med. Mater. Medica.* **8**, 122–124 (2006).
24. Zhao, W. L. *et al.* Distribution of habitat suitability for different sources of *fritillariae cirrhosae* bulb. *Chin. J. Ecol.* **37**, 1037–1042 (2018).
25. Liu, Y., Zhou, S., Xie, D., Huang, J. & Xingjin, H. E. Potential distribution of *Fritillaria Unibracteata* predicted by the Maxent model. *Guihaia.* **38**, 352–360 (2018).
26. Xu, B., Wang, J. N. & Shi, F. S. Impacts of ontogenetic and altitudinal changes on morphological traits and biomass allocation patterns of *Fritillaria Unibracteata*. *J. Mt. Sci.* **17**, 83–94 (2020).
27. Centenaro, G. *et al.* Silvicultural management and altitude prevail on soil properties and fungal community in shaping understorey plant communities in a mediterranean pine forest. *Sci. Total Environ.* **858**, 159860 (2023).
28. Dostalek, T., Rokaya, M. B. & Munzbergova, Z. Altitude, habitat type and herbivore damage interact in their effects on plant population dynamics. *PLoS ONE* **13**, e209149 (2018).
29. Yirga, F., Marie, M., Kassa, S. & Haile, M. Impact of altitude and anthropogenic disturbance on plant species composition, diversity, and structure at the wof-washa highlands of Ethiopia. *Heliyon.* **5**, e2284 (2019).
30. Hou, Y., Zhao, W., Hua, T. & Pereira, P. Mapping and assessment of recreation services in qinghai-tibet plateau. *Sci. Total Environ.* **838**, 156432 (2022).
31. Liu, J., Xin, Z., Huang, Y. & Yu, J. Climate suitability assessment on the qinghai-tibet plateau. *Sci. Total Environ.* **816**, 151653 (2022).
32. Körner, F. C. & Fabbro, T. Altitudinal differences in flower traits and reproductive allocation. *Flora* **199**, 70–81 (2004).
33. Chen, W. N. *et al.* Variation of phenology and bulbs of *Fritillaria Unibracteata* along altitudinal gradients. *Acta Pratacul. Sin.* **21**, 319–324 (2012).
34. Xu, B., Wang, J., Guo, H., Shi, F. & Wu, N. Morphological adaptation of wild *Fritillaria Unibracteata* to alpine conditions in the eastern Qinghai-Tibet plateau. *Chin. J. App. Environ. Biol.* **20**, 955–961 (2014).
35. Chen, W. N. *et al.* Reproductive organ variation in *Fritillaria Unibracteata* along an altitudinal gradient. *Acta Pratacul. Sin.* **24**, 155–163 (2015).
36. Xu, W. B., Svenning, J. C., Chen, G. K., Zhang, M. G. & Ma, K. P. Human activities have opposing effects on distributions of narrow-ranged and widespread plant species in China. *Proc. Natl. Acad. Sci.* **116**, 266–274 (2019).
37. Zhang, X. L., Deng, T., Luo, L. & Li, J. Y. Prediction of potential suitable area of *Rosa Persica* and study on its vulnerable mechanism. *Acta Botan. Boreali-Occiden. Sin.* **41**, 1570–1582 (2021).
38. Xu, Y. *et al.* Quality comparison of cultivated *fritillaria cirrhosa* and cultivated *F. Unibracteata* with different harvesting time. *West China J. Pharm. Sci.* **33**, 515–518 (2018).
39. Zhang, L. *et al.* *Fritillaria Unibracteata* in Barkam: the effect of cultivation methods on growth. *Chin. Agric. Sci. Bull.* **32**, 69–74 (2016).
40. Dang, Y. C., He, H. S., Zhao, D. D., Sunde, M. & Du, H. B. Quantifying the relative importance of climate change and human activities on selected wetland ecosystems in China. *Sustainability.* **12**, 912 (2020).
41. Liu, L. *et al.* Simulation of potential suitable distribution of original species of *fritillariae cirrhosae* bulb in China under climate change scenarios. *Environ. Sci. Pollut. Res.* **29**, 22237–22250 (2021).
42. Wei, B., Ma, S., Song, J., He, L. & Li, X. Prediction of the potential distribution and ecological suitability of *Fritillaria Walujewii*. *Acta Ecol. Sin.* **39**, 228–234 (2019).
43. Guan, X. Y., Shi, W. & Cao, K. F. Effect of climate change in future on geographical distribution of widespread *Quercus Acutissima* and analysis of dominant climatic factors. *J. Trop. Subtrop. Botany.* **26**, 661–668 (2018).
44. Cao, W., Wu, D., Huang, L., Pan, M. & Huhe, T. Determinizing the contributions of human activities and climate change on greening in the Beijing-Tianjin-Hebei region. *China. Sci Rep.* **11**, 21201 (2021).
45. Deng, G. *et al.* Response of vegetation variation to climate change and human activities in semi-arid swamps. *Front. Plant Sci.* **13**, 990592 (2022).
46. Hou, Q., Ji, Z., Yang, H. & Yu, X. Impacts of climate change and human activities on different degraded grassland based on Ndvi. *Sci Rep.* **12**, 15918 (2022).
47. Xiong, Y. *et al.* Influence of human activities and climate change on wetland landscape pattern—a review. *Sci. Total Environ.* **879**, 163112 (2023).
48. Wang, X. *et al.* Similarities and differences between the cultivated *Fritillaria Unibracteata* and *Fritillaria Unibracteata* Var. *Wabuensis*. *West China J. Pharm. Sci.* **35**, 650–654 (2020).
49. Meng, X. H. *et al.* Review of climate change and its environmental influence on the three-river regions. *Plateau Meteorol.* **39**, 1133–1143 (2020).

50. Wei, Y. Q., Lu, H. Y., Wang, J. N., Sun, J. & Wang, X. F. Responses of vegetation zones, in the Qinghai-Tibetan Plateau, to climate change and anthropogenic influences over the last 35 years. *Pratacult. Sci.* **36**, 1163–1176 (2019).
51. Sen, T. & Samanta, S. K. Medicinal plants, human health and biodiversity: a broad review. *Adv. Biochem. Eng. Biotechnol.* **147**, 59–110 (2015).
52. Yan, B. *et al.* A review on the plant resources of important medicinal licorice. *J. Ethnopharmacol.* **301**, 115823 (2023).
53. Zhang, H. J. Research progress of *Fritillaria Unibracteata* Hsiao Et K.C.Hsia. *J. Anhui Agricult. Sci.* **39**, 2067–2069 (2011).
54. Zhao, W. *et al.* Review on in pharmacological effects of *Fritillaria*. *Shanghai J. Tradit. Chin. Med.* **52**, 97–100 (2018).
55. Guo, F. X., Chen, Y., Zhang, J. F. & Bai, G. Abloom phenology and pollination mode of endangered plant *Fritillaria Unibracteata*. *Acta Agrestia Sinica.* **25**, 1324–1332 (2017).
56. Xu, B., Wang, J. N., Shi, F. S., Gao, J. & Wu, N. Adaptation of biomass allocation patterns of Wild *Fritillaria Unibracteata* to alpine environment in the eastern Qinghai-Xizang plateau. *Chin. J. Plant Ecol.* **37**, 187–196 (2013).
57. Guan, L. *et al.* Potential distribution of *Blumea Balsamifera* in China using maxent and the ex situ conservation based on its effective components and fresh leaf yield. *Environ. Sci. Pollut. Res.* **29**, 44003–44019 (2022).
58. Yang, J. *et al.* Potential geographical distribution of the edangered plant isoetes under human activities using maxent and garp. *Glob. Ecol. Conserv.* **38**, e2186 (2022).
59. Zhang, V. M., Punzalan, D. & Rowe, L. Climate change has different predicted effects on the range shifts of two hybridising ambush bug (Phymata) species. *Ecol. Evol.* **21**, 12036–12048 (2020).
60. Liu, L. *et al.* Modeling habitat suitability of *Houttuynia Cordata* thunb (Ceercao) using maxent under climate change in China. *Ecol. Inform.* **63**, 101324 (2021).
61. Wang, G. Z. *et al.* Predicting *Pseudolarix Amabilis* potential habitat based on four niche models. *Ecologica Sinica.* **40**, 6096–6104 (2020).

Author contributions

D. Z., J. C. and J. W. provided relevant data of *Fritillaria unibracteata* and required funds for the experiment. D. Z., W. D., K. Y. and J. W. planned and supervised the project. Q. L., H. L., B. Z. and X. Y. analyzed the data and performed simulations.

Funding

This work was funded by the China Agriculture Research System of MOF and MARA (CARS-21), the key project at central government level: the ability establishment of sustainable use for valuable chinese medicine resources(2060302), Sichuan Science and Technology Program (2022ZHXC0101 and R22ZYZF0012), the Tibetan Plateau Scientific Expedition and Research Program (STEP) (2019QZKK0608), the Natural Science Foundation of Sichuan Province (2022NSFSC0589) and the Heavy Rain and Drought-Flood Disasters in Plateau and Basin Key Laboratory of Sichuan Province (SCQXKJYJXZD202209).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-42988-0>.

Correspondence and requests for materials should be addressed to J.W.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023