



OPEN

Apigenin ameliorates doxorubicin prompted testicular damage: biochemical, spermatological and histological based study

Muhammad Umar Ijaz^{1✉}, Saba Yaqoob¹, Ali Hamza¹, Mehwish David², Tayyaba Afsar³, Fohad Mabood Husain⁴, Houda Amor⁵ & Suhail Razak^{3✉}

Doxorubicin (DOX) is a highly effective, commonly prescribed, potent anti-neoplastic drug that damages the testicular tissues and leads to infertility. Apigenin (APG) is an important flavonoid that shows diverse biological activities. The present research was designed to evaluate the alleviative role of APG against DOX-induced testicular damages in rats. Forty-eight adult male albino rats were randomly distributed into 4 groups, control, DOX administered (3 mgkg⁻¹), DOX + APG co-administered (3 mgkg⁻¹ of DOX; 15 mgkg⁻¹ of APG), and APG administered group (15 mgkg⁻¹). Results of the current study indicated that DOX treatment significantly reduced the activities of superoxide dismutase (SOD), glutathione reductase (GSR), catalase (CAT) and glutathione peroxidase (GPx), while increasing the levels of malondialdehyde (MDA) and reactive oxygen species (ROS). DOX treatment also reduced the sperm count, viability, and motility. Moreover, DOX significantly increased the sperm morphological anomalies and reduced the levels of plasma testosterone, luteinizing hormone (LH) and follicle-stimulating hormone (FSH). The administration of DOX significantly increased the expressions of Bax and Caspase-3, as well as the levels of inflammatory markers. Additionally, DOX treatment significantly downregulated the expressions of steroidogenic enzymes (StAR, 3β-HSD and 17β-HSD) and Bcl-2. Furthermore, DOX administration provoked significant histopathological abnormalities in the testicular tissues. However, APG supplementation significantly reversed all the testicular damages due to its androgenic, anti-apoptotic, anti-oxidant and anti-inflammatory nature. Therefore, it is concluded that APG may prove a promising therapeutic agent to treat DOX-induced testicular damages.

Keywords Doxorubicin, Apigenin, Oxidative stress, Testicular damage, Apoptosis

Chemotherapy is a commonly prescribed cancer treatment that refers to the use of chemical agents to destroy cancerous cells. It is considered to be highly effective in combating cancer¹. The long-term use of anti-tumor drugs can cause serious toxic effects that may damage the body, which ultimately restricts their clinical application². Unfortunately, the incidence of cancer has been reached to the point where it is considered to be the second leading cause of death worldwide, followed by heart disease³. Chemotherapeutic advancements have increased the life expectancy of cancer patients, 82% of cancer patients are currently enjoying the expectancy of life up to 5 years due to progressive advancements in chemotherapy⁴. Cancer is a serious public health challenge that causes profound morbidity and mortality globally⁵. DOX can induce hepatotoxicity, nephrotoxicity^{6,7} as well as testicular toxicity⁸.

According to the previous literature, DOX has the potential to induce oxidative stress (OS), lipid peroxidation (LP) and testicular damage⁹. The oxidative damage induced by DOX treatment is due to a combination of free radicals that contain superoxide, hydroxyl radicals and iron. These free radicals disrupt the mitochondrial

¹Department of Zoology, Wildlife and Fisheries, University of Agriculture, Faisalabad 38040, Pakistan. ²Department of Animal Sciences, Quaid-I-Azam University, Islamabad 45320, Pakistan. ³Department of Community Health Sciences, College of Applied Medical Sciences, King Saud University, P.O. Box 2455, 11451 Riyadh, Saudi Arabia. ⁴Department of Food Science and Nutrition, College of Food and Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia. ⁵Department of Obstetrics, Gynecology and Reproductive Medicine, Saarland University Clinic, Homburg, Germany. ✉email: umar.ijaz@uaf.edu.pk

membrane present in the sperm mid-piece and result in reduced ATP production¹⁰. DOX has the potential to reduce sperm quantity, motility and increase sperm abnormalities in rats¹¹.

Natural products are the primary source of the active ingredients of medicines. More than 80% medicines are derived from natural sources¹². Plant-based bioactive substances, such as flavonoids have received a lot of attention during the last few years due to their broad range of pharmacological effects, such as antioxidant, anti-apoptotic, and anti-inflammatory. Plants use flavonoids to protect themselves from biotic and abiotic stress and to defend against plaques¹³. They are considered as risk-free and show little adverse effects^{14,15}. Apigenin (APG) is an important naturally occurring flavonoid¹⁶ that is extracted from plant leaves, seeds, vegetables, fruits¹⁷ and several herbal medicines including *Teucrium gnaphalodes*, *Stachys tibetica* Vatke, and *Scutellaria baicalensis* Georgi¹⁸. APG shows anti-fungal, anti-oxidant, anti-mutagenic, anti-inflammatory¹⁹ and anti-cancer properties²⁰. To overcome DOX induced testicular toxicity, a detailed investigation on plant-based remedies is required. Therefore, the present study was designed to evaluate the ameliorative role of APG against DOX-induced testicular damage in rats.

Materials and methods

Chemicals

Doxorubicin (DOX) (Physical form: Liquid; CAS number: 25316–40-9; Molecular weight: 579.98) and apigenin (APG) (Physical form: Solid; CAS number: 578–74-5; Molecular weight: 432.38) were obtained from *Sigma Aldrich* (Germany).

Animals

In the current experiment, adult male albino rats (250 ± 20 g) were used. The rats were placed in steel cages maintained at standard temperature (22–25 °C), 12-h dark/light cycle and humidity (45 ± 5%) in the animal house of the University of Agriculture, Faisalabad (UAF). The animals were provided free access to commercial food and water and the protocol of the study was approved by UAF ethical committee of Animal Protection and Handling. The study procedures were performed by following the ARRIVE guidelines²¹ and all the methods were performed in accordance with the relevant guidelines and regulations.

Experimental protocol

Fourty eight rats were distributed into 4 groups, 12 rats per group. First group was designed as a control group. Second group was administered with DOX (3 mg/kg), Third group was co-administered with DOX (3 mg/kg) and APG (15 mg/kg) and the Forth group was administered with APG (15 mg/kg) only. The 3 doses of DOX were administered intraperitoneally after every 2 weeks, whereas the dose of APG was administered daily through oral gavage. The dose of DOX was selected according to the previous study²², while APG dose was selected in compliance with the study of Guo et al.²³ After the completion of 56 days of treatment, the rats were anesthetized using combined intraperitoneal injection of ketamine (75 mg/kg) and xylazine (2.5 mg/kg)²⁴. Anesthetized rats were secured in a supine position and a thoracotomy was performed to assure euthanasia. The euthanasia method was performed in accordance with AVMA guidelines²⁵. The blood was collected in sterile containers and serum samples were obtained by centrifuging blood at 1000 g for 15 min and stored at –20 °C for biochemical assays. Testes were removed, right testis was stored in zipper bags for biochemical observation at –80 °C, testicular tissues were homogenized in Na₃PO₄ buffer solution at 12,000 rpm for 14–15 min. Finally, various parameters were evaluated by using the supernatant. The right testis was fixed in formalin (10%) for histopathological analysis.

Assessment of catalase (CAT)

The activity of CAT in was determined in accordance with the technique of Chance and Maehly²⁶. 2.5 mL of 50 mM phosphate buffer (pH 5.0), 0.4 mL of 5.9 mM H₂O₂, and 0.1 mL enzyme extract were mixed to make reaction mixture. Absorbance changes in the mixture were observed at 240 nm. One unit of CAT activity was considered as an absorbance change of 0.01 as units/min.

Assessment of superoxide dismutase (SOD)

The activity of SOD was assessed by using the method of Kakkar et al.²⁷ Reaction mixture consisted of 1.2 mL of sodium pyrophosphate buffer (0.052 mM; pH 7.0) and 0.1 mL of phenazine methosulphate (186 mM). 0.3 mL of supernatant after centrifugation (1500 × g for 10 min followed by 10,000 × g for 15 min) of homogenate was added to the reaction solution. Then, 0.2 mL of NADH (780 mM) was added to initiate enzyme reaction, which was later stopped by adding 1 mL of glacial acetic acid. Finally, chromogen's amount was assessed by noticing the change in color intensity (at 560 nm). The values of SOD activity were presented as unit/mg protein.

Assessment of glutathione peroxidase (GPx)

The activity of GPx was evaluated according to the protocol of Rotruck et al.²⁸ Reaction mixture of GPx activity consisted of 0.01 mL of 10 mM sodium azide, 2.0 mL of 0.4 M Tris–HCl buffer (pH 7.0), 0.5 mL of 0.2 mM H₂O₂ and 0.2 mL of 10 mM glutathione. Incubation was performed at 37 °C for about 10 min. and then 0.4 mL 10% (v/v) TCA was added for termination of reaction. The mixture was centrifuged at 5000 rpm for about 5 min. and absorbance was noticed at 430 nm. Its final values were shown as unit/mg protein.

Assessment of glutathione reductase (GSR)

The activity of GSR was estimated according to the procedure of Carlberg and Mannervik²⁹. Reaction mixture consisted of 0.1 mL EDTA (0.5 mM), 0.05 mL oxidized glutathione (1 mM), 0.1 mL NADPH (0.1 mM), 1.65 mL

phosphate buffer (0.1 M, pH 7.6) and 0.1 mL of 10% homogenate in a volume of 2 mL. Enzymatic activity was measured at 25 °C by noticing NADPH disappearance at about 340 nm. The values obtained were presented as nM NADPH oxidized/min/mg tissue.

Assessment of MDA level

MDA level was measured by the method of Afsar et al.³⁰ The quantification of thiobarbituric acid reactive substances was assessed by comparing the absorption to the standard curve of MDA equivalents generated by acid-catalyzed hydrolysis of 1, 1, 3, 3 tetramethoxy propane. The values of MDA were displayed as nmol/mg protein.

Assessment of ROS level

ROS level was determined by 2', 7' -dichlorofluorescein diacetate (DCFH-DA). In brief, 100 µL cell lysates or renal homogenate were reacted with 100 µL 2 mg/mL DCFH-DA. A fluorescence plate reader was employed to monitor the fluorescence change after 30 min incubation at 37 °C. The excitation and emission wavelengths were respectively set at 488 nm and 525 nm. Result was displayed as relative fluorescence unit (RFU) per mg protein.

Semen analysis

The caudal portion of the epididymis was used to collect sperm samples. The caudal portion was minced and incubated in physiological saline for 30 min at 37 °C to release the spermatozoa. Then sperm motility was observed under a phase-contrast microscope (400X)³¹. Eosin or nigrosin staining and microscopic examination were used to evaluate the sperm viability. Moreover, epididymal sperm count was counted using a hemocytometer³². The sperm structural anomalies (head, tail and midpiece) were evaluated by using the Filler method³³.

Hypo-osmotic swelling test

HOS test was used to determine the sperm integrity by following the Correa and Zavos³⁴ protocol. The current experiment was performed by mixing 180 µL of fructose solution with 20 µL of semen sample by keeping the 80 mOsm/L osmotic pressure for twenty minutes. Following incubation, sperms were stained using eosin or nigrosin. Moreover, 200 sperm of swollen and non-swollen tails were calculated at 400X.

RNA extraction and real-time quantitative reverse transcription-polymerase chain reaction (qRT-PCR)

Steroidogenic enzymes and apoptotic markers expressions were evaluated by using RT-qPCR. The tissue samples were homogenized by employing a rotor stator homogenizer to isolate RNA. After centrifugation, the supernatant was collected and 70% ethanol was added. RNA spin columns were used for purification. DNase I treatment was accomplished, followed by multiple washes. Ethanol was removed and the isolated RNA was stored at -70 °C. RNA quality was determined with the help of NanoDrop Spectrophotometer, using RNA samples having A260/A280 ratio between 1.8 and 2.0. cDNA was synthesized from RNA samples using the Thermo Scientific Revert Aid RT Kit and reverse transcriptase enzyme. RT-qPCR was carried out in a LightCycler 480 instrument (Roche, Basel, Switzerland) using Syber green PCR master mix (TaKaRa, Biotech. Co., Ltd.). PCR amplification was started by denaturation for 3 min at 95 °C that was followed by 45 cycles of 95 °C for 10 s, 60 °C for 40 s, 72 °C for 32 s and a final incubation at 75 °C for 5 min. β -actin was considered as internal control and $2^{-\Delta\Delta CT}$ was used to assess the variations in the expressions of these parameters. Table 1 displays the primer sequences of the target genes as reported previously³⁵.

Gene	Primers 5'–3'	Accession number	Product size
3 β -HSD	Forward: GCCACCCCTTAACTGCCACT	NM_0010079	119
	Reverse: CTGTGCTGCTCCACTAGTGT		
17 β -HSD	Forward: TATCCAGGTGCTGACCCCTT	NM_054007	103
	Reverse: CAAGGCAGCCACAGGTTCA		
StAR	Forward: AGCGTAGAGGTTCCACCTGT	NM_031558	123
	Reverse: ATACTGAGCAGCCACGTGAG		
Bax	Forward: GCACTAAAGTGCCCGAGCTG	NM_017059.2	148
	Reverse: CCAGATGGTGAGTGAGGCAG		
Bcl-2	Forward: ACTGAGTACCTGAACCGGCA	NM_016993.1	139
	Reverse: CCCAGGTATGCACCCAGAGT		
Caspase-3	Forward: GTACAGAGCTGGACTGCGGT	NM_012922.2	137
	Reverse: TCAGCATGGCGCAAAGTGAC		
β -actin	Forward: AGGAGATTACTGCCCTGGCT	NM_031144	138
	Reverse: CATTGCGGTGCACGATGGA		

Table 1. Primers sequences for the real-time quantitative reverse transcription-polymerase chain reaction. 3 β -hydroxysteroid dehydrogenase (3 β -HSD); 17 β -hydroxysteroid dehydrogenase (17 β -HSD); Steroidogenic acute regulatory protein (StAR).

Hormonal analysis

Plasma testosterone (serial number H090), LH (serial number H206) and FSH (serial number H101) levels were measured using (ELISA) kits by following the company's directions (Los Angeles, CA, USA). The 96-well ELISA plate was filled with assay diluent (50 μ L) as well as plasma (10 μ L). It was then incubated at room temperature for two hours. Following a rinse with deionized water, the plates were incubated for a maximum of two hours before 100 μ L of peroxidase-conjugated immunoglobulin G (IgG) anti-LH, anti-FSH or anti-testosterone solution was added to each well. Plates were once again washed with deionized water, then wells were filled with substrate solution and incubated for 25 min at room temperature. To halt the reaction, stop solution (50 μ L) was added. Lastly, at 450 nm, the absorbance of plasma testosterone, LH, and FSH was measured. To prevent inter-assay variance, each sample was conducted in triplicate under the same circumstances at the same time³⁶.

Inflammatory indices

The levels of NF- κ B (CSB-E13148r), TNF- α (CSB-E07379r), IL-1 β (CSB-E08055r) and IL-6 (CSB-E04640r) and COX-2 (CSB-E13399r) activity were evaluated by ELISA kits (Cusabio Technology Llc, Houston, TX, USA) and the instructor's guidelines were followed. Firstly, 50 μ L of sample was dispensed to the microplate wells. After that, 50 μ L of antibody cocktail was poured to the wells. Plates were incubated at room temperature for the duration of 1 h. After washing properly with the help of wash buffer, 100 μ L of TMB substrates were dispensed to each well and incubated for about 10 min. After the addition of 100 μ L of stop solution, the color was developed. The optical density was noted at 450 nm using Tecan Multimode Reader³⁶.

Histopathology

The tissues were kept in 10% formalin after cleaning with a normal saline solution. The tissues were gradually dehydrated by using the rising grades of alcohol 70%, 90% and 100% and then fixed in paraffin wax. 4–5- μ m tissue slices were cut with the help of a microtome and stained with hematoxylin–eosin. Then, these slides were examined under a light microscope (400X) and microphotography was carried out using a Leica LB microscope connected to a camera. The photos were analyzed by using Image J2X software.

Statistical analysis

Data were presented as mean \pm SEM. One-way ANOVA followed by Tukey's test was performed by using Minitab software. The level of significance was set at $P < 0.05$.

Ethical approval

The protocol of the study was approved by UAF ethical committee of Animal Protection and Handling. The study procedures were performed by following the ARRIVE guidelines and all the methods were performed in accordance with the relevant guidelines and regulations.

Results

Effect of DOX and APG on biochemical parameters

In the current study DOX administration induced a significant ($P < 0.001$) decrease in the activities of GPx, GSR, CAT and SOD. On the other hand, a significant ($P < 0.001$) increase in the levels of MDA and ROS was observed as compared to control group. However, the co-administration of APG + DOX significantly ($P < 0.01$) increased the activities of GPx, GSR, CAT, and SOD. Moreover, MDA and ROS levels were significantly ($P < 0.01$) reduced in co-administered group as compared to DOX-administered group. Moreover, in APG only administered group these parameters were comparable to the control group (Table 2).

Effect of DOX and APG on spermatogenic parameters

DOX-intoxication prompted a significant ($P < 0.01$) reduction in the sperm viability, count and motility, whereas a significant ($P < 0.01$) increase in the dead sperm count and sperm structural anomalies (tail, head and mid-piece) were observed as compared to control. However, APG + DOX co-administration significantly ($P < 0.01$) reduced the sperm abnormalities induced by DOX and significantly ($P < 0.01$) increased the sperm motility,

Parameters	Groups			
	Control	DOX	DOX + APG	APG
CAT (Umg ⁻¹ protein)	9.35 \pm 0.29	4.94 \pm 0.12 ^{###}	7.53 \pm 0.58 ^{**}	9.38 \pm 0.30 ^{***}
SOD (Umg ⁻¹ protein)	7.02 \pm 0.17	3.35 \pm 0.14 ^{###}	5.28 \pm 0.18 ^{**}	7.07 \pm 0.12 ^{***}
GPx (Umg ⁻¹ protein)	23.73 \pm 0.99	7.19 \pm 0.21 ^{###}	18.42 \pm 0.73 ^{**}	23.79 \pm 0.98 ^{***}
GSR (nM NADPH oxidized/min/mg tissue)	6.31 \pm 0.28	1.48 \pm 0.14 ^{###}	4.77 \pm 0.10 ^{**}	6.35 \pm 0.29 ^{***}
ROS (RFU/mg protein)	0.72 \pm 0.08	8.12 \pm 0.28 ^{###}	2.54 \pm 0.24 ^{**}	0.68 \pm 0.09 ^{***}
MDA (nmol/mg protein)	0.69 \pm 0.14	6.06 \pm 0.17 ^{###}	1.51 \pm 0.13 ^{**}	0.66 \pm 0.07 ^{***}

Table 2. Effect of DOX and APG on biochemical parameters. Values based on mean \pm SEM values. Significant differences are displayed as ^{###} $P < 0.001$ compared to control group; ^{**} $P < 0.01$ compared to DOX-administered group, ^{***} $P < 0.001$ compared to control group.

viability and sperm number in comparison to DOX administered group. Moreover, these sperm indices in APG alone administered group were close to the control group (Table 3).

Effect of DOX and APG on steroidogenic enzymes

DOX-intoxication led to a significant ($P < 0.01$) decrease in the expressions of steroidogenic enzymes (17 β -HSD, 3 β -HSD and StAR) in contrast to the control group. Nonetheless, the co-administration of APG + DOX significantly ($P < 0.05$) increased these expressions in contrast to DOX-administered group. Moreover, the expressions of these steroidogenic enzymes in only APG administered group were close to the control group (Fig. 1).

Parameters	Groups			
	Control	DOX	DOX + APG	APG
Epididymal sperm count (million/mL)	25.68 \pm 1.21	7.33 \pm 0.53 ^{##}	18.36 \pm 0.77 ^{**}	25.93 \pm 1.33 ^{***}
Motility (%)	81.97 \pm 1.22	33.84 \pm 0.93 ^{##}	72.37 \pm 1.01 ^{**}	82.56 \pm 1.66 ^{***}
Dead sperms (%)	15.92 \pm 0.93	75.78 \pm 0.87 ^{##}	34.24 \pm 1.19 ^{**}	14.89 \pm 1.03 ^{***}
Head abnormality (%)	2.74 \pm 0.37	33.38 \pm 1.20 ^{##}	7.41 \pm 0.73 ^{**}	2.69 \pm 0.38 ^{***}
Mid sperm abnormality (%)	0.58 \pm 0.08	13.37 \pm 1.27 ^{##}	2.63 \pm 0.19 ^{**}	0.55 \pm 0.08 ^{***}
Tail abnormality (%)	4.71 \pm 0.53	23.12 \pm 1.89 ^{##}	9.88 \pm 0.45 ^{**}	4.69 \pm 0.52 ^{***}
Hypo-osmotic swelled sperm count (HOS) (%)	87.92 \pm 2.28	25.35 \pm 1.15 ^{##}	71.63 \pm 0.83 ^{**}	89.08 \pm 2.38 ^{***}

Table 3. Effect of DOX and APG on sperm parameters. Values based on mean \pm SEM values. Significant differences are displayed as ^{##} $P < 0.01$ compared to control group; ^{**} $P < 0.01$ compared to DOX-administered group, ^{***} $P < 0.001$ compared to control group.




Figure 1. Effect of DOX and APG on the expression of (a) 3 β -HSD, (b) 17 β -HSD and (c) StAR. The bars are based on mean \pm SEM values. Significant differences are displayed as ^{##} $P < 0.01$ compared to control group; ^{*} $P < 0.05$ compared to DOX-administered group, ^{**} $P < 0.01$ compared to control group.

Effect of DOX and APG on hormones

DOX exposure significantly ($P < 0.001$) decreased the levels of LH, FSH and testosterone in contrast to the control group. However, co-administration of APG + DOX resulted in a considerable ($P < 0.01$) increase in the hormonal level in contrast to DOX-administered group. Moreover, in only APG administered group the hormonal level was close to the control group (Fig. 2).

Effect of DOX and APG on apoptotic markers

DOX-intoxication resulted in a significant ($P < 0.01$) reduction in the expression of Bcl-2, while a significant ($P < 0.01$) upsurge in the expressions of Caspase-3 and Bax as compared to control group. However, the co-administration of APG + DOX considerably ($P < 0.05$) decreased Caspase-3 and Bax expressions, while significantly ($P < 0.05$) increasing the Bcl-2 expression as compared to DOX-administered group. Additionally, APG only administered group showed the expressions of these markers close to the control group (Fig. 3).

Effect of DOX and APG on inflammatory indices

DOX administration significantly ($P < 0.001$) increased the inflammatory marker level such as, TNF- α , IL-6, NF- κ B, IL-1 β and COX-2 activity in comparison to control animals. However, the co-administration of APG with DOX significantly ($P < 0.01$) decreased the level of inflammatory indices as compared to DOX-administered group. Additionally, the level of inflammatory indices in APG alone administered group were similar to the control group (Fig. 4).




Figure 2. Effect of DOX and APG on the level of (a) FSH, (b) LH and (c) Plasma testosterone. The bars are based on mean \pm SEM values. Significant differences are displayed as ### $P < 0.001$ compared to control group; ** $P < 0.01$ compared to DOX-administered group, *** $P < 0.001$ compared to control group.




Figure 3. Effect of DOX and APG on the expression of (a) Bax, (b) Bcl-2 and (c) Caspase-3. The bars are based on mean \pm SEM values. Significant differences are displayed as ## $P < 0.01$ compared to control group; * $P < 0.05$ compared to DOX-administered group, ** $P < 0.01$ compared to control group.

Effect of DOX and APG on testicular histopathology

DOX-intoxication induced a significant ($P < 0.001$) decrease in seminiferous tubular diameter as well as height and width of tunica propria. While a significant ($P < 0.001$) increase in the interstitial space and luminal diameter was observed as compared to the control group. Additionally, DOX treatment caused a significant ($P < 0.001$) reduction in the germ cells count. However, the co-administration of APG with DOX resulted in significant ($P < 0.01$) increase in seminiferous tubular diameter as well as height and width of tunica propria. Conversely, a significant ($P < 0.01$) decrease in the luminal diameter and interstitial spaces was observed as compared to DOX administered group. Furthermore, histopathological profile in only APG administered group was similar to the control group (Table 4, Fig. 5).

Discussion

DOX is a powerful, commonly prescribed and a strong anti-cancer drug. However, its clinical application is limited due to its potential side effects such as, cardiotoxicity³⁷, hepatotoxicity³⁸, nephrotoxicity³⁹ and testicular toxicities⁴⁰. DOX intoxication is reported to disturb hormone levels, sperm parameters, apoptotic markers, antioxidant enzymes, steroidogenic enzymes and testicular histopathology⁴¹. DOX exposure may damage the quantity and quality of sperm in adults⁴². It is reported that natural anti-oxidant can be used to treat OS-related disorders^{43,44}. Apigenin (APG) is a natural flavonoid that is present in the leaves, seeds and fruits of *Teucrium gnaphalodes*, *Stachys tibetica* Vatke, and *Scutellaria baicalensis* Georgi⁴⁵. APG shows anti-hyperglycemic, anti-oxidant, anti-carcinogenic, anti-inflammatory and anti-viral effects⁴⁶. Therefore, the present research was planned to assess the therapeutic role of APG in mitigating DOX-induced testicular damage in rats.

DOX exposure significantly reduced the activities of antioxidant enzymes (CAT, GSR, SOD and GPx), while increasing the levels of ROS and MDA. ROS includes superoxide radicals (O_2^-), hydroxyl radicals (OH^-) and hydrogen peroxide (H_2O_2), which are metabolic byproducts that may cause harm to cells of the body. Our results are further supported by the previous study in which it was reported that DOX showed the potential to induce OS in rats and reduced the activities of anti-oxidant enzymes⁴⁷. The production of ROS leads to OS that ultimately damage the quality and quantity of sperms⁴⁸. Elevated ROS production impairs spermatozoa's




Figure 4. Effect of DOX and APG on the level of (a) NF-κB, (b) TNF-α, (c) IL-1β, (d) IL-6 and (e) COX-2. The bars are based on mean ± SEM values. Significant differences are displayed as ###*P* < 0.001 compared to control group; ***P* < 0.01 compared to DOX-administered group, ****P* < 0.001 compared to control group.

anti-oxidant defense mechanism⁴⁹. CAT is one of the major antioxidant enzyme that plays a substantial part in H₂O₂ catabolism⁵⁰. It separates NADPH oxidase-generated superoxide radicals and safeguards the spermatozoa from oxidative damage⁵¹. SOD transforms superoxide into H₂O₂ and O₂⁵². GSR facilitates the transformation of GSH from its reduced state to its oxidized state⁵³. GPx helps in lowering the lipid peroxide (LP) and H₂O₂ concentrations and plays an important role in reducing OS⁵⁴. CAT, GPx, SOD and GST are anti-oxidant enzymes that are pivotal in alleviating ROS, OS and eventually reduce the damage⁵⁵. MDA is a byproduct of ROS-induced LP and it is used as a biomarker of LP and OS⁵⁶. In order to minimize OS, plant sources can also be used as a supplement⁵⁷. The co-administration of APG with DOX lowered the OS in rat testes and increased the anti-oxidant enzymes activities, besides lessened the level of ROS and MDA due to its anti-oxidant potential. Moreover, our results are

Parameters	Groups			
	Control	DOX	DOX + APG	APG
Interstitial spaces (μm)	6.55 \pm 0.68	73.90 \pm 1.85 ^{***}	19.48 \pm 1.49 ^{b**}	6.52 \pm 0.67 ^{***}
Tunica propria (μm)	85.63 \pm 1.19	15.40 \pm 1.23 ^{***}	64.34 \pm 1.52 ^{**}	86.53 \pm 1.62 ^{***}
Seminiferous tubules diameter (μm)	372.44 \pm 5.17	138.41 \pm 5.12 ^{***}	241.06 \pm 6.08 ^{**}	375.21 \pm 6.09 ^{***}
Seminiferous tubule epithelial height (μm)	74.44 \pm 1.97	28.46 \pm 1.81 ^{***}	73.57 \pm 1.84 ^{**}	73.39 \pm 2.59 ^{***}
Tubular lumen (μm)	32.68 \pm 0.97	97.51 \pm 3.06 ^{***}	57.14 \pm 1.57 ^{**}	32.17 \pm 0.33 ^{***}
Spermatogonia (n)	65.59 \pm 1.32	24.80 \pm 0.76 ^{***}	54.35 \pm 1.05 ^{**}	66.33 \pm 1.15 ^{***}
Primary spermatocytes (n)	55.99 \pm 1.83	19.56 \pm 1.62 ^{***}	44.22 \pm 1.87 ^{**}	58.61 \pm 2.74 ^{***}
Secondary spermatocytes (n)	46.77 \pm 2.55	15.74 \pm 1.58 ^{***}	34.78 \pm 1.74 ^{**}	48.64 \pm 3.18 ^{***}
Spermatids (n)	63.73 \pm 0.90	24.95 \pm 1.84 ^{***}	37.3 \pm 15.85 ^{**}	64.55 \pm 1.22 ^{***}

Table 4. Effect of DOX and APG on testicular histopathology. Values based on mean \pm SEM values. Significant differences are displayed as ^{***} $P < 0.001$ compared to control group; ^{**} $P < 0.01$ compared to DOX-administered group, ^{***} $P < 0.001$ compared to control group.




Figure 5. Micro-images of the adult albino rat testicles (H&E, 400X): (A) Control group presenting impenetrable germinal epithelium exhibiting germ cells & tapered luminal area with sperms; (B) APG treated group showing compacted seminiferous tubules with less IS and luminal part filled with germ cells; (C) DOX exposed group showing sloughing of the epithelial layer, vacant lumen & degenerated area of IS; (D) DOX + APG co-administered group demonstrating renovation in epithelial part as well as TL filled with ST and recovered deteriorated IS. DOX: Doxorubicin; APG: Apigenin TL: Tubular lumen; EH: Seminiferous epithelial height; IS: Interstitial spaces; SG: Spermatogonia; PS: Primary spermatocytes; SS: Secondary Spermatocytes; ST: Spermatids.

further endorsed by the study of Kale et al.⁵⁸ who stated that double bonds between C2 and C3 in the structure of APG are responsible for its anti-oxidant activity.

DOX intoxication significantly reduced the sperm viability, number and motility. Besides dead sperm count and sperm tail, head and mid-piece abnormalities were increased in the DOX exposed rats. Our findings are in line with the research of Ahmad et al.⁵⁹ who stated that DOX administration resulted in reduced sperm count, impaired motility, induced spermatotoxicity, and spermatogenic failure. Sperm quality is a crucial factor in ascertaining male reproductive capability, where the primary indicators are sperm count and mortality rate. Reduced sperm count can be due to the delayed maturation and abnormal germ cell proliferation⁶⁰. According to previous literature, large amount of polyunsaturated fatty acids (PUFAs) make spermatozoa susceptible to OS. The motility of sperm and the integrity of sperm plasma membrane depend on the PUFAs⁶¹. The fluidity of sperm plasma membrane and the process of spermatogenesis are severely affected by OS⁶². Furthermore, OS can directly affect sperm cell mitochondria and results in reduced ATP production, flagellar activity and eventually leads to apoptotic death and immobility⁶³. However, the supplementation of APG mitigated all these DOX prompted impairments due to its ROS scavenging nature.

DOX exposure significantly reduced the expressions of 3β -HSD, 17β -HSD and StAR. These enzymes play a pivotal role in the production of steroid hormones⁶⁴. StAR acts as a regulatory protein during the production of testosterone and it plays a key role in the transportation of cholesterol inside the mitochondrial membrane^{65,66}. 17β -HSD and 3β -HSD catalyzes the conversion of cholesterol to testosterone⁶⁷, while lowered expressions of these enzymes, decrease the level of testosterone⁶⁸. Nevertheless, the supplementation of APG significantly increased the steroidogenic enzymes expression and restored the level of testosterone. Moreover, it is stated that flavonoids have chemical structure similar to cholesterol that could regulate the production of androgens⁶⁹.

In the current investigation, DOX intoxication significantly decreased the level of plasma testosterone, FSH and LH. According to the study conducted by Mustafa et al.²² DOX possesses the potential to reduce the hormonal level in rats. Spermatogenesis is controlled by a proper ratio of testosterone, FSH and LH⁷⁰. The production of testosterone is crucial for the reproductive health of male⁷¹. Both FSH and LH are vital to control spermatogenesis⁷². LH induces LCs to generate testosterone, besides FSH encourages Sertoli cell proliferation⁷³. However, the supplementation of APG improved the levels of these hormones and eventually recovered the spermatogenesis. Our findings are supported by the research of Agrawal⁷⁴ who stated that flavonoids have the potential to control the production of hormones i.e., estrogen and androgen.

DOX exposure augmented the expressions of Bax and Caspase-3, while reducing Bcl-2 expression. Bcl-2 acts as anti-apoptotic protein that safeguards the cells from apoptotic death, Bax accelerates the process of apoptosis⁷⁵. A reduction in Bcl-2 and elevation in Bax expression induce significant alteration in mitochondrial membrane permeability, which eventually stimulates the liberation of cytochrome C into the cytoplasm⁷⁶. Caspase-3 belongs to the cysteine aspartic protease family and plays a pivotal role in initiating and executing cellular apoptosis⁷⁷. Caspase-3 is activated by the augmented level of cytochrome C that leads to cell death or apoptosis⁷⁸. However, the administration of APG augmented the Bcl-2 expression, whereas lowered the expression of Caspase-3 and Bax due to its anti-apoptotic nature.

DOX administration significantly increased the levels of inflammatory markers, such as TNF- α , IL-1 β , NF- κ B, IL-6 and COX-2 activity. The results of our research are supported by the research of Hu et al.⁷⁹ who stated that DOX exposure augmented the level of inflammatory markers in the testicular tissue of rats. Inflammation is one of the major cause of male infertility⁸⁰. NF- κ B instigation is pivotal for the expression of proinflammatory cytokines (IL-6, TNF- α , COX-2 and IL-1 β) that are related with severe inflammation and other ROS associated disorders⁸¹. COX-2 is also an inflammatory marker that remarkably contributes to inflammation⁸². However, the administration of APG with DOX resulted in a considerable reduction in inflammatory indices due to its anti-inflammatory nature. Moreover, our results are further supported by the previous study in which it was reported that APG has the potential to attenuate lipopolysaccharide induced inflammation in rats²³.

DOX exposure significantly increased the diameter of tubular lumen and interstitial spaces. Besides, it reduced the seminiferous tubular diameter and epithelial height, tunica propria width and germ cells count. The findings of our current histopathological investigation are in line with an earlier study, in which DOX-intoxication reduced seminiferous tubular diameter as well as germ cells number. It was also reported that DOX-intoxication reduced the testosterone level, which ultimately reduced the germ cells count and induced histopathological damages⁸³. According to Adejuwon et al.⁸⁴ an imbalance between pro-oxidants and anti-oxidants results in the overproduction of ROS that leads to histopathological damage. The disruption of steroidogenic enzymes expression as well as a decrease in testosterone level is one of the major reasons of spermatogenesis failure⁸⁵. However, APG treatment considerably augmented the germ cells count and restored all the testicular anomalies due to its anti-apoptotic, anti-oxidative, anti-inflammatory and androgenic properties.

Conclusion

The findings of the current study indicated that APG exhibited excellent ameliorative role against DOX-induced testicular damage by reducing the oxidative stress. APG administration significantly regulated the activities of anti-oxidant enzymes, the levels of MDA ROS and inflammatory indices, the expressions of steroidogenic enzymes, apoptotic and anti-apoptotic proteins as well as histological profile of testes. Therefore, our findings revealed that APG may be used to cure testicular damage. However, this study was performed by using rats as animal model so, its clinical trials on humans are recommended in future.

Data availability

All the data is contained in the manuscript.

References

- Punia, R., Raina, K., Agarwal, R. & Singh, R. P. Acacetin enhances the therapeutic efficacy of doxorubicin in non-smallcell lung carcinoma cells. *PLoS ONE* **12**, 1–19. <https://doi.org/10.1371/journal.pone.0182870> (2017).
- Cao, T. *et al.* Rare earth fluorescent nanoprobes with minimal side effects enable tumor microenvironment activation for chemotherapy. *J. Rare Earths*. <https://doi.org/10.1016/j.jre.2022.11.005> (2022).
- Rhee, P. *et al.* Increasing trauma deaths in the United States. *Ann. Surg.* **260**, 13–21. <https://doi.org/10.1097/SLA.0000000000000600> (2014).
- Gatta, G. *et al.* Childhood cancer survival in Europe 1999–2007: results of EURO CARE-5—a population-based study. *Lancet Oncol.* **15**, 35–47. [https://doi.org/10.1016/S1470-2045\(13\)70548-5](https://doi.org/10.1016/S1470-2045(13)70548-5) (2014).
- Mirzaei, S. *et al.* Resveratrol augments doxorubicin and cisplatin chemotherapy: a novel therapeutic strategy. *Curr. Mol. Pharmacol.* **16**, 280–306. <https://doi.org/10.2174/1874467215666220415131344> (2022).
- Song, S. *et al.* Protective effects of dioscin against doxorubicin-induced hepatotoxicity via regulation of Sirt1/FOXO1/NF- κ B signal. *Front. Pharmacol.* **10**, 1030. <https://doi.org/10.3389/fphar.2019.01030> (2019).
- Afsar, T., Razak, S., Almajwal, A. & Al-Disi, D. Doxorubicin-induced alterations in kidney functioning, oxidative stress, DNA damage, and renal tissue morphology; Improvement by Acacia hydaspicca tannin-rich ethyl acetate fraction. *Saudi J. Biol. Sci.* **27**, 2251–2260. <https://doi.org/10.1016/j.sjbs.2020.07.011> (2020).
- Aksu, E. H. *et al.* Palliative effect of curcumin on doxorubicin-induced testicular damage in male rats. *J. Biochem. Mol. Toxicol.* **33**, 22384. <https://doi.org/10.1002/jbt.22384> (2019).
- Öztürk, E. *et al.* Thymoquinone is a protective agent that reduces the negative effects of doxorubicin in rat testis. *Hum. Exp. Toxicol.* **39**, 1364–1373. <https://doi.org/10.1177/09603271209241> (2020).
- Türedi, S., Yuluğ, E., Alver, A., Kutlu, Ö. & Kahraman, C. Effects of resveratrol on doxorubicin induced testicular damage in rats. *Exp. Toxicol. Pathol.* **67**, 229–235. <https://doi.org/10.1016/j.etp.2014.12.002> (2015).
- Xin, Y. F. *et al.* Protective effect of Lycium barbarum polysaccharides against doxorubicin-induced testicular toxicity in rats. *Phytother. Res.* **26**, 716–721. <https://doi.org/10.1002/ptr.3633> (2012).
- Park, E. S. *et al.* Cardioprotective effects of rhamnetin in H9c2 cardiomyoblast cells under H₂O₂-induced apoptosis. *J. Ethnopharmacol.* **153**, 552–560 (2014).
- Hamza, A., Ijaz, M. U. & Anwar, H. Rhamnetin alleviates polystyrene microplastics-induced testicular damage by restoring biochemical, steroidogenic, hormonal, apoptotic, inflammatory, spermatogenic and histological profile in male albino rats. *Hum. Exp. Toxicol.* **42**, 9603271231173378. <https://doi.org/10.1177/09603271231173378> (2023).
- Jučá, M. M. *et al.* Flavonoids: Biological activities and therapeutic potential. *Nat. Prod. Res.* **34**, 692–705. <https://doi.org/10.1080/14786419.2018.1493588> (2020).
- Baskar, V., Venkatesh, R., Ramalingam, S. Flavonoids (antioxidants systems) in higher plants and their response to stresses. *Antioxidants Antioxidant Enzym Higher Plants* 2018:253–268
- Sun, Q., Lu, N. N. & Feng, L. Apigenin inhibits gastric cancer progression through inducing apoptosis and regulating ROS-modulated STAT3/JAK2 pathway. *Biochem. Biophys. Res. Commun.* **498**, 164–170. <https://doi.org/10.1016/j.bbrc.2018.02.009> (2018).
- Funakoshi-Tago, M., Nakamura, K., Tago, K., Mashino, T. & Kasahara, T. Anti-inflammatory activity of structurally related flavonoids, Apigenin Luteolin and Fisetin. *Int. Immunopharmacol.* **11**, 1150–1159. <https://doi.org/10.1016/j.intimp.2011.03.012> (2011).
- Lim, H. S., Kim, O. S., Kim, B. Y. & Jeong, S. J. Apigenin from scutellaria baicalensis georgi inhibits neuroinflammation in BV-2 microglia and exerts neuroprotective effect in HT22 hippocampal cells. *J. Med. Food.* **19**, 1032–1040. <https://doi.org/10.1089/jmf.2016.0074> (2016).
- Pandey, K. B. & Rizvi, S. I. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid. Med. Cell Long.* **2**, 270–278. <https://doi.org/10.4161/oxim.2.5.9498> (2009).
- Rodríguez-Mateos, A. *et al.* Bioavailability, bioactivity and impact on health of dietary flavonoids and related compounds: An update. *Arch. Toxicol.* **88**, 1803–1853. <https://doi.org/10.1007/s00204-014-1330-7> (2014).
- PercieduSert, N. *et al.* The ARRIVE guidelines 20: Updated guidelines for reporting animal research. *J. Cereb. Blood Flow Metab.* **40**, 1769–1777 (2020).
- Mustafa, S. *et al.* Isorhamnetin: a flavonoid, attenuated doxorubicin-induced testicular injury via regulation of steroidogenic enzymes and apoptotic signaling gene expression in male rats. *Toxicol. Res. (Camb)*. **23**, 475–485. <https://doi.org/10.1093/toxres/ffac024> (2022).
- Guo, H., Li, M. & Xu, L. J. Apigenin treatment attenuates LPS-induced acute otitis media through suppressing inflammation and oxidative stress. *Biomed. Pharmacother.* **109**, 1978–1987. <https://doi.org/10.1016/j.biopha.2018.07.022> (2019).
- Afsar, T. *et al.* Prevention of testicular damage by indole derivative MMINA via upregulated StAR and CatSper channels with coincident suppression of oxidative stress and inflammation. *In Silico In Vivo Valid.* **11**(10), 2063 (2022).
- Underwood W, Anthony RJROM: AVMA guidelines for the euthanasia of animals: 2020 edition. 2020, 2013(30):2020–2021.
- Maehly, A. C. & Chance, B. The assay of catalases and peroxidases. *Methods Biochem. Anal.* **1**, 357–424. <https://doi.org/10.1002/9780470110171> (1954).
- Kakkar, P., Das, B. & Viswanathan, P. N. A modified spectrophotometric assay of superoxide dismutase. *Indian J. Biochem. Biophys.* **21**, 130–132 (1984).
- Rotruck, J. T. *et al.* Selenium: biochemical role as a component of glutathione peroxidase. *Science*. **179**, 588–590. <https://doi.org/10.1126/science.179.4073.588> (1973).
- Carlberg, I. & Mannervik, B. Purification and characterization of the flavoenzyme glutathione reductase from rat liver. *J. Biol. Chem.* **250**, 5475–5480. [https://doi.org/10.1016/S0021-9258\(19\)41206-4](https://doi.org/10.1016/S0021-9258(19)41206-4) (1975).
- Afsar, T. *et al.* Prevention of testicular damage by indole derivative MMINA via upregulated StAR and CatSper channels with coincident suppression of oxidative stress and inflammation: In silico and in vivo validation. *Antioxidants (Basel)*. **11**, 2063. <https://doi.org/10.3390/antiox11102063> (2022).
- Kenjale, R., Shah, R. & Sathaye, S. Effects of Chlorophytum borivilianum on sexual behaviour and sperm count in male rats. *Phytother. Res.* **22**, 796–801. <https://doi.org/10.1002/ptr.2369> (2008).
- Yokoi, K., Uthus, E. O. & Nielsen, F. H. Nickel deficiency diminishes sperm quantity and movement in rats. *Biol. Trace Elem. Res.* **93**, 141–153. <https://doi.org/10.1385/BTER:93:1-3:141> (2003).
- Filler, R. Methods for evaluation of rat epididymal sperm morphology. *Toxicol. Mech. Methods*. **3**, 334–343 (1993).
- Correa, J. R. & Zavos, P. M. The hypoosmotic swelling test: its employment as an assay to evaluate the functional integrity of the frozen-thawed bovine sperm membrane. *Theriogenology*. **42**, 351–360. [https://doi.org/10.1016/0093-691X\(94\)90280-1](https://doi.org/10.1016/0093-691X(94)90280-1) (1994).
- Ijaz, M. U. *et al.* Chemoprotective effect of vitexin against cisplatin-induced biochemical, spermatological, steroidogenic, hormonal, apoptotic and histopathological damages in the testes of Sprague-Dawley rats. *Saudi Pharm. J.* **30**(5), 519–526. <https://doi.org/10.1016/j.jsps.2022.03.001> (2022).

36. Rizwan, A., Ijaz, M. U., Hamza, A. & Anwar, H. Attenuative effect of astilbin on polystyrene microplastics induced testicular damage: Biochemical, spermatological and histopathological-based evidences. *Toxicol. Appl. Pharmacol.* **471**, 116559. <https://doi.org/10.1016/j.taap.2023.116559> (2023).
37. Manechote, C., Chattipakorn, S. C. & Chattipakorn, N. Recent advances in mitochondrial fission/fusion-targeted therapy in doxorubicin-induced cardiotoxicity. *Pharmaceutics*. **15**, 1182. <https://doi.org/10.3390/pharmaceutics15041182> (2023).
38. Scarlata, E. & O'Flaherty, C. Antioxidant enzymes and male fertility: Lessons from knockout models. *Antioxid. Redox Signal.* **32**, 569–580. <https://doi.org/10.1089/ars.2019.7985> (2020).
39. Antar, S. A. *et al.* Modulatory role of autophagy in metformin therapeutic activity toward doxorubicin-induced nephrotoxicity. *Toxics*. **11**, 273. <https://doi.org/10.3390/toxics11030273> (2023).
40. Baokbah, T. A. Protective effects of selenium nanoparticles against doxorubicin-induced testicular apoptosis in rats. *Med sci.* **27**, e188ms2975 (2023).
41. Pugazhendhi, A., Edison, T. N., Karuppusamy, I. & Kathirvel, B. Inorganic nanoparticles: a potential cancer therapy for human welfare. *Int. J. Pharm.* **539**, 104–111. <https://doi.org/10.1016/j.ijpharm.2018.01.034> (2018).
42. Das, J., Ghosh, J., Manna, P. & Sil, P. C. Taurine protects rat testes against doxorubicin-induced oxidative stress as well as p53, Fas and caspase 12-mediated apoptosis. *Amino acids*. **42**, 1839–1855. <https://doi.org/10.1007/s00726-011-0904-4> (2012).
43. Hamza, A., Ijaz, M. U. & Anwar, H. Rhamnetin alleviates polystyrene microplastics-induced testicular damage by restoring biochemical, steroidogenic, hormonal, apoptotic, inflammatory, spermatogenic and histological profile in male albino rats. *Hum. Exp. Toxicol.* **42**, 9603271231173378. <https://doi.org/10.1177/09603271231173378> (2023).
44. Ijaz, M. U. *et al.* Pinostrobin alleviates testicular and spermatological damage induced by polystyrene microplastics in adult albino rats. *Biomed. Pharmacother.* **162**, 114686. <https://doi.org/10.1016/j.biopha.2023.114686> (2023).
45. Zhang, R. *et al.* Apigenin ameliorates streptozotocin-induced pancreatic β -cell damages via attenuating endoplasmic reticulum stress. *In Vitro Cell Dev. Biol.* **56**, 622–634. <https://doi.org/10.1007/s11626-020-00478-x> (2020).
46. Bhosale, P. B. *et al.* Apigenin promotes TNF α -induced apoptosis, necroptosis, G2/M phase cell cycle arrest, and ROS generation through inhibition of NF- κ B pathway in Hep3B liver cancer cells. *Cells*. **11**, 2734. <https://doi.org/10.3390/cells11172734> (2022).
47. Shi, S., Chen, Y., Luo, Z., Nie, G. & Dai, Y. Role of oxidative stress and inflammation-related signaling pathways in doxorubicin-induced cardiomyopathy. *Cell Commun. Signal.* **21**, 61. <https://doi.org/10.1186/s12964-023-01077-5> (2023).
48. Seddiki, Y., da Silva, F. M. & da Silva, F. M. Antioxidant properties of polyphenols and their potential use in improvement of male fertility: A review. *Biomed. J. Sci. Tech. Res.* **1**, 612–617. <https://doi.org/10.26717/BJSTR.2017.01.000259> (2017).
49. Ortiz-García, C. I. *et al.* Calpain regulates reactive oxygen species production during capacitation through the activation of NOX2 and NOX4. *Int. J. Mol. Sci.* **24**, 3980. <https://doi.org/10.3390/ijms24043980> (2023).
50. Tuzet, A., Rahantaniaina, M. S. & Noctor, G. Analyzing the function of catalase and the ascorbate–glutathione pathway in H2O2 processing: Insights from an experimentally constrained kinetic model. *Antioxid. Redox Signal.* **30**, 1238–1268. <https://doi.org/10.1089/ars.2018.7601> (2019).
51. Siva Kumar, T. & Neeraja, P. Factors associated with oxidative stress in the testes and the mitigating role of antioxidants: A review. *Int. J. Recent Innov. Clin. Med. Res.* **1**, 6–12 (2019).
52. Asadi, N., Bahmani, M., Kheradmand, A. & Rafeian-Kopaei, M. The impact of oxidative stress on testicular function and the role of antioxidants in improving it: A review. *J. Clin. Diagn. Res.* **11**, 1–5 (2017).
53. Guerriero, G., Trocchia, S., Abdel-Gawad, F. K. & Ciarcia, G. Roles of reactive oxygen species in the spermatogenesis regulation. *Front. Endocrinol.* **5**, 56. <https://doi.org/10.3389/fendo.2014.00056> (2014).
54. Shen, Y., Huang, H., Wang, Y., Yang, R. & Ke, X. Antioxidant effects of Se-glutathione peroxidase in alcoholic liver disease. *J. Trace Elem. Med. Biol.* **74**, 127048. <https://doi.org/10.1016/j.jtemb.2022.127048> (2022).
55. Aksu, E. H. *et al.* Rutin ameliorates cisplatin-induced reproductive damage via suppression of oxidative stress and apoptosis in adult male rats. *Andrologia*. **49**, 12593. <https://doi.org/10.1111/and.12593> (2017).
56. Agarwal, A. & Sengupta, P. Oxidative stress and its association with male infertility. *Male Infert. Contemp. Clin. Approach. Androl. ART Antioxidants*. **1**, 57–68. https://doi.org/10.1007/978-3-030-32300-4_6 (2020).
57. Nahid, A., Neelabh, C. & Navneet, K. Antioxidant and antimicrobial potentials of Artemisia Indica collected from the Nepal region. *J. Pharm. Sci. Res.* **9**, 1822–1826 (2017).
58. Kale, A., Gawande, S. & Kotwal, S. Cancer phytotherapeutics: Role for flavonoids at the cellular level. *Phytother. Res.* **22**, 567–577. <https://doi.org/10.1002/ptr.2283> (2008).
59. Ahmed, Z. A., Abtar, A. N., Othman, H. H. & Aziz, T. A. Effects of quercetin, sitagliptin alone or in combination in testicular toxicity induced by doxorubicin in rats. *Drug Des. Devel. Ther.* **13**, 3321–3329. <https://doi.org/10.2147/DDDT.S22127> (2019).
60. Sergeant, N., Briand-Amirat, L., Bencharif, D. & Delehedde, M. The sperm specific protein proAKAP4 as an innovative marker to evaluate sperm quality and fertility. *J. Dairy Vet. Sci.* **11**, 555803. <https://doi.org/10.19080/JDVS.2019.11.555803> (2019).
61. Saïd, L., Banni, M., Kerkeni, A., Saïd, K. & Messaoudi, I. Influence of combined treatment with zinc and selenium on cadmium induced testicular pathophysiology in rat. *Food Chem. Toxicol.* **48**, 2759–2765. <https://doi.org/10.1016/j.fct.2010.07.003> (2010).
62. Barati, E., Nikzad, H. & Karimian, M. Oxidative stress and male infertility: Current knowledge of pathophysiology and role of antioxidant therapy in disease management. *Cell Mol Life Sci.* **77**, 9313. <https://doi.org/10.1007/s00018-019-03253-8> (2020).
63. Dutta, S., Majzoub, A. & Agarwal, A. Oxidative stress and sperm function: A systematic review on evaluation and management. *Arab. J. Urol.* **17**, 87–97. <https://doi.org/10.1080/2090598X.2019.1599624> (2019).
64. Barbagallo, F. *et al.* Effects of bisphenols on testicular steroidogenesis. *Front. Endocrinol.* **11**, 373. <https://doi.org/10.3389/fendo.2020.00373> (2020).
65. Das, J., Ghosh, J., Manna, P. & Sil, P. C. Taurine protects rat testes against doxorubicin-induced oxidative stress as well as p53, Fas and caspase 12-mediated apoptosis. *Amino Acids* **42**, 1839–1855. <https://doi.org/10.1007/s00726-011-0904-4> (2012).
66. Castillo, A. F., Orlando, U., Helfenberger, K. E., Poderoso, C. & Podesta, E. J. The role of mitochondrial fusion and StAR phosphorylation in the regulation of StAR activity and steroidogenesis. *Mol. Cell Endocrinol.* **408**, 73–79. <https://doi.org/10.1016/j.mce.2014.12.011> (2015).
67. Couture, R. *et al.* Luteolin modulates gene expression related to steroidogenesis, apoptosis, and stress response in rat LC540 tumor Leydig cells. *Cell Biol. Toxicol.* **36**, 31–49. <https://doi.org/10.1007/s10565-019-09481-9> (2020).
68. Sun, S., Jin, Y., Luo, P. & Shi, X. Polystyrene microplastics induced male reproductive toxicity and transgenerational effects in freshwater prawn. *Sci. Total Environ.* **842**, 156820. <https://doi.org/10.1016/j.scitotenv.2022.156820> (2022).
69. Martin, L. J. & Touaibia, M. Improvement of testicular steroidogenesis using flavonoids and isoflavonoids for prevention of late-onset male hypogonadism. *Antioxidants*. **9**, 237. <https://doi.org/10.3390/antiox9030237> (2020).
70. Wisniewski, P. *et al.* Adult exposure to bisphenol A (BPA) in Wistar rats reduces sperm quality with disruption of the hypothalamic–pituitary–testicular axis. *Toxicology*. **329**, 1–9. <https://doi.org/10.1016/j.tox.2015.01.002> (2015).
71. Walker, W. H. Testosterone signaling and the regulation of spermatogenesis. *Spermatogenesis*. **1**, 116–120. <https://doi.org/10.4161/spmg.1.2.16956> (2011).
72. Song, Y. J. *et al.* Evaluation of the efficacy of immunocastration vaccine composed of gonadotrophin-releasing hormone conjugated with Salmonella typhimurium flagellin in rats. *Reprod. Domest. Anim.* **47**, 47–50. <https://doi.org/10.1111/j.1439-0531.2011.01931.x> (2012).
73. Walker, W. H. & Cheng, J. FSH and testosterone signaling in Sertoli cells. *Reproduction*. **130**, 15–28. <https://doi.org/10.1530/rep.1.0035> (2005).

74. Agrawal, A. D. Pharmacological activities of flavonoids: A review. *PCI- Approv. IJPSN* **4**, 1394–1398. <https://doi.org/10.37285/ijpsn.2011.4.2.3> (2011).
75. Peña-Blanco, A. & García-Sáez, A. J. Bax, Bak and beyond—mitochondrial performance in apoptosis. *FEBS J.* **285**, 416–431. <https://doi.org/10.1111/febs.14186> (2018).
76. Yang, C. C. *et al.* Assessment of doxorubicin-induced mouse testicular damage by the novel second-harmonic generation microscopy. *Am. J. Transl. Res.* **9**, 5275–5288 (2017).
77. Xu, D. *et al.* TBK1 suppresses RIPK1-driven apoptosis and inflammation during development and in aging. *Cell.* **174**, 1477–1491. <https://doi.org/10.1016/j.cell.2018.07.041> (2018).
78. Lindqvist, L. M., Heinlein, M., Huang, D. C. & Vaux, D. L. Prosurvival Bcl-2 family members affect autophagy only indirectly, by inhibiting Bax and Bak. *Proc. Natl. Acad. Sci.* **111**, 8512–8517. <https://doi.org/10.1073/pnas.140642511> (2014).
79. Hu, C. *et al.* Osteocin attenuates inflammation, oxidative stress, apoptosis, and cardiac dysfunction in doxorubicin-induced cardiotoxicity. *Clin. Transl. Med.* **10**, 124. <https://doi.org/10.1002/ctm2.124> (2020).
80. Dutta, S., Sengupta, P., Slama, P. & Roychoudhury, S. Oxidative stress, testicular inflammatory pathways, and male reproduction. *Int. J. Mol. Sci.* **22**, 10043. <https://doi.org/10.3390/ijms221810043> (2021).
81. Akhter, S., Irfan, H. M., Alamgeer, J. S., Shahzad, M. & Latif, M. B. Nerolidol: a potential approach in rheumatoid arthritis through reduction of TNF- α , IL-1 β , IL-6, NF-kB, COX-2 and antioxidant effect in CFA-induced arthritic model. *Inflammopharmacology.* **30**, 537–548. <https://doi.org/10.1007/s10787-022-00930-2> (2022).
82. Sethi, R. *et al.* Neurobiology and therapeutic potential of cyclooxygenase-2 (COX-2) inhibitors for inflammation in neuropsychiatric disorders. *Front. Psychiat.* **10**, 605. <https://doi.org/10.3389/fpsy.2019.00605> (2019).
83. Chouhan, S. *et al.* Increase in the expression of inducible nitric oxide synthase on exposure to bisphenol A: A possible cause for decline in steroidogenesis in male mice. *Environ. Toxicol. Pharmacol.* **39**, 405–416. <https://doi.org/10.1016/j.etap.2014.09.014> (2015).
84. Adejuwon, S. A., Femi-Akinlosotu, O. M. & Omirinde, J. O. Cisplatin-induced testicular dysfunction and its amelioration by *L. aumaea taraxacifolia* leaf extract. *Andrologia.* **47**, 553–559. <https://doi.org/10.1111/and.12302> (2015).
85. Jin, H. *et al.* Polystyrene microplastics induced male reproductive toxicity in mice. *J Hazard Mater.* **401**, 123430. <https://doi.org/10.1016/j.jhazmat.2020.123430> (2021).

Acknowledgements

The authors extend their appreciation to the Researchers Supporting project number (RSPD2024R729), King Saud University, Riyadh Saudi Arabia for funding this project.

Author contributions

M.U.I., S.Y., A.H. and M.D. designed the study, conceived the study, and analyzed the results. M.U.I. conceived an initial part of the study, performed the experiment and helped in compiling the results. S.Y. wrote the manuscript. M.U.I., A.T., H.A., S.R., and T.A. helped in writing the results. M.U.I., A.H., S.R., T.A., H.A. and A.A. made a substantial contribution in the interpretation of data and revising the manuscript for intellectual content. All authors have read and approved the final manuscript.

Funding

The authors extend their appreciation to the Researchers Supporting project number (RSPD2024R729), King Saud University, Riyadh Saudi Arabia for funding this project. The funding body has no role in designing the study.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.U.I. or S.R.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024