









An efficient Julia framework for hierarchical equations of motion in open quantum systems

Yi-Te Huang ^{1,2,7}, Po-Chen Kuo ^{1,2,7}, Neill Lambert ³✉, Mauro Cirio ⁴✉, Simon Cross ³, Shen-Liang Yang ^{1,2}, Franco Nori ^{3,5,6} & Yueh-Nan Chen ^{1,2}✉

The hierarchical equations of motion (HEOM) approach can describe the reduced dynamics of a system simultaneously coupled to multiple bosonic and fermionic environments. The complexity of exactly describing the system-environment interaction with the HEOM method usually results in time-consuming calculations and a large memory cost. Here, we introduce an open-source software package called HierarchicalEOM.jl: a Julia framework integrating the HEOM approach. HierarchicalEOM.jl features a collection of methods to compute bosonic and fermionic spectra, stationary states, and the full dynamics in the extended space of all auxiliary density operators (ADOs). The required handling of the ADOs multi-indexes is achieved through a user-friendly interface. We exemplify the functionalities of the package by analyzing a single impurity Anderson model, and an ultra-strongly coupled charge-cavity system interacting with bosonic and fermionic reservoirs. HierarchicalEOM.jl achieves a significant speedup with respect to the corresponding method in the Quantum Toolbox in Python (QuTiP), upon which this package is founded.

¹Department of Physics, National Cheng Kung University, 701 Tainan, Taiwan. ²Center for Quantum Frontiers of Research and Technology, NCKU, 701 Tainan, Taiwan. ³Theoretical Quantum Physics Laboratory, Cluster for Pioneering Research, RIKEN, Wakoshi, Saitama 351-0198, Japan. ⁴Graduate School of China Academy of Engineering Physics, Haidian District, Beijing 100193, China. ⁵Center for Quantum Computing (RQC), RIKEN, Wakoshi, Saitama 351-0198, Japan. ⁶Physics Department, The University of Michigan, Ann Arbor, MI 48109-1040, USA. ⁷These authors contributed equally: Yi-Te Huang, Po-Chen Kuo. ✉email: nwlambert@gmail.com; cirio.mauro@gmail.com; yuehnan@mail.ncku.edu.tw

The time evolution of a closed quantum system can offer important insights about its nature and properties. However, the dynamics are inevitable affected by interactions with external environments^{1,2}, which can involve the exchange of energy or particles, and the suppression of quantum coherence. Due to the effective continuum of degrees of freedom present in these external baths^{3,4}, modeling the dynamics of an open quantum system can be challenging. This is especially the case when perturbative approaches⁵ are no longer valid due to non-Markovian effects emerging in the presence of strong interaction with the bath^{6,7}. In this regime, standard Markovian master equations are no longer applicable, and non-perturbative techniques are required^{8–12}.

In particular, here we consider the hierarchical equations of motion (HEOM) approach, which offers a non-perturbative¹³ characterization of all the environmental effects on the system. This is achieved by using a hierarchy of auxiliary density operators (ADOs) to model system-bath correlations and entanglement^{14,15}. This makes the HEOM suitable for studying complex systems strongly coupled to either bosonic (with applications in quantum biology^{16,17}, quantum optics¹⁸, quantum thermodynamics¹⁹, and quantum information²⁰), or fermionic environments (for the analysis of quantum transport^{21–23} and condensed matter physics²⁴). Additionally, quantum systems interacting simultaneously with both bosonic and fermionic environments can be found in the study of electron transport through both natural and artificial molecules^{25,26}. Naturally, such an increase in the complexity of the environment leads to an increase in computational resources which, in the case of the HEOM method, corresponds to an increase in the size of the HEOM Liouvillian superoperator (HEOMLS) matrix. To deal with this issue, it is beneficial to explore the numerical efficiency of programming languages designed to optimize different computational resources.

In this work, we use Julia^{27,28}: a dynamic general-purpose, high-level programming language capable of high-performance in scientific applications. It was created in 2012 with the goal to build a language with the speed of C²⁹, the dynamism of Ruby³⁰, the practicality of Python³¹, and able to solve statistics and linear algebra tasks like R³² and Matlab³³, respectively. Julia is fast because it uses a just-in-time compiler³⁴ to convert source code into machine code before running it. Recently, many researchers are turning to Julia to build high-performance simulations and models for open quantum system dynamics³⁵, quantum optics³⁶, quantum algorithms³⁷ and quantum information theories³⁸. Moreover, the variable name in Julia supports UTF-8 encoding (e.g., Greek symbols and mathematical symbols) which allows to write equations in code more elegantly.

HierarchicalEOM.jl is developed following the Julia design philosophy: one can have machine performance without sacrificing human convenience²⁷. While integrating many of the features presented in other open-source HEOM packages, HierarchicalEOM.jl also includes other functionalities, such as the estimation of importance values for all ADOs, the calculation of spectra for both bosonic and fermionic systems, the construction of HEOMLS matrices for even- or odd-parity auxiliary density operators, and a user-friendly interface (which interrogates the ADOs multi-indexes) for gaining access to bath properties. By wrapping some functions from other Julia packages^{39–41}, we could further optimize the computations of the dynamics and stationary states for all ADOs. The entire ecosystem of HierarchicalEOM.jl package is summarized in Fig. 1.

Results

Preliminaries for hierarchical equations of motion. Throughout this work, we consider an open quantum system (s) interacting

with *fermionic* (f) and *bosonic* (b) environments described by the following total (T) Hamiltonian (\hbar is set to unity throughout this work):

$$H_T = H_s(t) + H_f + H_b + H_{sf} + H_{sb}, \quad (1)$$

where $H_s(t)$ is the (possibly time-dependent) system Hamiltonian containing boson and fermion particles. Here, we allow the fermionic environment to be composed by multiple baths of non-interacting fermionic degrees of freedom described by the Hamiltonian

$$H_f = \sum_{\alpha} \sum_k \epsilon_{\alpha,k} c_{\alpha,k}^{\dagger} c_{\alpha,k}, \quad (2)$$

where $c_{\alpha,k}$ ($c_{\alpha,k}^{\dagger}$) annihilates (creates) a fermion (f) in the state k (with energy $\epsilon_{\alpha,k}$) of the α th fermionic bath. Analogously,

$$H_b = \sum_{\beta} \sum_k \omega_{\beta,k} b_{\beta,k}^{\dagger} b_{\beta,k}, \quad (3)$$

describes a generic bosonic environment which can accommodate multiple non-interacting bosonic baths in which $b_{\beta,k}$ ($b_{\beta,k}^{\dagger}$) is the bosonic annihilation (creation) operator associated to the k th mode (with frequency $\omega_{\beta,k}$) in the β th bosonic bath. This bosonic environment can represent a broad range of physical environments, e.g., electromagnetic environments^{42,43}, phonon environments^{44–46}, surface plasmon modes^{47,48} and vibronic environments in molecules (e.g., nuclear motion in a photosynthetic complex²²). The interaction Hamiltonian between a fermionic system and the fermionic environments can be written as

$$H_{sf} = \sum_{\alpha,k} \left(g_{\alpha,k} c_{\alpha,k}^{\dagger} d_s + g_{\alpha,k}^* d_s^{\dagger} c_{\alpha,k} \right), \quad (4)$$

in terms of the *coupling strengths* $g_{\alpha,k}$. Analogously, the interaction between a bosonic or fermionic system and the exterior bosonic environments can be modeled by

$$H_{sb} = V_s \sum_{\beta,k} g_{\beta,k} (b_{\beta,k} + b_{\beta,k}^{\dagger}), \quad (5)$$

in terms of the coupling strengths $g_{\beta,k}$. Here, d_s and V_s refer to the coupling operators acting on the system's degrees of freedom. In particular, d_s is a odd-parity operator destroying a fermion in the system, while V_s is in general a Hermitian operator which can act on both fermionic and bosonic systems. When V_s is acting on the fermionic system, it must have even-parity to be compatible with charge conservation. Furthermore, one can easily generalize to the case where the system contains multiple bosonic and fermionic quantum numbers (such as frequency, energy, or spin) interacting with either individual or shared environment(s) [see Example 1 and Example 2].

To derive the HEOM, we assume the following three conditions: (1) The system and the environments (baths) are initialized in a separable state. (2) Each of the fermionic (bosonic) baths is initially in thermal equilibrium characterized by a Fermi-Dirac (Bose-Einstein) distribution. (3) The bath operator within the system-bath interaction Hamiltonian should be linear in the bath annihilation and creation operators, as shown in Eq. (4) and Eq. (5).

Without resorting to any approximation, the reduced density matrix for the system at time t can be written in terms of the following Dyson series

$$\rho_s^{p=\pm}(t) = \hat{\mathcal{G}}(t) [\rho_s^{p=\pm}(0)], \quad (6)$$

where, the propagator $\hat{\mathcal{G}}(t)[\cdot]$ is a *canonical superoperator*⁴⁹ which propagates the even-parity ($p=+$) or odd-parity ($p=-$) operator initially in $\rho_s^p(0)$. Furthermore, the reduced density matrix of the system has even-parity and can be written as $\rho_s^+(0)$.




Fig. 1 The ecosystem of the HierarchicalEOM.jl package. **a** Users should specify the system Hamiltonian $H_s(t)$, coupling operators (V_s or d_s), and the bath correlation function $C(t)$. For the exponent $\{\eta_k, \gamma_k\}$, users can either specify the physical parameters characterizing the spectral density of the bath by built-in functions, or directly providing a list of exponents. **b** Construction of the bath-object which includes the system coupling operator and a list of exponents characterizing the bath correlation function. **c** Construction of the HEOM Liouvillian superoperator (HEOMLS) matrix $\hat{\mathcal{M}}$ which defines the hierarchical equations of motion from the system Hamiltonian and the bath-objects. **d** Computation of the dynamics and stationary states for all auxiliary density operators using $\hat{\mathcal{M}}$. **e** The hierarchy dictionary translates the index of each ADO into the corresponding multi-index ensembles together with the exponents of the bath, and vice-versa. **f** The hierarchy dictionary allows a high-level interpretation of the ADOs to compute some physical properties. **g** Logo of HierarchicalEOM.jl package.

The propagator takes the form

$$\hat{\mathcal{G}}(t)[\cdot] = \hat{T} \exp \left\{ - \int_0^t dt_1 \int_0^{t_1} dt_2 [\hat{\mathcal{W}}_f(t_1, t_2)[\cdot] + \hat{\mathcal{W}}_b(t_1, t_2)[\cdot]] \right\}, \quad (7)$$

in terms of the time-ordering operator \hat{T} , together with fermionic⁴⁹ and bosonic^{15,16} operators explicitly written as

$$\begin{aligned} \hat{\mathcal{W}}_f(t_1, t_2)[\cdot] = & \sum_{\alpha} \sum_{p=\pm} \sum_{\nu=\pm 1} \left\{ C_{\alpha}^{\nu}(t_1, t_2) [d_s^{\nu}(t_2), d_s^{\nu}(t_1) \cdot]_{-p} \right. \\ & \left. + C_{\alpha}^{\nu}(t_2, t_1) [\cdot d_s^{\nu}(t_2), d_s^{\nu}(t_1)]_{-p} \right\}, \end{aligned} \quad (8)$$

and

$$\hat{\mathcal{W}}_b(t_1, t_2)[\cdot] = \sum_{\beta} [V_s(t_1), (C_{\beta}^R(t_1, t_2)[V_s(t_2), \cdot]_{-} + iC_{\beta}^I(t_1, t_2)[V_s(t_2), \cdot]_{+})]_{-}, \quad (9)$$

respectively. For simplicity, we define ν to denote the presence ($\nu = +1$) or absence ($\nu = -1$) of a Hermitian conjugation and $\bar{\nu} := -\nu$. Here, $[\cdot, \cdot]_{-}$ and $[\cdot, \cdot]_{+}$ represent the commutator and the anti-commutator introduced to allow the propagator to keep track of the parity sector (labeled by $p = \pm$) it is acting on. This dependence intuitively originates from the properties of partial traces over composite fermionic spaces, where operators do not necessarily commute.

As an example, for an environment made out of a single fermion, the reduced matrix elements $\langle i | \rho_s^p | j \rangle$ (in a basis labeled by $\langle i |$ and $| j \rangle$) involve the perturbative sum of expressions of the form $\langle i | (c \tilde{\rho}_e \tilde{\rho}_s^p c^{\dagger} + \tilde{\rho}_e \tilde{\rho}_s^p) | j \rangle$ (in terms of environmental operators $\tilde{\rho}_e$, system operators $\tilde{\rho}_s^p$ with parity p , and the environment-annihilation operator c). These quantities depend on the commutator between $\tilde{\rho}_s^p$ and c , which is trivial only for even-parity ($p = +$). In the odd-parity ($p = -$) case, the partial trace over the environment requires further anti-commutations, ultimately resulting in extra minus signs in the expression for the effective propagator describing the reduced dynamics⁴⁹. It is important to explicitly note that, here, by parity we do not refer to the presence of an odd or even number of fermions in the system but, rather, to the number of fermionic operators needed to represent ρ_s^p . In summary, while commutators ($-p = -$) appear

for the propagation of even-parity operators $\rho_s^{p=+}(0)$ (which include the reduced density operator of the system), anti-commutators ($-p = +$) appear for the propagation of odd-parity operators $\rho_s^{p=-}(0)$ [e.g., $d_s \rho_s^{+}(0)$ or $d_s^{\dagger} \rho_s^{+}(0)$], which appear in the computation of observables such as the density of states (see the Methods section: Numerical computation of the spectrum).

As it can be seen in Eq. (8) and Eq. (9), the effects of fermionic and bosonic environments [initially in thermal equilibrium (eq) and linearly coupled to the system] are completely encoded in the *two-time correlation functions* $C(t_1, t_2)$. In the fermionic case, they depend on the spectral density $J_{\alpha}(\omega) = 2\pi \sum_k |g_{\alpha,k}|^2 \delta(\omega - \omega_k)$ and the Fermi-Dirac distribution $n_{\alpha}^{\text{eq}}(\omega) = \{\exp[(\omega - \mu_{\alpha})/k_B T_{\alpha}] + 1\}^{-1}$ as

$$C_{\alpha}^{\nu}(t_1, t_2) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega J_{\alpha}(\omega) \left[\frac{1 - \nu}{2} + \nu n_{\alpha}^{\text{eq}}(\omega) \right] e^{\nu i\omega(t_1 - t_2)}. \quad (10)$$

Analogously, in the bosonic case, they depend on the spectral density $J_{\beta}(\omega) = 2\pi \sum_k a_{\beta,k}^2 \delta(\omega - \omega_k)$ and the Bose-Einstein distribution $n_{\beta}^{\text{eq}}(\omega) = \{\exp[\omega/k_B T_{\beta}] - 1\}^{-1}$ as

$$C_{\beta}(t_1, t_2) = \frac{1}{2\pi} \int_0^{\infty} d\omega J_{\beta}(\omega) \left[n_{\beta}^{\text{eq}}(\omega) e^{i\omega(t_1 - t_2)} + (n_{\beta}^{\text{eq}}(\omega) + 1) e^{-i\omega(t_1 - t_2)} \right]. \quad (11)$$

Here, k_B is the Boltzmann constant and $T_{\alpha}(T_{\beta})$ represents the absolute temperature of the α -fermionic (β -bosonic) bath. A non-zero chemical potential ($\mu_{\alpha} \neq 0$) in the α -fermionic bath can account for non-equilibrium physics.

A more practical representation of Eq. (6) can be found by expressing the bath correlation functions as a sum of exponential terms (exponents). This allows to define an iterative procedure which leads to the celebrated hierarchical equations of motion (HEOM). Specifically, different approaches, such as the Matsubara⁵⁰ or the Padé⁵¹ spectral decompositions, can be used (see Methods) to write the fermionic and bosonic correlation functions as, respectively,

$$C_{\alpha}^{\nu}(\tau) = \sum_{h=1}^{N_{\alpha}} \eta_{\alpha,h}^{\nu} \exp(-\gamma_{\alpha,h}^{\nu} \tau), \quad (12)$$

and

$$C_\beta(\tau) = \sum_{l=1}^{N_\beta} \xi_{\beta,l} \exp(-\chi_{\beta,l}\tau), \quad (13)$$

where $\tau = t_1 - t_2$, and N_α (N_β) is the total number of exponentials for the α -fermionic (β -bosonic) bath. When $\chi_{\beta,l} \neq \chi_{\beta,l}^*$, a closed form for the HEOM can be obtained by further decomposing the bosonic correlation function into its real (\mathbb{R}) and imaginary (\mathbb{I}) parts as

$$C_\beta(\tau) = \sum_{u \in \mathbb{R}, \mathbb{I}} (\delta_{u,\mathbb{R}} + i\delta_{u,\mathbb{I}}) C_\beta^u(\tau) \quad (14)$$

where δ is the Kronecker delta function and where

$$C_\beta^u(\tau) = \sum_{l=1}^{N_\beta^u} \xi_{\beta,l}^u \exp(-\chi_{\beta,l}^u \tau). \quad (15)$$

The expressions for the fermionic [Eq. (12)] and bosonic [Eq. (14)] correlation functions can now be used in Eq. (7). The resulting expression can be recursively differentiated in time^{18,49,52} to define a local master equation in an enlarged space made out of *auxiliary density operators* (ADOs) $\rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}(t)$. Intuitively, these variables encode environmental effects related to different exponential terms present in the correlation function and provide an iterative description of high-order system-baths memory effects. In $\rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}(t)$, the tuple (m, n, p) represents the m th-level-bosonic-and- n th-level-fermionic ADO with parity p , and \mathbf{j} (\mathbf{q}) denotes a vector $[j_m, \dots, j_1]$ ($[q_m, \dots, q_1]$), where each j (q) represents a specific multi-index ensemble $\{\beta, u, l, \sigma_b\}$ ($\{\alpha, v, h, \sigma_f\}$). The indexes σ_b and σ_f are used to specify other quantum numbers (such as energy or spin) of the system interacting with a bosonic or fermionic bath, respectively. In this way, the hierarchical equations of motion in the Schrödinger picture can be expressed as

$$\begin{aligned} \partial_t \rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}(t) &\equiv \hat{\mathcal{M}} \rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}(t) \\ &= - \left(i\hat{\mathcal{L}}_s + \sum_{r=1}^m \chi_r + \sum_{w=1}^n \gamma_w \right) \rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}(t) \\ &\quad - i \sum_{q' \notin \mathbf{q}} \hat{\mathcal{A}}_{q'} \rho_{\mathbf{j}|\mathbf{q}'}^{(m,n+1,p)}(t) - i \sum_j \hat{\mathcal{B}}_j \rho_{\mathbf{j}^+|\mathbf{q}}^{(m+1,n,p)}(t) \\ &\quad - i \sum_{w=1}^n (-1)^{n-w} \hat{\mathcal{C}}_w \rho_{\mathbf{j}|\mathbf{q}_w}^{(m,n-1,p)}(t) - i \sum_{r=1}^m \hat{\mathcal{D}}_r \rho_{\mathbf{j}^-|\mathbf{q}}^{(m-1,n,p)}(t), \end{aligned} \quad (16)$$

where we used the multi-index notations:

$$\begin{aligned} \mathbf{j}^+ &= [j', j_m, \dots, j_1], \\ \mathbf{q}^+ &= [q', q_n, \dots, q_1], \\ \mathbf{j}^- &= [j_m, \dots, j_{r+1}, j_{r-1}, \dots, j_1], \\ \mathbf{q}_w^- &= [q_n, \dots, q_{w+1}, q_{w-1}, \dots, q_1]. \end{aligned} \quad (17)$$

Here, j' can be chosen from any one of the elements in \mathbf{j} or any other bosonic multi-index ensemble that has not yet been considered in \mathbf{j} , whereas $q' \notin \mathbf{q}$ due to the Pauli exclusion principle.

Furthermore, we defined $\hat{\mathcal{L}}_s[\cdot] = [H_s(t), \cdot]_-$ characterizing the bare system dynamics. System-bath interactions are encoded by the fermionic superoperators $\hat{\mathcal{A}}_q, \hat{\mathcal{C}}_q$ (to couple the n th-level-fermionic ADOs to the $(n+1)$ th-level- and $(n-1)$ th-level-fermionic ADOs, respectively), and the bosonic superoperators $\hat{\mathcal{B}}_j, \hat{\mathcal{D}}_j$ (to couple the m th-level-bosonic ADOs to the $(m+1)$ th-level- and $(m-1)$ th-level- bosonic ADOs, respectively). Their

explicit forms are given by

$$\begin{aligned} \hat{\mathcal{A}}_q[\cdot] &= (-1)^{\delta_{p,-}} \left\{ d_{\sigma_f}^\nu[\cdot] - \hat{\mathcal{P}}_s \left[[\cdot] d_{\sigma_f}^\nu \right] \right\}, \\ \hat{\mathcal{C}}_q[\cdot] &= (-1)^{\delta_{p,-}} \left\{ \eta_{\alpha,h}^\nu d_{\sigma_f}^\nu[\cdot] + (\eta_{\alpha,h}^\nu)^* \hat{\mathcal{P}}_s \left[[\cdot] d_{\sigma_f}^\nu \right] \right\}, \\ \hat{\mathcal{B}}_j[\cdot] &= [V_{\sigma_b}, \cdot]_-, \\ \hat{\mathcal{D}}_j[\cdot] &= \delta_{u,\mathbb{R}} \xi_{\beta,l}^{\mathbb{R}} [V_{\sigma_b}, \cdot]_- + i\delta_{u,\mathbb{I}} \xi_{\beta,l}^{\mathbb{I}} [V_{\sigma_b}, \cdot]_+, \\ \hat{\mathcal{P}}_s \left[\rho_{\mathbf{j}|\mathbf{q}}^{(m,n,\pm)}(t) d_{\sigma_f}^\nu \right] &= \mp (-1)^n \rho_{\mathbf{j}|\mathbf{q}}^{(m,n,\pm)}(t) d_{\sigma_f}^\nu. \end{aligned} \quad (18)$$

The form of the HEOM in Eq. (16) can be applied to auxiliary density operators with arbitrary parity symmetry. For example, $\rho_1^{(0,0,+)}(t)$ is the system reduced density operator, and the odd-parity operator $\rho_1^{(0,0,-)}(t)$ can take the form $d_s^\nu \rho_1^{(0,0,+)}(t)$ as mentioned after Eq. (9). To project in the different parity system sub-spaces, we introduced the superoperator

$$\hat{\mathcal{P}}_s[\cdot] = \left(\prod_{\sigma_f} \exp[i\pi d_{\sigma_f}^\dagger d_{\sigma_f}] \right) [\cdot] \left(\prod_{\sigma_f} \exp[i\pi d_{\sigma_f}^\dagger d_{\sigma_f}] \right). \quad (19)$$

Overall, the HEOM *Liouvillian superoperator* (HEOMLS) $\hat{\mathcal{M}}$ characterizes the dynamics in the full ADOs space and it can, numerically, be expressed as a matrix.

In practice, the dimensions of this matrix must be finite and the hierarchical equations must be truncated at a suitable bosonic-level (m_{\max}) and fermionic-level (n_{\max}). Previous works^{23,53} showed that the computational effort can be optimized by associating an *importance value* to each ADO and then discarding all ADOs (in the second and higher levels) whose importance value is smaller than a threshold value \mathcal{I}_{th} . The importance value \mathcal{I} for a given ADO $\rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)}$ is defined as

$$\mathcal{I} \left(\rho_{\mathbf{j}|\mathbf{q}}^{(m,n,p)} \right) = \left| \prod_{r=1}^m \left(\frac{\xi_{j_r}}{\text{Re}[\chi_{j_r}]} \frac{1}{\sum_{x=1}^r \text{Re}[\chi_{j_x}]} \right) \right| \times \left| \prod_{w=1}^n \left(\frac{\eta_{q_w}}{\text{Re}[\gamma_{q_w}]} \frac{1}{\sum_{x=1}^w \text{Re}[\gamma_{q_x}]} \right) \right|, \quad (20)$$

where the first and second absolute value represents the importance of the bosonic part²⁶ and fermionic part⁵⁴ of the hierarchy, respectively. Note that for hybrid (consider bosonic and fermionic baths at the same time) cases, we retain all the ADOs where their hierarchy levels $(m, n) \in \{(0,0), (0,1), (1,0), (1,1)\}$, and all the other high-level ADOs may be neglected if their importance value is less than \mathcal{I}_{th} . Moreover, the full hierarchical equations can be recovered in the limiting case $\mathcal{I}_{\text{th}} \rightarrow 0$. We will provide a concrete example and detailed discussion later.

Package architecture and design philosophy. The package `HierarchicalEOM.jl` is designed to integrate the efficiency of Julia with the functionalities provided by other existing HEOM packages^{15,24,55–57}. This leads to an intuitive interface to construct arbitrary Hamiltonians and initial states (which can be accomplished directly with the built-in library for linear algebra in Julia or by taking advantage of the package `QuantumOptics.jl`³⁶). We now introduce the ecosystem of the package as summarized in Fig. 1.

Following Fig. 1(a) and Fig. 1(b), users should specify the system Hamiltonian $H_s(t)$, system coupling operators V_s (d_s) describing the interaction with bosonic (fermionic) baths, and the bath correlation functions. For the bath correlation functions in Eq. (10) and Eq. (11), users can specify them by an exponential series as shown in Eqs. ((12)–(15)). `HierarchicalEOM.jl` provides built-in functions to construct these series from physical

parameters (e.g., coupling strength, temperature, etc.). For example, it is possible to choose a Drude-Lorentz spectral density for bosonic baths and a Lorentzian spectral density for fermionic baths. For each of these cases, the correlations are computed using either the Matsubara⁵⁰ or Padé spectral decomposition⁵¹. Alternatively, `HierarchicalEOM.jl` offers the possibility to manually define the correlation functions (simply supplying the list of exponents). Additional spectral densities could be incorporated into the built-in functions in future releases. We explicitly note that the package allows for any combination of fermionic and bosonic baths.

After the definition of the bath, users can further construct the HEOMLS matrix \hat{M} in Eq. (16) together with the system Hamiltonian $H_s(t)$, see Fig. 1c. At this stage, `HierarchicalEOM.jl` offers the possibility to optimize this matrix by neglecting all the ADOs (at second or higher level) whose importance values in Eq. (20) are below the user-specified threshold \mathcal{I}_{th} . It is also possible to construct \hat{M} within the even- or odd-parity sector depending on the parity of the auxiliary density operator upon which the HEOMLS matrix is acting on (see the Methods section: Numerical computation of the spectrum). Moreover, our package allows users to further add a *Lindbladian* (superoperator) to \hat{M} . The Lindbladian describes the dissipative interaction between the system and extra environment while its explicit form is given by

$$\tilde{\mathcal{J}}(F)[\cdot] = F[\cdot]F^\dagger - \frac{1}{2}[F^\dagger F, \cdot]_+, \quad (21)$$

where F is the jump operator and describes the dissipation dynamics of the system. Including such Lindbladians when some of the baths are weakly coupled to the system is more efficient than solving the full HEOM since it does not require any ADOs for such baths and hence reduces the size of \hat{M} (see Example 2 for more details).

As shown in Fig. 1d, with this information it is possible to proceed with solving for the dynamics of all the ADOs as described in Eq. (16). `HierarchicalEOM.jl` provides two distinct methods to compute the dynamics. The first one relies on `DifferentialEquations.jl`³⁹ which provides a set of low-level solvers for ordinary differential equations. The second method directly builds the propagator when the system Hamiltonian is time-independent: $\hat{G}(t) = \exp(\hat{M}t)$ using `FastExp.jl`⁴¹ which is optimized for the exponentiation of either large-dense or sparse matrices. In addition, `HierarchicalEOM.jl` can also directly solve for the stationary states of all the ADOs using `LinearSolve.jl`⁴⁰, which offers a unified interface to solve linear equations in Julia.

To allow further analysis of specific physical properties, `HierarchicalEOM.jl` provides a hierarchy dictionary, as depicted in Fig. 1e, f. This dictionary translates the index of each ADO in terms of the corresponding exponential terms of the bath, and vice-versa. This feature is designed to allow a high-level description of the ADOs, which can be useful in the analysis of electronic currents^{21,23}, heat currents^{19,58,59}, and higher-order moments of heat currents⁶⁰. Moreover, `HierarchicalEOM.jl` can calculate the spectrum for both bosonic and fermionic systems using `LinearSolve.jl` (see the Methods section: Numerical computation of the spectrum).

We now demonstrate the use of `HierarchicalEOM.jl` with two examples: a single impurity strongly coupled to fermionic reservoirs and an ultra-strongly coupled charge-cavity system interacting with both bosonic and fermionic reservoirs.

Example 1: Electronic system strongly interacting with fermionic reservoirs. The investigation of the Kondo effect in

electronic systems is crucial as it serves both as a valuable testing ground for the theories of the Kondo effect and has the potential to lead to a better understanding of this intrinsic many-body phenomena^{61–63}, with applications in building new generation nanoscale electronic devices^{64–66} and quantum computation⁶⁷. In this sense, we here consider a single-level electronic system [which can be populated by a spin up, $\sigma_f = \uparrow$, or spin down electron, $\sigma_f = \downarrow$] coupled to two [left (L) and right (R)] fermionic reservoirs, as illustrated in Fig. 2. In this case, the different Hamiltonian terms in Eq. (1) take the form

$$\begin{aligned} H_s &= \epsilon \left(d_\uparrow^\dagger d_\uparrow + d_\downarrow^\dagger d_\downarrow \right) + U \left(d_\uparrow^\dagger d_\uparrow d_\downarrow^\dagger d_\downarrow \right), \\ H_f &= \sum_{\alpha=L,R} \sum_{\sigma_f=\uparrow,\downarrow} \sum_k \epsilon_{\alpha,\sigma_f,k} c_{\alpha,\sigma_f,k}^\dagger c_{\alpha,\sigma_f,k}, \\ H_{sf} &= \sum_{\alpha=L,R} \sum_{\sigma_f=\uparrow,\downarrow} \sum_k g_{\alpha,k} c_{\alpha,\sigma_f,k}^\dagger d_{\sigma_f} + g_{\alpha,k}^* d_{\sigma_f}^\dagger c_{\alpha,\sigma_f,k}, \end{aligned} \quad (22)$$

where ϵ is the energy of the impurity and U is the Coulomb repulsion energy for double occupation. We further assume both fermionic reservoirs to have a Lorentzian-shaped spectral density. As we discussed above, the correlation function in Eq. (10) can be expressed as a sum of exponential terms, see Eq. (12), using the Padé decomposition (see Methods). These spectral densities depend on the following physical parameters: the coupling strength $\Gamma_L = \Gamma_R = \Gamma$ between system and reservoirs, the band-width W_ω , the temperature T , and the chemical potential $\mu_L = -\mu_R = e\Phi/2$, where e is the elementary charge, and Φ is the bias voltage. We summarize the values of these parameters in Table. 1. With these choices, it is now possible to use `HierarchicalEOM.jl` to construct the HEOMLS matrix \hat{M} and solve for the stationary states of all the auxiliary density operators (ADOs).

By using the built-in function in `HierarchicalEOM.jl`, we first compute the spectrum of the system (also known as density of states (DOS); see Methods) in the stationary state and plot the results in Fig. 3a. The DOS of the system exhibits two




Fig. 2 Schematic illustration of a light-matter system (an electron coupled to a bosonic mode in a cavity) interacting with bosonic and fermionic reservoirs. Here, g is the coupling strength between the electron and cavity, Δ is the coupling strength between the cavity and the bosonic reservoir, and $\Gamma_L = \Gamma_R = \Gamma$ are the coupling strengths between the electron and its fermionic reservoirs.

Table 1 Parameters used in the examples.

| Example | ϵ | ω_c | U | ω_c | g | Γ | Δ | W_β | W_α | $e\Phi$ | $k_B T$ | N_{photon} | N_β | N_α | m_{max} | n_{max} | \mathcal{I}_{th} |
|---------|------------|------------|-----|------------|----------|----------|----------|-----------|------------|---------|---------|---------------------|-----------|------------|------------------|------------------|---------------------------|
| (1) | -5 | x | 10 | x | x | 1 | x | x | 10 | [0, 4] | 0.5 | x | x | 7 | x | 4 | 10^{-7} |
| (2) | -3 | 1 | x | x | [0, 0.5] | 1 | 0.01 | 0.2 | 10 | [0, 10] | 0.5 | x | 5 | 7 | 4 | 3 | 10^{-6} |

Here, ϵ and ω_c are, respectively, the energies of the electronic state and the frequency of the cavity mode, while U represents the Coulomb repulsion energy. The electron-cavity, electron-(fermionic reservoir), and cavity-(bosonic reservoir) couplings strengths are specified by g , Γ , and Δ , respectively. W_α and W_β are the band-widths of the bosonic and fermionic reservoir, respectively. The bias voltage Φ is used to characterize the chemical potential of the fermionic reservoirs as $\mu_L = -\mu_R = e\Phi/2$ (in terms of the elementary charge e). All the reservoirs are initially at equilibrium at the same value of $k_B T$, where k_B is the Boltzmann constant and T is the absolute temperature. Note that all the values are considered in units of millielectronvolt (meV). The truncation parameters are: N_{photon} for the Fock space dimension, N_β for the number of exponential terms for the bosonic reservoirs, N_α for the number of exponential terms for the fermionic reservoirs, m_{max} for the tier of the bosonic hierarchy, n_{max} for the tier of the fermionic hierarchy, and \mathcal{I}_{th} for the importance threshold.

Hubbard peaks and an additional central peak. The negative frequency Hubbard peak is a consequence of the singly occupied electron state with either spin-up or spin-down. Similarly, double occupation of the state contributes to the resonance at energy $\epsilon + U$. When the system is in equilibrium, the Kondo-resonance central peak is a consequence of the many-body entanglement between the single electron populating the system and the electrons in the fermionic reservoirs. As the bias voltage Φ is increased, the central peak splits into two⁶⁸. This ultimately results in two distinct energies for the Kondo cloud which depend on the Fermi-level of the fermionic reservoirs^{69,70}.

Alternatively, the Kondo effect can also be probed by calculating the differential conductance $G = dI/d\Phi$ between the system and fermionic reservoirs. To compute this quantity, we first compute the electronic current in the stationary states of the ADOs (see Methods) as a function of the bias voltage Φ and then take the derivative numerically. The zero-bias conductance peak shown in Fig. 3b is another manifestation of the existence of a single-channel spin-1/2 Kondo resonance. In fact, in this regime, the system effectively acts as a one-dimensional channel for the moving electrons thereby increasing the conductance⁷¹.

Example 2: The ultra-strongly coupled charge-cavity system interacting with bosonic and fermionic reservoirs. A crucial aspect of quantum transport research involves investigating impurities, coupled to normal metal contacts, whose charge degree of freedom is strongly coupled to cavity photons⁷²⁻⁷⁴. By integrating quantum dots with microwave cavities, researchers can investigate such novel cavity quantum electrodynamics systems⁷⁵, yielding insights into a diverse range of quantum phenomena such as out-of-equilibrium state occupation in double quantum dots⁷⁶, reservoir-induced charge relaxation in single dots⁷⁷, and photo-assisted tunneling processes^{72,78}. Going beyond the strong coupling regime, we here consider a single-level electronic system ultra-strongly coupled to a single-mode cavity⁷⁹. The electronic impurity is in contact with two distinct, left (L) and right (R), fermionic reservoirs, while the cavity is in contact with a bosonic reservoir, as illustrated in Fig. 2. In order to focus on the charge-photon interaction, this model neglects the electronic spin degree of freedom. As a consequence, the different contributions to the total Hamiltonian, as given in Eq. (1), can be expressed as

$$\begin{aligned}
 H_s &= \epsilon d^\dagger d + \omega_c a^\dagger a + g d^\dagger d (a + a^\dagger), \\
 H_f &= \sum_{\alpha=L,R} \sum_k \epsilon_{\alpha,k} c_{\alpha,k}^\dagger c_{\alpha,k}, \\
 H_b &= \sum_k \omega_k b_k^\dagger b_k, \\
 H_{sf} &= \sum_{\alpha=L,R} \sum_k g_{\alpha,k} c_{\alpha,k}^\dagger d + g_{\alpha,k}^* d^\dagger c_{\alpha,k}, \\
 H_{sb} &= \sum_k g_k (a + a^\dagger) (b_k + b_k^\dagger),
 \end{aligned} \tag{23}$$

in terms of the energy ϵ of the electronic state, the frequency ω_c of the single-mode cavity, the electron-cavity coupling strength g , and the annihilation (creation) operator a (a^\dagger) of the single-mode cavity. We assume a Drude-Lorentz (Lorentzian) spectral density for the bosonic (fermionic) reservoir(s). As previously discussed, the correlation function in Eq. (10) and Eq. (11) can be expressed as a sum of exponential terms using the Padé decomposition, as shown in Eq. (12) and Eq. (13), respectively.

The environmental spectral densities depend on the following physical parameters: the coupling strength $\Gamma_L = \Gamma_R = \Gamma$ between the electron and the fermionic reservoirs, the coupling strength Δ between the single-mode cavity and the bosonic reservoir, the band-width W_α (W_β) of the fermionic (bosonic) reservoirs, the




Fig. 3 The Kondo resonance from equilibrium to non-equilibrium. In **a**, the black dashed, red dash-dotted, orange dotted, green solid, and blue dashed curves represent the density of states $A(\omega)$ for bias voltage Φ at 0, 1, 2, 3, and 4 mV, respectively. One can observe that the density of states of the electronic system exhibits two Hubbard peaks and an additional central peak at equilibrium (i.e., bias voltage $\Phi = 0$). By increasing the bias voltage Φ to non-equilibrium, the central peak is suppressed and splits into two. In **b**, the black curve shows the effects of the Kondo resonance on the differential conductance as a function of bias voltage Φ . The conductance G has a peak at $\Phi = 0$ as the Kondo resonance acts as a transport channel for the electron.




Fig. 4 Effects of the electron-cavity coupling on the spectra of the charge and cavity system. In **a**, the black dashed, red dotted, and blue solid curves represent the density of states $A(\omega)$ for the charge-cavity coupling strength g at 0, 0.25Γ , and 0.5Γ , respectively. The resonance in the density of states of the charge system is reduced and shifts towards lower energies as g increases. In **b**, we plot the power spectral density $S(\omega)$ of the cavity and uses two different methods [HEOM (red dotted and blue solid curves for g at 0.25Γ and 0.5Γ , respectively) and perturbative master equation (abbreviated as ME, which is represented by orange dash-dotted and green dashed curves for g at 0.25Γ and 0.5Γ , respectively)] to describe the interaction between the cavity and the bosonic reservoir. Both approaches show a single peak at $\omega = \omega_c$ for $g \in (0, 0.5]$. The Born-Markov approximation used to derive the ME may result in an inaccurate estimation of the magnitude of the $S(\omega)$ compared to the HEOM, especially in the high-frequency range.

chemical potentials $\mu_L = -\mu_R = e\Phi/2$ for the left and right fermionic reservoirs, and the temperature T common to all reservoirs. All values of the parameters are summarized in Table 1. Using these parameters as input in `HierarchicalEOM.jl`, we can construct the HEOMLS matrix $\hat{\mathcal{M}}$ and solve for the stationary states of all the ADOs.

We now compute the spectra (see Methods) of both the charge (density of states; DOS) and cavity (power spectral density; PSD) using the built-in function in `HierarchicalEOM.jl`, and plot the results in Fig. 4. Here, we consider the bias voltage $\Phi = 6$, such that the site energy of the charge system aligns with the Fermi-level of one of the fermionic reservoirs, increasing the likelihood of electron tunneling to the fermionic reservoirs. We first discuss the DOS of the charge system, as shown in Fig. 4a. As the coupling strength g between the charge and the cavity increases, the DOS of the charge system decreases and its peak shifts towards lower energies. This can be attributed to the energy shift induced by the cavity field on the electronic level. To explore the charge-cavity interaction, we now analyze the PSD of the cavity system, see Fig. 4b. The PSD of the cavity exhibits a single

large peak at $\omega = \Gamma = \omega_c$ for $g \in (0, 0.5]$, indicating that the cavity resonance is largely unaffected by the electronic system, but that it does induce additional broadening.

It is interesting to further compare the differences between the HEOM and a perturbative master equation (ME) approach. We do this by reducing the interaction strength $\Delta = 0.01\omega_c$ between the cavity and its bosonic reservoir to a weak-coupling regime. We model the interaction between the system and bosonic reservoir with a Born-Markov master equation⁸⁰ using the Lindbladian given in Eq. (21). More specifically, we replace the bosonic part of HEOM with the following term

$$J_\beta(\omega_c) \left[\left(n_\beta^{\text{eq}}(\omega_c) + 1 \right) \tilde{\mathcal{J}}(a) + n_\beta^{\text{eq}}(\omega_c) \tilde{\mathcal{J}}(a^\dagger) \right] \rho_{\text{iq}}^{(0,n,p)}(t) \quad (24)$$

on the right-hand side of Eq. (16). Here, J_β is the spectral density of the bosonic bath (assumed to be Drude-Lorentz in this case) and n_β^{eq} is the Bose-Einstein distribution. In this specific scenario, it is evident that the outcomes derived from the master equation approach may overestimate the magnitude when compared to the HEOM results in the high-frequency range of the PSD. The origin




Fig. 5 Effect of the electron-cavity coupling on the conductance between the charge system and fermionic reservoirs. The black dashed, red dotted, and blue solid curves represent the conductance G for the charge-cavity coupling strength g at 0, 0.25Γ , and 0.5Γ , respectively. The conductance depends on the absolute value of the bias voltage Φ and reaches its maximum when the average energy of the charge system is equal to the Fermi-level of one of the fermionic reservoirs, i.e., when $e|\Phi|/2 = \epsilon$. As the electron-cavity coupling g increases, both the conductance G and the average energy of the charge system decrease, resulting in a wider splitting of the conductance peaks.

of this inconsistency can be attributed to the non-Markovian effect^{5,9,81,82} arising from the bosonic reservoir, characterized by the narrow band-width of the Drude-Lorentz spectral density ($W_\beta = 0.2$) we use to describe it; even though the coupling strength is weak, this narrow width implies the environment can still have memory. In addition, the master equation has limited capability in addressing the combined influence of electron-cavity interactions and electron-fermionic-reservoir hybridization. Nevertheless, these challenges can be effectively tackled using the HEOM approach.

Applying the same methodology as in the previous example, one can also calculate the conductance between the charge system and the fermionic reservoirs, as shown in Fig. 5. We begin by examining the case when the charge system is not coupled to the cavity ($g = 0$). In this scenario, due to the finite band-width of the fermionic reservoirs W_α and a site energy of the charge system $\epsilon = -3$ being far away from the Fermi-level, a local minimum of the conductance is observed at $\Phi = 0$. Additionally, the conductance increases with increasing $|\Phi|$, reaching its maximum value when the site energy ϵ coincides with the Fermi-level of one of the fermionic reservoirs, resulting in the splitting of the peaks in the conductance.

Conversely, when the charge is coupled to the cavity ($g > 0$), the average energy of the charge in the electronic system decreases away from the Fermi-levels of the fermionic reservoirs, resulting in a decrease in conductance. It should be noted that the maximum conductance will be achieved with larger $|\Phi|$, leading to a wider splitting of the conductance peak.

Capabilities of HierarchicalEOM.jl and comparison with other packages. Recent developments in the field of HEOM have led to an explosion of open source software, such as PHI⁵⁵, HEOM-QUICK²⁴, DM-HEOM⁵⁶, PyHEOM⁵⁷, and QuTiP-BoFin^{15,83,84}. These packages are written in different programming languages which have different capabilities. Thus, in Table 2, we briefly

compare their differences and similarities with respect to HierarchicalEOM.jl.

We first emphasize the advantage of specifying the importance threshold \mathcal{I}_{th} and estimating the importance for each ADOs with Eq. (20), which are not available for other packages listed in Table 2. The main computational complexity can be quantified by the total number of ADOs because it directly affects the size of $\hat{\mathcal{M}}$. The importance criterion $\mathcal{I}(\rho_{j|q}^{(m,n,p)}) \geq \mathcal{I}_{\text{th}}$ allows us to only consider the ADOs which affects the dynamics more, and thus, reduce the size of $\hat{\mathcal{M}}$. Table 3 summarizes the number of ADOs with respect to different truncation level n_{max} and importance thresholds \mathcal{I}_{th} . We observe a reduction in the number of auxiliary density operators when \mathcal{I}_{th} increases.

We further discuss the consistency of the importance criterion and consider the parameter settings in example 1 (see Table 1). In this case, we first increase the truncation level until we find out that the values of electronic current converges at $n_{\text{max}} = 4$, which requires 396,607 ADOs (as shown in Table. 3) to describe the dynamics. The effects on the convergence of the electronic current while using the importance criterion are shown in Fig. 6. For the equilibrium case ($\Phi = 0$), one can observe that the electronic current already converged at $\mathcal{I}_{\text{th}} = 10^{-5}$. However, by increasing the bias voltage to non-equilibrium $\Phi > 0$, the results obtained with a higher \mathcal{I}_{th} show larger deviation from the converged results. The results converge at $\mathcal{I}_{\text{th}} = 10^{-7}$ and require only 49,713 ADOs (as shown in Table. 3). For too large thresholds $\mathcal{I}_{\text{th}} \geq 10^{-4}$, we observe unphysical negative electronic currents or even fail to obtain the stationary states of the ADOs.

We now compare the performance in constructing HEOMLS matrix $\hat{\mathcal{M}}$, solving time evolution (TE) of ADOs, and solving the stationary states (SS) of ADOs between HierarchicalEOM.jl and QuTiP-BoFin^{15,83,84}. This choice is motivated by the fact that QuTiP-BoFin served as an inspiration and guide for the development of this package, and thus has similar features, and by the fact that they are both purely written in high-level languages (Python and Julia). For this comparison, we consider the same setting in example 1 (see Table 1). The benchmark result of constructing $\hat{\mathcal{M}}$ are shown in Fig. 7a. HierarchicalEOM.jl not only improves the computational time on single-thread processing but also supports multi-threading to make the process of constructing $\hat{\mathcal{M}}$ faster. Note that we only consider the importance threshold $\mathcal{I}_{\text{th}} = 0.0$ in this benchmark because QuTiP-BoFin does not support this functionality at this time. On top of the speed up in constructing $\hat{\mathcal{M}}$, we emphasize that HierarchicalEOM.jl is also faster in computing the time evolution and stationary states of the ADOs, as shown in Fig. 7(b), largely because of underlying performance benefits of the numerical libraries in Julia^{39–41}. For example, DifferentialEquations.jl³⁹ provides a unified user interface to solve the differential equations with various choices of solver. HierarchicalEOM.jl wrapped a subset of the functions in DifferentialEquations.jl for solving TE of the ADOs, and users can optimize their solver choice depending on the problem. We have also wrapped some of the functions in LinearSolver.jl⁴⁰ for users to solve the stationary states of ADOs and the spectrum of the system with different solvers. These functionalities improve both runtime and accuracy compared to the libraries currently available in Python.

Conclusion

In conclusion, the HierarchicalEOM.jl software package provides a user-friendly and efficient tool for simulating complex open quantum systems, including non-Markovian effects due to

Table 2 Comparison between different open source HEOM packages.

| Package Name | PHI ⁵⁵ | HEOM-QUICK ²⁴ | DM-HEOM ⁵⁶ | PyHEOM ⁵⁷ | QuTiP-BoFin ¹⁵ | HierarchicalEOM.jl |
|--------------------|-------------------|--------------------------|-----------------------|----------------------|---------------------------|--------------------|
| Language | Fortran | Fortran | C/C++ | C++/Python | Python | Julia |
| Documentation | ✓ | ✓ | ✓ | × | ✓ | ✓ |
| RunTests | × | × | × | × | ✓ | ✓ |
| Bosonic Bath | ✓ | × | ✓ | ✓ | ✓ | ✓ |
| Fermionic Bath | × | ✓ | × | × | ✓ | ✓ |
| Hybrid Bath | × | × | × | × | ✓ | ✓ |
| T-D system support | × | ✓ | × | × | ✓ | ✓ |
| Parity support | × | ✓ | × | × | × | ✓ |
| Importance support | × | × | × | × | × | ✓ |

For each package, the check (cross) mark represents the presence (absence) of support for the listed functionalities. Language specifies the programming language. Documentation labels the availability of instructions for installing and running the code. RunTests describes the availability of a suite of tests to ensure the correct functionality of the package. T-D system describes the availability of the system Hamiltonian H_s to be time-dependent (T-D). Parity describes the availability of constructing the HEOMLS matrices for even- or odd-parity auxiliary density operators.

Table 3 Number of auxiliary density operators for different truncation values of the fermionic hierarchy level and different thresholds of the importance value.

| $n_{\max} \setminus \mathcal{I}_{\text{th}}$ | 10^{-3} | 10^{-4} | 10^{-5} | 10^{-6} | 10^{-7} | 10^{-8} | 10^{-9} | 10^{-10} | 0.0 |
|--|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|------------|
| 1 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57 |
| 2 | 249 | 873 | 1193 | 1421 | 1569 | 1597 | 1597 | 1597 | 1597 |
| 3 | 305 | 1489 | 4241 | 12,713 | 18,933 | 23,693 | 27,161 | 28,645 | 29,317 |
| 4 | 305 | 1489 | 5011 | 18,901 | 49,713 | 126,715 | 205,803 | 274,249 | 396,607 |
| 5 | 305 | 1489 | 5011 | 19,013 | 55,173 | 170,297 | 418,589 | 931,551 | 4,216,423 |
| 6 | 305 | 1489 | 5011 | 19,013 | 55,201 | 171,557 | 444,979 | 1,137,239 | 36,684,859 |

Here, we use the parameters from example 1 (see Table 1), where n_{\max} is the truncation tier of the fermionic hierarchy and \mathcal{I}_{th} is the importance threshold. Note that we only neglect the ADOs in the second and higher levels ($n \geq 2$) whose importance value is smaller than \mathcal{I}_{th} .




Fig. 6 Convergence of the electronic current with respect to the importance threshold. Here, we consider the parameters setting in example 1 (see Table 1) and show the electronic current I_L (from the left-hand side fermionic reservoir into the system) as a function of bias voltage Φ where the red dashed, green dash-dotted, blue solid curves, and black dots correspond to the importance threshold \mathcal{I}_{th} at 10^{-5} , 10^{-6} , 10^{-7} , and 0, respectively. The values of electronic current converge when $\mathcal{I}_{\text{th}} = 10^{-7}$, which reduces the number of ADOs from 396,607 ($\mathcal{I}_{\text{th}} = 0$) to 49,713, as shown in Table 3.

non-perturbative interaction with one, or multiple, environments. The following characteristics define the package:

- It decreases the number of ADOs and the size of the HEOMLS matrices by allowing the users to specify importance thresholds^{23,26}.

- It provides built-in functions to calculate the spectrum for both bosonic and fermionic systems based on `LinearSolve.jl`⁴⁰.
- It provides a dictionary translating the index of the ADOs to the corresponding bath exponential-expansion series, and vice versa. This allows an easy way to interrogate the ADOs to gain access to bath properties, like electronic and heat currents.
- Users can choose to construct the HEOMLS matrices within even- or odd-parity. It depends on the parity of the operator which the HEOMLS matrix is acting on.
- It provides built-in functions for the users to further add Lindbladians to the HEOMLS matrices while keeping the size of HEOMLS matrices unchanged (it does not extend the ADO space). The Lindbladian describes the dissipative interactions (under Born-Markov approximation⁸⁰) between the system and extra environments.

Moreover, `HierarchicalEOM.jl` also takes advantage of other available packages for the following features:

- It is written in Julia.
- It supports different choices of spectral densities and spectral decomposition methods to accurately compute bath correlation functions.
- It supports time-dependent system Hamiltonians.
- It constructs the HEOMLS matrices for different types (bosonic, fermionic, or hybrid) of baths.
- The HEOMLS matrices are constructed using multi-threading.
- It provides different methods based on `DifferentialEquations.jl`³⁹, `LinearSolve.jl`⁴⁰, and `FastExp.jl`⁴¹ to compute the dynamics and stationary states of all ADOs.




Fig. 7 Benchmark of the performance between HierarchicalEOM.jl and QuTiP-BoFiN. We consider the parameters used in example 1 (see Table 1). Note that QuTiP-BoFiN^{15,83,84} currently does not support the estimation of importance for ADOs, and thus, we set $\mathcal{I}_{\text{th}} = 0$ here. The benchmark was done on a workstation with two Intel(R) Xeon(R) Silver 4110 @2.10GHz CPUs and 256 GB memory. The software versions were QuTiP 4.7.1, HierarchicalEOM.jl 0.1.0, Python 3.7.6, and Julia 1.8.2. The operating system was CentOS Linux 8.1.1911. In **a**, we show the total runtime versus truncation level n_{\max} for the construction of the HEOMLS matrices $\hat{\mathcal{M}}$ using QuTiP-BoFiN (red crosses) and HierarchicalEOM.jl (black triangles and blue circles represent 1 and 8 threads used during the process of constructing the matrices, respectively). Note that QuTiP-BoFiN currently only supports single-thread process for the construction of $\hat{\mathcal{M}}$. In **b**, we show the total runtime versus truncation level n_{\max} for calculating the stationary states (SS), represented as dashed lines, and time evolution (TE), represented as solid lines, of the auxiliary density operators using QuTiP-BoFiN (red crosses and green squares) and HierarchicalEOM.jl (blue circles and black triangles).

We exemplified the functionalities of HierarchicalEOM.jl using two physics examples where we observed order of magnitudes speedups with respect to QuTiP-BoFiN in constructing HEOMLS matrices and solving dynamics and stationary states for all ADOs. As a result, we believe that HierarchicalEOM.jl will be a valuable tool for researchers working in different fields such as quantum biology, quantum optics, quantum thermodynamics, quantum information, quantum transport, and condensed matter physics.

We further plan to extend the software for future releases in two different directions. First, we aim to add fitting routines to the decomposition methods used to compute correlation functions, thereby expanding the domain of spectral densities supported by the HEOM approach. Second, we aim to improve the computational efficiency of the package by reducing the complexity of the HEOM (for example imposing Hermiticity^{23,24}) and by including support for distributed computing on clusters and GPU computing.

Methods

Table of symbols. Table 4 summarizes the most relevant symbols used throughout the article.

Fermion correlation function with Lorentzian spectral density.

In this subsection, we present an example for the decomposition of a α -fermionic reservoir correlation function as a sum of N_α -exponential terms. In particular, we focus on the following Lorentzian spectral density

$$J_\alpha(\omega) = \frac{\Gamma_\alpha W_\alpha^2}{(\omega - \mu_\alpha)^2 + W_\alpha^2}, \quad (25)$$

where Γ_α represents the coupling strength between system and α -fermionic reservoir with band-width W_α and chemical potential μ_α . We now express the Fermi-Dirac distribution $n_\alpha^{\text{eq}}(x) = \{\exp(x) + 1\}^{-1}$ as a series by employing the Padé decomposition⁵¹ (which has been reported^{15,52} to enjoy a faster convergence than the Matsubara one⁵⁰) to obtain

$$n_\alpha^{\text{eq}}(x) \approx n_\alpha^{\text{Padé}}(x) = \frac{1}{2} - \sum_{h=2}^{N_\alpha} \frac{2\kappa_h x}{x^2 + \zeta_h^2}, \quad (26)$$

where the parameters κ_h and ζ_h are described in previous work⁵¹ and depend on the choice of N_α . Thus, the integration in Eq. (10)

can be analytically performed using the residue theorem and results in Eq. (12), which explicitly reads

$$C_\alpha^\nu(\tau) \approx \sum_{h=1}^{N_\alpha} \eta_{\alpha,h}^\nu \exp(-\gamma_{\alpha,h}^\nu \tau) \quad (27)$$

with

$$\begin{aligned} \eta_{\alpha,1}^\nu &= \frac{\Gamma_\alpha W_\alpha}{2} n_\alpha^{\text{Padé}}\left(\frac{iW_\alpha}{k_B T}\right), \\ \gamma_{\alpha,1}^\nu &= W_\alpha - \nu i\mu_\alpha, \\ \eta_{\alpha,h \neq 1}^\nu &= -i\kappa_h k_B T \cdot \frac{\Gamma_\alpha W_\alpha^2}{-(\zeta_h k_B T)^2 + W_\alpha^2}, \\ \gamma_{\alpha,h \neq 1}^\nu &= \zeta_h k_B T - \nu i\mu_\alpha. \end{aligned} \quad (28)$$

Boson correlation function with Drude-Lorentz spectral density. In this subsection, we present an example for the decomposition of a β -bosonic reservoir correlation function as a sum of N_β -exponential terms. In particular, we focus on the following Drude-Lorentz spectral density

$$J_\beta(\omega) = \frac{4\Delta_\beta W_\beta \omega}{\omega^2 + W_\beta^2}, \quad (29)$$

in which Δ_β represents the coupling strength between the system and the β -bosonic reservoir with band-width W_β . Similarly as in the fermionic case, we can now express the Bose-Einstein distribution $n_\beta^{\text{eq}}(x) = \{\exp(x) - 1\}^{-1}$ as a series by employing the Padé decomposition⁵¹, to obtain

$$n_\beta^{\text{eq}}(x) \approx n_\beta^{\text{Padé}}(x) = \frac{1}{x} - \frac{1}{2} + \sum_{l=2}^{N_\beta} \frac{2\kappa_l x}{x^2 + \zeta_l^2}, \quad (30)$$

where the parameters κ_l and ζ_l are described in previous work⁵¹ and depend on the choice of N_β . Using this decomposition, the integration in Eq. (11) can be analytically performed using the residue theorem and results in Eq. (13), which explicitly reads

$$C_\beta(\tau) \approx \sum_{l=1}^{N_\beta} \xi_{\beta,l} \exp(-\chi_{\beta,l} \tau) \quad (31)$$

Table 4 List of symbols.

| Symbol | Description |
|--|--|
| σ_b | specification of different quantum numbers of the bosonic system |
| σ_f | specification of different quantum numbers of the fermionic system |
| β | specification of different bosonic baths |
| α | specification of different fermionic baths |
| $[\cdot, \cdot]_-$ | commutator |
| $[\cdot, \cdot]_+$ | anti-commutator |
| $\nu \in \{+1, -1\}$ | presence (+1) or absence (-1) of Hermitian conjugation |
| $\bar{\nu}$ | opposite of ν : $\bar{\nu} \equiv -\nu$ |
| $p \in \{+, -\}$ | even-parity (+) or odd-parity (-) of the operator |
| $u \in \{\mathbb{R}, \mathbb{I}\}$ | Real (\mathbb{R}) or imaginary (\mathbb{I}) part of bosonic correlation function |
| $C(t_1, t_2)$ | two time correlation function of the bath |
| $n^{\text{eq}}(\omega)$ | Fermi-Dirac or Bose-Einstein equilibrium distribution |
| $J(\omega)$ | spectral density of the bath |
| $\hat{\mathcal{G}}[\cdot]$ | superoperator which propagates all auxiliary density operators (ADOs) |
| $\hat{\mathcal{M}}[\cdot]$ | HEOM Liouvillian superoperator (matrix) |
| $\tilde{\mathcal{J}}(F)[\cdot]$ | Lindbladian superoperator with jump operator F |
| $\rho_{\mathbf{j}\mathbf{q}}^{(m,n,p)}$ | auxiliary density operator (ADO) |
| $\rho_{\mathbf{j}}^{(0,0,+)}$ | reduced density operator |
| m | bosonic hierarchy level |
| n | fermionic hierarchy level |
| \mathbf{j} | vector of bosonic multi-index ensembles: $[j_m, \dots, j_1]$ |
| \mathbf{q} | vector of fermionic multi-index ensembles: $[q_n, \dots, q_1]$ |
| $\mathcal{I}(\rho_{\mathbf{j}\mathbf{q}}^{(m,n,p)})$ | importance value of a given auxiliary density operator (ADO) |
| \mathcal{I}_{th} | threshold of importance value |
| $A(\omega)$ | spectrum of fermionic system (density of states, DOS) |
| $S(\omega)$ | spectrum of bosonic system (power spectral density, PSD) |
| I_α | electronic current from α -fermionic bath into the system |
| G | conductance |

We summarize the most relevant symbols used in this article.

with

$$\begin{aligned} \xi_{\beta,1} &= \Delta_\beta W_\beta \left[-i + \cot\left(\frac{W_\beta}{2k_B T}\right) \right], \\ \chi_{\beta,1} &= W_\beta, \\ \xi_{\beta,l \neq 1} &= -2\kappa_l k_B T \cdot \frac{2\Delta_\beta W_\beta \cdot \zeta_l k_B T}{-(\zeta_l k_B T)^2 + W_\beta^2}, \\ \chi_{\beta,l \neq 1} &= \zeta_l k_B T. \end{aligned} \tag{32}$$

Contrary to the fermionic case, we note that, here, the correlation function does not have to be further decomposed into its real and imaginary part since $\chi_{\beta,l} = \chi_{\beta,l}^* \forall l$.

Numerical computation of the spectrum. In this subsection, we briefly summarize how to numerically compute the spectrum associated with the system degree of freedom. Previous work¹³ showed that the spectrum can be evaluated either in time or frequency domain. `HierarchicalEOM.jl` provides a built-in function which performs the calculation in frequency domain. In the following, we first focus on the bosonic case (to compute the power spectral density) and, next, on fermionic case (to compute the density of states).

In order to compute the power spectral density in the time-domain, we write the system correlation function in terms of the propagator $\hat{\mathcal{G}}(t) = \exp(\hat{\mathcal{M}}t)$, for $t > 0$, when the system

Hamiltonian is time-independent. The power spectral density associated with σ_b can be obtained as

$$\begin{aligned} \pi S_{\sigma_b}(\omega) &= \text{Re} \left\{ \int_0^\infty dt \left\langle a_{\sigma_b}^\dagger(t) a_{\sigma_b}(0) \right\rangle^* e^{i\omega t} \right\} \\ &= \text{Re} \left\{ \int_0^\infty dt \left\langle a_{\sigma_b}^\dagger(t) a_{\sigma_b}(0) \right\rangle e^{-i\omega t} \right\} \\ &= \text{Re} \left\{ \int_0^\infty dt \left\langle a_{\sigma_b}^\dagger e^{\hat{\mathcal{M}}t} a_{\sigma_b} \right\rangle e^{-i\omega t} \right\} \\ &= \text{Re} \left\{ \int_0^\infty dt \left\langle a_{\sigma_b}^\dagger e^{(\hat{\mathcal{M}}-i\omega)t} a_{\sigma_b} \right\rangle \right\} \\ &= \text{Re} \left\{ -\left\langle a_{\sigma_b}^\dagger (\hat{\mathcal{M}} - i\omega)^{-1} a_{\sigma_b} \right\rangle \right\} \\ &= -\text{Re} \left\{ \text{Tr} \left[a_{\sigma_b}^\dagger (\hat{\mathcal{M}} - i\omega)^{-1} a_{\sigma_b} \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)} \right] \right\} \\ &= -\text{Re} \left\{ \text{Tr} \left[a_{\sigma_b}^\dagger \mathbf{x} \right] \right\}, \end{aligned} \tag{33}$$

where a half-Fourier transform has been introduced in the fifth line. To determine \mathbf{x} at a fixed frequency ω , one can express all of the ADOs in a vector form and then solve the linear problem $\mathbf{A}\mathbf{x} = \mathbf{b}$ (where $\mathbf{A} = \hat{\mathcal{M}} - i\omega$ and $\mathbf{b} = a_{\sigma_b} \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)}$) using the package `LinearSolve.jl`⁴⁰. We note that, while in principle all the ADOs are required to solve for \mathbf{x} , only the reduced density operator ($m=0$ and $n=0$) is considered when taking the final trace.

In order to compute the density of states, we start from the retarded Green's function¹³

$$G_{\sigma_f}^R(t) = -i\Theta(t) \left\{ \left\langle d_{\sigma_f}(t) d_{\sigma_f}^\dagger(0) \right\rangle + \left\langle d_{\sigma_f}^\dagger(t) d_{\sigma_f}(0) \right\rangle^* \right\}, \tag{34}$$

in which the Heaviside function $\Theta(t)$ is needed to impose causality. Similarly to the bosonic case, the density of states associated with σ_f can be written in compact form as

$$\begin{aligned} \pi A_{\sigma_f}(\omega) &= -\text{Im} \int_{-\infty}^\infty G_{\sigma_f}^R(t) e^{i\omega t} dt \\ &= \text{Re} \left\{ \int_0^\infty dt \left[\left\langle d_{\sigma_f}(t) d_{\sigma_f}^\dagger(0) \right\rangle + \left\langle d_{\sigma_f}^\dagger(t) d_{\sigma_f}(0) \right\rangle^* \right] e^{i\omega t} \right\} \\ &= -\text{Re} \left\{ \text{Tr} \left[d_{\sigma_f} (\hat{\mathcal{M}} + i\omega)^{-1} d_{\sigma_f}^\dagger \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)} \right] \right. \\ &\quad \left. + \text{Tr} \left[d_{\sigma_f}^\dagger (\hat{\mathcal{M}} - i\omega)^{-1} d_{\sigma_f} \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)} \right] \right\} \\ &= -\text{Re} \left\{ \text{Tr} \left[d_{\sigma_f} \mathbf{x}_+ \right] + \text{Tr} \left[d_{\sigma_f}^\dagger \mathbf{x}_- \right] \right\}, \end{aligned} \tag{35}$$

where \mathbf{x}_+ is determined by solving

$$(\hat{\mathcal{M}} + i\omega) \mathbf{x}_+ = d_{\sigma_f}^\dagger \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)}, \tag{36}$$

and \mathbf{x}_- is determined by solving

$$(\hat{\mathcal{M}} - i\omega) \mathbf{x}_- = d_{\sigma_f} \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)}. \tag{37}$$

Here, the HEOMLS matrix $\hat{\mathcal{M}}$ acts on the odd-parity ($p = -$) space, compatibly with the parity of both the operators $d_{\sigma_f} \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)}$ and $d_{\sigma_f}^\dagger \rho_{\mathbf{j}\mathbf{q}}^{(m,n,+)}$. As for the computation of the power spectral density, we only need the reduced density operator ($m=0$ and $n=0$) to take the final trace operation.

Electronic current. In this subsection, we discuss how to compute an environmental observable: the electronic current. Within the influence functional approach^{21,23}, the expectation value of the electronic current from the α -fermionic bath into the system can be written in terms of the first-level-fermionic ($n=1$)

auxiliary density operators, namely

$$\begin{aligned} \langle I_\alpha(t) \rangle &= (-e) \frac{d \langle \mathcal{N}_\alpha \rangle}{dt} \\ &= ie \sum_{\nu, h, \sigma_f} (-1)^{\delta_{\nu,-}} \text{Tr} \left[d_{\sigma_f}^{\bar{\nu}} \rho_{\{\alpha, \nu, h, \sigma_f\}}^{(0,1,+)}(t) \right], \end{aligned} \quad (38)$$

where e represents the value of the elementary charge, and $\mathcal{N}_\alpha = \sum_k c_{\alpha,k}^\dagger c_{\alpha,k}$ is the occupation number operator for the α -fermionic bath. To compute this quantity, it is possible to first use `HierarchicalEOM.jl` to obtain the stationary states for the ADOs $\rho_{\text{jq}}^{(m,n,+)}(t)$, and then use the hierarchy dictionary [as shown in Fig. 1e] to identify all the zeroth-level-bosonic-and-first-level-fermionic ($m=0$ and $n=1$) ADOs and their corresponding indices α , ν , h , and σ_f to be used in Eq. (38).

Data availability

The detailed information for reproducing the numerical data in both examples is available through a public GitHub repo (<https://github.com/ytdHuang/HierarchicalEOM.jl-Examples>). All other data are available upon reasonable request.

Code availability

The `HierarchicalEOM.jl` package is available through a public GitHub repo (<https://github.com/NCKU-QFort/HierarchicalEOM.jl>). It is also registered in the Julia package registry and can be installed by the Julia package manager. Moreover, detailed information (documentation and other examples) is available through a public website (<https://ncku-qfort.github.io/HierarchicalEOM.jl>).

Received: 13 June 2023; Accepted: 10 October 2023;

Published online: 23 October 2023

References

- Zwanzig, R. Ensemble method in the theory of irreversibility. *J. Chem. Phys.* **33**, 1338 (1960).
- Feynman, R. & Vernon, F. The theory of a general quantum system interacting with a linear dissipative system. *Ann. Phys.* **24**, 118 (1963).
- Caldeira, A. & Leggett, A. Path integral approach to quantum Brownian motion. *Physica A* **121**, 587 (1983).
- Hedegård, P. & Caldeira, A. O. Quantum dynamics of a particle in a Fermionic environment. *Phys. Scripta* **35**, 609 (1987).
- Chen, H.-B., Lambert, N., Cheng, Y.-C., Chen, Y.-N. & Nori, F. Using non-Markovian measures to evaluate quantum master equations for photosynthesis. *Sci. Rep.* **5**, 12753 (2015).
- Tanimura, Y. & Kubo, R. Time evolution of a quantum system in contact with a nearly Gaussian-Markoffian noise bath. *J. Phys. Soc. Jpn.* **58**, 101 (1989).
- Tanimura, Y. Nonperturbative expansion method for a quantum system coupled to a harmonic-oscillator bath. *Phys. Rev. A* **41**, 6676 (1990).
- Bulla, R., Costi, T. A. & Pruschke, T. Numerical renormalization group method for quantum impurity systems. *Rev. Mod. Phys.* **80**, 395 (2008).
- Zhang, W.-M., Lo, P.-Y., Xiong, H.-N., Tu, M. W.-Y. & Nori, F. General non-Markovian dynamics of open quantum systems. *Phys. Rev. Lett.* **109**, 170402 (2012).
- Strasberg, P., Schaller, G., Lambert, N. & Brandes, T. Nonequilibrium thermodynamics in the strong coupling and non-Markovian regime based on a reaction coordinate mapping. *New J. Phys.* **18**, 073007 (2016).
- Brenes, M. et al. Tensor-network method to simulate strongly interacting quantum thermal machines. *Phys. Rev. X* **10**, 031040 (2020).
- Sowa, J. K., Lambert, N., Seideman, T. & Gauger, E. M. Beyond Marcus theory and the Landauer-Büttiker approach in molecular junctions. II. A self-consistent Born approach. *J. Chem. Phys.* **152**, 064103 (2020).
- Li, Z. et al. Hierarchical Liouville-space approach for accurate and universal characterization of quantum impurity systems. *Phys. Rev. Lett.* **109**, 266403 (2012).
- Tanimura, Y. Numerically “exact” approach to open quantum dynamics: The hierarchical equations of motion (HEOM). *J. Chem. Phys.* **153**, 020901 (2020).
- Lambert, N. et al. QuTiP-BoFiN: A bosonic and fermionic numerical hierarchical-equations-of-motion library with applications in light-harvesting, quantum control, and single-molecule electronics. *Phys. Rev. Res.* **5**, 013181 (2023).
- Lambert, N., Ahmed, S., Cirio, M. & Nori, F. Modelling the ultra-strongly coupled spin-boson model with unphysical modes. *Nat. Commun.* **10**, 3721 (2019).
- Fay, T. P. & Limmer, D. T. Coupled charge and energy transfer dynamics in light harvesting complexes from a hybrid hierarchical equations of motion approach. *J. Chem. Phys.* **157**, 174104 (2022).
- Ma, J., Sun, Z., Wang, X. & Nori, F. Entanglement dynamics of two qubits in a common bath. *Phys. Rev. A* **85**, 062323 (2012).
- Kato, A. & Tanimura, Y. Quantum heat current under non-perturbative and non-Markovian conditions: Applications to heat machines. *J. Chem. Phys.* **145**, 224105 (2016).
- Chen, X.-Y. et al. Global correlation and local information flows in controllable non-Markovian open quantum dynamics. *npj Quantum Inf.* **8**, 22 (2022).
- Jin, J., Zheng, X. & Yan, Y. Exact dynamics of dissipative electronic systems and quantum transport: Hierarchical equations of motion approach. *J. Chem. Phys.* **128**, 234703 (2008).
- Ishizaki, A. & Fleming, G. R. Unified treatment of quantum coherent and incoherent hopping dynamics in electronic energy transfer: Reduced hierarchy equation approach. *J. Chem. Phys.* **130**, 234111 (2009).
- Härtle, R., Cohen, G., Reichman, D. R. & Millis, A. J. Decoherence and lead-induced interdot coupling in nonequilibrium electron transport through interacting quantum dots: A hierarchical quantum master equation approach. *Phys. Rev. B* **88**, 235426 (2013).
- Ye, L. et al. HEOM-quick: a program for accurate, efficient, and universal characterization of strongly correlated quantum impurity systems. *WIREs Comput. Mol. Sci.* **6**, 608 (2016).
- Schinabeck, C., Härtle, R. & Thoss, M. Hierarchical quantum master equation approach to electronic-vibrational coupling in nonequilibrium transport through nanosystems: Reservoir formulation and application to vibrational instabilities. *Phys. Rev. B* **97**, 235429 (2018).
- Bätge, J., Ke, Y., Kaspar, C. & Thoss, M. Nonequilibrium open quantum systems with multiple bosonic and fermionic environments: A hierarchical equations of motion approach. *Phys. Rev. B* **103**, 235413 (2021).
- Bezanson, J., Karpinski, S., Shah, V. B. & Edelman, A. Julia: A fast dynamic language for technical computing. *arXiv preprint arXiv:1209.5145*, (2012).
- Bezanson, J., Edelman, A., Karpinski, S. & Shah, V. B. Julia: A fresh approach to numerical computing. *SIAM Rev.* **59**, 65 (2017).
- Kernighan, B. W. & Ritchie, D. M. *The C programming language* (Prentice Hall Professional Technical Reference, 2006).
- Flanagan, D. & Matsumoto, Y. *The Ruby Programming Language* (O’Reilly Media, Inc., 2007).
- Van Rossum, G. & Drake, F. L. *Python 3 Reference Manual* (CreateSpace, Scotts Valley, CA, 2009).
- R Core Team. *R: A Language and Environment for Statistical Computing* (R Foundation for Statistical Computing, Vienna, Austria, 2021).
- Higham, D. J. & Higham, N. J. *MATLAB Guide* (Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2017).
- Lattner, C. & Adve, V. LLVM: a compilation framework for lifelong program analysis & transformation. In *International Symposium on Code Generation and Optimization, 2004. CGO 2004.*, 75, <https://doi.org/10.1109/CGO.2004.1281665> (2004).
- Chen, H. & Lidar, D. A. Hamiltonian open quantum system toolkit. *Commun. Phys.* **5**, 112 (2022).
- Krämer, S., Plankensteiner, D., Ostermann, L. & Ritsch, H. QuantumOptics.jl: A Julia framework for simulating open quantum systems. *Comput. Phys. Commun.* **227**, 109 (2018).
- Luo, X.-Z., Liu, J.-G., Zhang, P. & Wang, L. Yao.jl: Extensible, Efficient Framework for Quantum Algorithm Design. *Quantum* **4**, 341 (2020).
- Gawron, P., Kurzyk, D. & Pawela, Ł. QuantumInformation.jl—A Julia package for numerical computation in quantum information theory. *PLOS ONE* **13**, e0209358 (2018).
- Rackauckas, C. & Nie, Q. DifferentialEquations.jl – A performant and feature-rich ecosystem for solving differential equations in Julia. *J. Open Res. Software* **5**, 15 (2017).
- Kimmerer, W., Puri, V. & Rackauckas, C. Linearsolve.jl. <https://github.com/SciML/LinearSolve.jl>.
- Hogben, H., Krzysztyniak, M., Charnock, G., Hore, P. & Kuprov, I. Spinach - A software library for simulation of spin dynamics in large spin systems. *J. Magn. Reson.* **208**, 179 (2011).
- Cirio, M., De Liberato, S., Lambert, N. & Nori, F. Ground state electroluminescence. *Phys. Rev. Lett.* **116**, 113601 (2016).
- Stockklauser, A. et al. Strong coupling cavity QED with gate-defined double quantum dots enabled by a high impedance resonator. *Phys. Rev. X* **7**, 011030 (2017).

44. Gustafsson, M. V. et al. Propagating phonons coupled to an artificial atom. *Science* **346**, 207 (2014).
45. Manenti, R. et al. Circuit quantum acoustodynamics with surface acoustic waves. *Nat. Commun.* **8**, 975 (2017).
46. Iorsh, I., Poshakinskiy, A. & Poddubny, A. Waveguide quantum optomechanics: Parity-time phase transitions in ultrastrong coupling regime. *Phys. Rev. Lett.* **125**, 183601 (2020).
47. Benz, F. et al. Single-molecule optomechanics in “picocavities”. *Science* **354**, 726 (2016).
48. Kuo, P. C. et al. Collectively induced exceptional points of quantum emitters coupled to nanoparticle surface plasmons. *Phys. Rev. A* **101**, 013814 (2020).
49. Cirio, M., Kuo, P. C., Chen, Y. N., Nori, F. & Lambert, N. Canonical derivation of the fermionic influence superoperator. *Phys. Rev. B* **105**, 035121 (2022).
50. Shi, Q., Chen, L., Nan, G., Xu, R.-X. & Yan, Y. Efficient hierarchical Liouville space propagator to quantum dissipative dynamics. *J. Chem. Phys.* **130**, 084105 (2009).
51. Hu, J., Luo, M., Jiang, F., Xu, R.-X. & Yan, Y. Padé spectrum decompositions of quantum distribution functions and optimal hierarchical equations of motion construction for quantum open systems. *J. Chem. Phys.* **134**, 244106 (2011).
52. Kuo, P.-C. et al. Kondo QED: The Kondo effect and photon trapping in a two-impurity Anderson model ultra-strongly coupled to light. *arXiv preprint arXiv:2302.01044* (2023).
53. Wenderoth, S., Bätge, J. & Härtle, R. Sharp peaks in the conductance of a double quantum dot and a quantum-dot spin valve at high temperatures: A hierarchical quantum master equation approach. *Phys. Rev. B* **94**, 121303 (2016).
54. Härtle, R., Cohen, G., Reichman, D. R. & Millis, A. J. Transport through an Anderson impurity: Current ringing, nonlinear magnetization, and a direct comparison of continuous-time quantum monte carlo and hierarchical quantum master equations. *Phys. Rev. B* **92**, 085430 (2015).
55. Strümpfer, J. & Schulten, K. Open quantum dynamics calculations with the hierarchy equations of motion on parallel computers. *J. Chem. Theory Comput.* **8**, 2808 (2012).
56. Kramer, T., Noack, M., Reinefeld, A., Rodríguez, M. & Zelinskyy, Y. Efficient calculation of open quantum system dynamics and time-resolved spectroscopy with distributed memory HEOM (DM-HEOM). *J. Comput. Chem.* **39**, 1779 (2018).
57. Ikeda, T. & Scholes, G. D. Generalization of the hierarchical equations of motion theory for efficient calculations with arbitrary correlation functions. *J. Chem. Phys.* **152**, 204101 (2020).
58. Velizhanin, K. A., Wang, H. & Thoss, M. Heat transport through model molecular junctions: A multilayer multiconfiguration time-dependent hartree approach. *Chem. Phys. Lett.* **460**, 325 (2008).
59. Kato, A. & Tanimura, Y. Quantum heat transport of a two-qubit system: Interplay between system-bath coherence and qubit-qubit coherence. *J. Chem. Phys.* **143**, 064107 (2015).
60. Song, L. & Shi, Q. Hierarchical equations of motion method applied to nonequilibrium heat transport in model molecular junctions: Transient heat current and high-order moments of the current operator. *Phys. Rev. B* **95**, 064308 (2017).
61. Sprinzak, D., Ji, Y., Heiblum, M., Mahalu, D. & Shtrikman, H. Charge distribution in a Kondo-correlated quantum dot. *Phys. Rev. Lett.* **88**, 176805 (2002).
62. Keller, A. J. et al. Emergent SU(4) Kondo physics in a spin-charge-entangled double quantum dot. *Nat. Phys.* **10**, 145 (2013).
63. Le Hur, K. Quantum dots and the Kondo effect. *Nature* **526**, 203 (2015).
64. Park, J. et al. Coulomb blockade and the Kondo effect in single-atom transistors. *Nature* **417**, 722 (2002).
65. Wingreen, N. S. Quantum many-body effects in a single-electron transistor. *Science* **304**, 1258 (2004).
66. Yu, L. H. et al. Inelastic electron tunneling via molecular vibrations in single-molecule transistors. *Phys. Rev. Lett.* **93**, 266802 (2004).
67. Smith, A., Kim, M. S., Pollmann, F. & Knolle, J. Simulating quantum many-body dynamics on a current digital quantum computer. *npj Quantum Inf.* **5**, 106 (2019).
68. Wang, S., Zheng, X., Jin, J. & Yan, Y. Hierarchical Liouville-space approach to nonequilibrium dynamical properties of quantum impurity systems. *Phys. Rev. B* **88**, 035129 (2013).
69. Kouwenhoven, L. & Glazman, L. Revival of the Kondo effect. *Phys. World* **14**, 33 (2001).
70. Borzenets, I. V. et al. Observation of the Kondo screening cloud. *Nature* **579**, 210 (2020).
71. Smith, L. W. et al. Electrically controllable Kondo correlation in spin-orbit-coupled quantum point contacts. *Phys. Rev. Lett.* **128**, 027701 (2022).
72. van der Wiel, W. G. et al. Electron transport through double quantum dots. *Rev. Mod. Phys.* **75**, 1 (2002).
73. Bruhat, L. E. et al. Circuit QED with a quantum-dot charge qubit dressed by Cooper pairs. *Phys. Rev. B* **98**, 155313 (2018).
74. van Woerkom, D. J. et al. Microwave photon-mediated interactions between semiconductor qubits. *Phys. Rev. X* **8**, 041018 (2018).
75. Scarlino, P. et al. Coherent microwave-photon-mediated coupling between a semiconductor and a superconducting qubit. *Nat. Commun.* **10**, 3011 (2019).
76. Viennot, J. J., Delbecq, M. R., Dartiailh, M. C., Cottet, A. & Kontos, T. Out-of-equilibrium charge dynamics in a hybrid circuit quantum electrodynamics architecture. *Phys. Rev. B* **89**, 165404 (2014).
77. Bruhat, L. E. et al. Cavity photons as a probe for charge relaxation resistance and photon emission in a quantum dot coupled to normal and superconducting continua. *Phys. Rev. X* **6**, 021014 (2016).
78. Souquet, J. R., Woolley, M. J., Gabelli, J., Simon, P. & Clerk, A. A. Photon-assisted tunnelling with nonclassical light. *Nat. Commun.* **5**, 5562 (2014).
79. Kockum, A. F., Miranowicz, A., Liberato, S. D., Savasta, S. & Nori, F. Ultrastrong coupling between light and matter. *Nat. Rev. Phys.* **1**, 295 (2019).
80. Breuer, H.-P. & Petruccione, F. *The Theory of Open Quantum Systems* (Oxford University Press, 2007).
81. Chen, Y.-N., Chen, G.-Y., Liao, Y.-Y., Lambert, N. & Nori, F. Detecting non-Markovian plasmonic band gaps in quantum dots using electron transport. *Phys. Rev. B* **79**, 245312 (2009).
82. Xiong, H.-N., Lo, P.-Y., Zhang, W.-M., Feng, D. H. & Nori, F. Non-Markovian complexity in the quantum-to-classical transition. *Sci. Rep.* **5**, 13353 (2015).
83. Johansson, J., Nation, P. & Nori, F. QuTiP: An open-source Python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **183**, 1760 (2012).
84. Johansson, J., Nation, P. & Nori, F. QuTiP 2: A Python framework for the dynamics of open quantum systems. *Comput. Phys. Commun.* **184**, 1234 (2013).

Acknowledgements

The authors acknowledge fruitful discussions with Yuan-Ching Lin and Chung-Hsuan Tung. N.L. acknowledges the Information Systems Division, RIKEN, for the use of their facilities. M.C. acknowledges support from NSFC (Grants No. 12050410264 and No. 11935012) and NSAF (Grant No. U1930403). F.N. is supported in part by: Nippon Telegraph and Telephone Corporation (NTT) Research, the Japan Science and Technology Agency (JST) [via the Quantum Leap Flagship Program (Q-LEAP), and the Moonshot R&D Grant Number JPMJMS2061], Office of Naval Research (ONR), and the Asian Office of Aerospace Research and Development (AOARD) (via Grant No. FA2386-20-1-4069). F.N. and N.L. acknowledge the Foundational Questions Institute Fund (FQXi) via Grant No. FQXi-IAF19-06. YNC acknowledges the support of the National Center for Theoretical Sciences and the National Science and Technology Council, Taiwan (NSTC Grants No. 112-2123-M-006-001).

Author contributions

Y.-T.H. and P.-C.K. programmed the package, conceived the research, carried out the numerical calculations and wrote the manuscript. N.L., M.C. and S.C. contributed to analyzing the results and writing the manuscript. S.C. contributed to the development of the package. S.-L.Y. contributed to the numerical analysis of the results. F.N. and Y.-N.C. supervised the project. All authors participated in the discussion of the results and reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s42005-023-01427-2>.

Correspondence and requests for materials should be addressed to Neill Lambert, Mauro Cirio or Yueh-Nan Chen.

Peer review information *Communications Physics* thanks the anonymous reviewers for their contribution to the peer review of this work. A peer review file is available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023