



# Phagotrophic protists preserve antibiotic-resistant opportunistic human pathogens in the vegetable phyllosphere

Chenshuo Lin<sup>1,2</sup>, Li-Juan Li<sup>1,2</sup>, Kexin Ren<sup>1</sup>, Shu-Yi-Dan Zhou<sup>3</sup>, Alain Isabwe<sup>1</sup>, Le-Yang Yang<sup>1,2</sup>, Roy Neilson<sup>4</sup>, Xiao-Ru Yang<sup>1</sup>, Eddie Cytryn<sup>5</sup> and Yong-Guan Zhu<sup>1,6</sup>✉

© The Author(s) 2023

Food safety of leafy greens is an emerging public health issue as they can harbor opportunistic human pathogens (OHPs) and expose OHPs to consumers. Protists are an integral part of phyllosphere microbial ecosystems. However, our understanding of protist-pathogen associations in the phyllosphere and their consequences on public health remains poor. Here, we examined phyllosphere protists, human pathogen marker genes (HPMGs), and protist endosymbionts from four species of leafy greens from major supermarkets in Xiamen, China. Our results showed that *Staphylococcus aureus* and *Klebsiella pneumoniae* were the dominant human pathogens in the vegetable phyllosphere. The distribution of HPMGs and protistan communities differed between vegetable species, of which Chinese chive possessed the most diverse protists and highest abundance of HPMGs. HPMGs abundance positively correlated with the diversity and relative abundance of phagotrophic protists. Whole genome sequencing further uncovered that most isolated phyllosphere protists harbored multiple OHPs which carried antibiotic resistance genes, virulence factors, and metal resistance genes and had the potential to HGT. *Colpoda* were identified as key phagotrophic protists which positively linked to OHPs and carried diverse resistance and virulence potential endosymbiont OHPs including *Pseudomonas nitroreducens*, *Achromobacter xylooxidans*, and *Stenotrophomonas maltophilia*. We highlight that phyllosphere protists contribute to the transmission of resistant OHPs through internalization and thus pose risks to the food safety of leafy greens and human health. Our study provides insights into the protist-OHP interactions in the phyllosphere, which will help in food safety surveillance and human health.

ISME Communications; <https://doi.org/10.1038/s43705-023-00302-z>

## INTRODUCTION

Consuming raw or minimally processed leafy greens has become a popular healthy lifestyle choice for consumers [1]. However, the vegetable phyllosphere is a known reservoir of microbes [2] that can also host human pathogens without exhibiting any sign of spoilage or deterioration of the phyllosphere tissues, raising concerns about food safety [3]. Several outbreaks of severe food poisoning have been attributed to the consumption of raw leafy greens contaminated by pathogenic microorganisms, such as *Salmonella* spp., *E. coli*, and *Listeria monocytogenes* [4, 5]. According to the USDA Economic Research Service in 2016, consumers tend to believe that leafy greens produced with organic fertilizer (usually animal manure) without pesticides are safer and healthier than conventionally produced leafy greens [6]. However, they may ignore the potential health risks of organic vegetables, such as increased ARGs and human pathogens in leafy greens due to the application of animal manure [7]. Therefore, leafy greens especially organically produced ones are potential hot spots for the transmission of

food-borne opportunistic human pathogens (OHPs) [8–10] in light of the “One Health” concept.

Protists are one of the overlooked phyllosphere microbes that potentially affect human health by their potential pathogenicity [11] or shaping phyllosphere bacterial composition and function [12] (including bacterial human pathogens). Some phyllosphere protists are responsible for causing human diseases, including *Acanthamoeba* spp., *Giardia duodenalis*, *Entamoeba* spp., *Cryptosporidium*, etc. [13, 14]. More than 100 diverse environmental amoebae were reported to reduce the number of OHPs such as *Acinetobacter baumannii*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* [15, 16] by phagocytosis, killing OHPs intracellularly with toxic materials such as copper and arsenic [17]. In response to predation stress, specific pathogens evolve virulence (e.g., Shiga toxin and Violacein) [18–21] or mental resistance (e.g., copper resistance) [22] to resist digestion or internal shuttling to the contractile vacuole to replicate before being exocytosed to the environment [23]. Efflux pump systems closely associated with antibiotic resistance are also involved in

<sup>1</sup>Key Laboratory of Urban Environment and Health, Institute of Urban Environment, Chinese Academy of Sciences, 1799 Jimei Road, Xiamen 361021, China. <sup>2</sup>University of Chinese Academy of Sciences, 19A Yuquan Road, 100049 Beijing, China. <sup>3</sup>Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Xingke Road 723, Tianhe District, Guangzhou 510650, China. <sup>4</sup>Ecological Sciences, The James Hutton Institute, Dundee DD2 5DA Scotland, UK. <sup>5</sup>Department of Soil Chemistry, Plant Nutrition and Microbiology, Institute of Soil, Water and Environmental Sciences, The Volcani Institute, Agriculture Research Organization, 7528809 Rishon LeZion, Israel. <sup>6</sup>State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, 100085 Beijing, China. ✉email: [ygzhu@rcees.ac.cn](mailto:ygzhu@rcees.ac.cn)

Received: 19 April 2023 Revised: 21 August 2023 Accepted: 23 August 2023

Published online: 02 September 2023

grazing defense strategies. For example, the *C. jejuni* RND-type efflux pump, CmeABC, associated with multidrug resistance, may be involved in virulence and survival in *A. polyphaga* [24]. Meanwhile, protist-adapted pathogens may possess enhance environmental persistence, resistance, and transmission, thus enhancing the colonization in eukaryotic organisms of these pathogens [25]. For example, *Vibrio cholerae* that grew intracellularly in *Acanthamoeba castellanii* protists displayed *f1rA* mutations that increased the competitive fitness of *V. cholerae* and colonization potential in zebrafish [26]. Similarly, amoeba-adapted *L. pneumophila* showed increased resistance to antibiotics and chlorine and were more infectious than in vitro-grown *L. pneumophila* [27]. Therefore, protists can be “Trojan horses” (reservoir, shelters, and vectors) and “training grounds” that select for enhanced survival and virulence traits of OHPs [28–31]. To date, researches on phyllosphere protists in leafy greens have mainly focused on pathogens [32, 33]. Studies evaluating the endosymbiotic mechanisms of protists are generally based on co-cultivation assays between a single protist host and a specific pathogen [34, 35]. Therefore, our understanding of *in situ* pathogen-protist associations in the vegetable phyllosphere and their possible impact on human health remains poor. It is also not clear to us whether the composition of the protist community and its relationship with OHPs are different in the phyllosphere of organically and conventionally produced vegetables.

To reveal the associations between phyllosphere protists and OHPs in leafy greens, we characterized the protistan community in four vegetable species that applied two different production systems (organic vs conventional) sourced from four major supermarket chains in Xiamen, China. Subsequently, we quantified selected human pathogen marker genes (HPMGs) in the phyllosphere of leafy greens using high-throughput quantitative PCR (HT-qPCR) and investigated the correlations between protists and HPMGs. Finally, we isolated endosymbionts of phyllosphere protists and identified their potential pathogenicity, virulence, and resistance traits using whole genome sequencing.

## MATERIALS AND METHODS

### Sample collection

To investigate the microbial communities of vegetable phyllosphere, we collected four species of both conventionally produced leafy greens (CPLGs) and organically produced leafy greens (OPLGs), including Chinese chive (*Allium tuberosum*), Chinese cabbage (*Brassica rapa subsp. pekinensis*), cabbage (*Brassica Oleracea var. capitata*), and lettuce (*Lactuca sativa*) with three replicates of each during September and November 2020 from four supermarket chains located in Xiamen city, China (24°48'N, 118°08'E). All organically produced leafy greens are China Organic Standard GB/T 19630-2019 certified products. The China Organic Standard GB/T 19630-2019 certification guarantees the absence of growth regulating agents, chemical and synthetic products including pesticides, etc., and the presence of a strict organic management system. This standard recommends the use of biofertilizers from organic sources, which are mainly derived from animal manures or manures mixed with crop residues in China. Chemical fertilizer has been reported to affect some specific microbial taxa in the phyllosphere by altering photosynthetic activity and carbon assimilation by leaves, leading to changing available resources on the leaf, which are thought to shape the phyllosphere microbiome [36]. Before being transported by cleaned specialized transportation, organically produced vegetables should be cleaned, sorted, preserved, etc. by cleaned-up equipment to avoid contamination. The Chinese chives and cabbage were grown in Putian, Fujian, whereas the lettuce and Chinese cabbage were from Zhangjiakou, Hebei. The OPLGs were wrapped, packaged in sealed plastic bags and refrigerated (~4°C), the CPLGs were not processed or packaged and displayed at room temperature. A total of 96 samples were immediately transferred to the lab and processed as follows.

After removing the outermost leaves, approximately 60 g of fresh vegetable leaves (both spear leaves and mature leaves) from each sample were placed into individual conical flasks (250 mL) containing 150 mL 0.01 M sterile buffer solution (120 mg NaCl, 4 mg MgSO<sub>4</sub>·7H<sub>2</sub>O, 4 mg CaCl<sub>2</sub>·2H<sub>2</sub>O, 359 mg Na<sub>2</sub>HPO<sub>4</sub>·12H<sub>2</sub>O, and 130 mg KH<sub>2</sub>PO<sub>4</sub> in 1 L of

deionized water) and shaken (180 rpm) at 30 °C for 2 h. The suspensions were then sonicated for 10 min and then divided equally into two groups, one for DNA extraction and the other for protists enrichment. Thereafter, each sample was filtered through a 0.22-μm cellulose membrane to collect the microbial community.

### DNA extraction, PCR assays, and amplicon sequencing

Membranes with adhering microbes were cut into small pieces with a sterilized cutter. DNA was extracted from them using the FastDNA<sup>®</sup> Spin Kit for Soil (MP Biomedical, Santa Ana, CA), following the instructions listed by the manufacturer. To assess the concentration and quality of DNA, a NanoDrop Spectrophotometer (Nanodrop ND-1000, Thermo Scientific, Waltham, MA) and 1.0% agarose gel electrophoresis were used. DNA was stored at -20 °C until further use.

The V4 region of the eukaryotic 18S rRNA gene was amplified using the primer set TAREukFWD1F and TAREukREV3R [37]. To avoid amplification of chloroplast sequences [38], the V5-V7 region of bacterial 16S rRNA gene was selected for amplification with primers pair 799F and 1193R [39]. The protocols describing the eukaryotic 18S rRNA gene and bacterial 16S rRNA gene amplifications are detailed in Supplementary Methods S1. High-throughput sequencing was performed on an Illumina MiSeqPE300 sequencer (Majorbio, Shanghai, China).

### Sequencing processing

After sequencing, Illumina data was processed using the QIIME2 pipeline (version 2018.11, <https://qiime2.org>) [40]. Amplicon sequence variants (ASVs) were identified using DADA2 [41]. All eukaryotic ASVs were taxonomically assigned using the Protist Ribosomal Reference (PR2) database version 4.14.0 (<https://github.com/pr2database/pr2database>) [42] and blast-2.9.0 (<http://ftp.ncbi.nlm.nih.gov/blast/executables/blast+/2.9.0/>). Bacterial taxonomic identity was determined using the Silva 138.1 database (<https://www.arb-silva.de/>) [43]. After removing mitochondria, chloroplast, Fungi, Metazoa, Rhodophyta, Streptophyta, unclassified Opisthokonta, and embryophyceae sequences [44–46], and samples with less than 455 protist sequences (the minimum number of sequences in samples to reach the plateau based on the rarefaction curve Supplementary Fig. S1), we retained 209,942 valid protist sequences assigned to 640 ASVs in 72 samples. ASVs with an identity value lower than 80% were discarded. Here, we defined core bacteria as bacterial ASVs that presenting in all samples.

### Cultivation of phyllosphere protists

Phyllosphere-derived protists collected on the membranes described above were repeatedly washed with 5 ml 0.01 M sterile Page's amoeba saline (PAS) (Supplementary Fig. S2A). Protists were isolated by serial dilution with PAS in the resulting suspensions of three replicates and were fed with (OD<sub>600</sub> 1.5) heat-inactivated *E. coli* (DH5α) in 96-well plates at 28 °C in the dark, which inevitably selects against non-bacterivorous and small protistan taxa [47]. Microscopic observations were performed daily for the presence of phyllosphere protists and their morphology and locomotion by an inverted microscope with 400 times magnification (Nikon Eclipse Ti-U, Tokyo, Japan). Each isolated protist was then subcultured three times in fresh PAS medium with heat-inactivated *E. coli* every 4–7 days. Subculture might lower the presence of false endosymbionts (e.g. incompletely digested bacteria or bacteria that accidentally appear within protists) despite the inevitable possibility of preferential enrichment. Protists were then identified via 18S rRNA gene sequencing using the PSSU (5'-CTT TCG ATG GTA GTG TAT TGG ACT AC-3') and EukB (5'-TGATCC TTC TGC AGG TTC ACC TAC-3') primers [48, 49] (Supplementary Table S1). A maximum likelihood tree was constructed using the 18S rRNA gene sequences of phyllosphere protists isolated in this study, together with eight reference sequences downloaded from the NCBI GenBank using IQ-TREE (<http://iqtree.cibiv.univie.ac.at/>) with 1000 bootstrap iterations.

### Isolation of bacterial endosymbionts in phyllosphere protists

To obtain the bacterial endosymbionts that can stably survive in the protists, protists were separated from extracellular bacteria using S3e Cell Sorter (Bio-Rad, USA) equipped by 488 nm laser systems as previously described [44]. Briefly, protists were filtered through a sterile nylon 100 μm cell strainer (Solarbio, China) to avoid plugging the cell sorter before sorting. A green fluorescing probe Lysotracker Green DND-26 fluorescent signal was excited by a 488 nm laser and collected in bandpass Filter1 (525/30 nm). Protists stained with Lysotracker Green DND-26 (75 nmol/L;

Invitrogen, USA) [50] were sorted with a purity yield mode based on the higher internal complexity (side scatter, SSC) versus bigger cell size (forward scatter, FSC) as well as higher green fluorescence compared to bacteria [50] (Supplementary Fig. S2B). Sorted protists were confirmed by an inverted microscope (Nikon Eclipse Ti-U, Tokyo, Japan) immediately. To further kill extracellular bacteria without harming the protists and their bacterial endosymbionts, sorted protists were then incubated for 1 h with  $50 \mu\text{g ml}^{-1}$  each of gentamicin, ampicillin, and meropenem [51] at room temperature. After rinsing twice with PAS, antibiotic-treated protists were passed through flow cytometry again to further remove extracellular bacteria. Each sample had four replicates. Sorted protists were collected by centrifuging at  $400 \times g$  for 2 min and were lysed with 0.4% Triton X-100 to release the intracellular bacteria cells from protists in three replicates. Then,  $100 \mu\text{L}$  of lysates were plated on LB agar plates [22, 51] in triplicate and incubated for 36 h at  $37^\circ\text{C}$ . Samples not amended with 0.4% Triton X-100 served as a control to validate that there were no extracellular bacteria.

### Genome sequencing and analyses

Finally, a total of 206 single-colony bacterial endosymbionts were obtained and identified by PCR using 16S rRNA gene (27F and 1492R) primers. By blasting the full-length 16S rRNA gene against the NCBI-nr database, 50 different endosymbionts were selected. Genomic DNA of 50 bacterial endosymbionts was extracted using the Wizard<sup>®</sup> Genomic DNA Purification Kit (Promega) according to the manufacturer's instructions and quantified by TBS-380 fluorometer (Turner BioSystems Inc., Sunnyvale, CA). Sequencing libraries were then sequenced at Shanghai Majorbio Bio-pharm Technology Co., Ltd, using the Illumina HiSeq X platform that generated paired-end reads of 150 bp. Genome sequences were de novo assembled using SPAdes v.3.6.2. We successfully obtained a total of 36 high-quality genomes of bacterial endosymbionts. Open reading frames (ORFs) were predicted using Prodigal v.2.6.3 [52]. Taxonomy affiliation was determined by MetaPhlan2 [53]. Marker genes of isolated bacterial endosymbionts and 38 reference genomes (30 clinical genomes and eight environmental genomes) downloaded from the NCBI GenBank were used to build a maximum likelihood tree by IQ-TREE (<http://iqtree.cibiv.univie.ac.at/>) with 1000 bootstrap iterations. The OHPs were defined according to the A-to-Z database and previous reports (Table S2). Antibiotic resistance genes (ARGs) and metal resistance genes (MRGs) were identified by comparing the amino acid sequence of each gene with sequences in the CARD database [54] and the BacMet predicted database (version 1.1; <http://bacmet.biomedicine.gu.se>) [55] with an e-value  $\leq 10^{-10}$ , identity  $\geq 80\%$  and a minimal coverage  $\geq 70\%$  [56], respectively. Virulence factor genes (VFGs) were identified with BLASTP (e-value  $\leq 10^{-10}$  with minimal identity and coverage of 80%) homology searches against the VFDB dataset, which includes only genes associated with experimentally verified virulence factors [57]. PlasFlow was used to predict plasmid sequences from ARG-carrying contigs [58]. ICEfinder online tool was used to estimate the transferability of ARGs involving integrative and conjugative elements (ICEs) and integrative and mobilizable elements (IMEs) [59]. ARGs have the potential to be transmitted if mobile genetic components (MGEs) are present within 10 open reading frames upstream or downstream of ARGs in the same contig [60].

### HT-qPCR assays

A previously described TaqMan probe-based HT-qPCR method using the WaferGen SmartChip Real-Time PCR system platform (WaferGen Inc. USA) [61], was used to quantify human pathogen genes. A total of 68 primer pairs targeting 33 human pathogens were used in addition to a universal primer pair and probe of bacterial 16S rRNA gene to act as a reference gene (Supplementary Table S3). HT-qPCR assays were performed in triplicate using the TaqMan<sup>®</sup> Gene Expression Master Mix kit. More information about amplification conditions is available in Supplementary Methods S2. The absolute abundance (copies/ $\mu\text{L}$ ) of microbial markers was calculated according to standard curves (Supplementary Table S4).

### Statistical analysis

The  $\alpha$ -diversity of bacterial and protistan communities was estimated using: the Shannon index and the number of ASVs using QIIME2. Protistan trophic groups were assigned including phagotrophic protist, phototrophic protist, plant parasitic protist, animal parasitic protist, and human parasitic protist based on functional guilds at the genus level [62, 63]. One-way analysis of variance (ANOVA) was used to test differences in  $\alpha$ -diversity

using SPSS. Venn diagrams were created using the suite of "OmicStudio" online tools (<https://www.omicstudio.cn/tool/>) [64]. Permutational multivariate analysis of variance (PERMANOVA) was performed using the Adonis function with 999 permutations and sample ordination was visualized in a Principal coordinate analysis (PCoA). Heatmaps were performed by the "vegan" package in R 3.4.4 [65]. We evaluated the relationship between the absolute abundance of Top4 pathogen marker genes and microbial diversity (independently for bacteria, phagotrophic protists, and phototrophic protists) by linear regression. Random forest analysis was then conducted to identify predictors of the absolute abundance of prevalent pathogen marker genes using the "randomForest" R package [66]. To test the relationships between the diversity (observed richness and Shannon index) of microbe (bacteria and phagotrophic/phototrophic protists) or relative abundance of phyllosphere phagotrophic/phototrophic protists and the overall abundance or diversity of HPM, linear regression model, Pearson correlations, and associated significance were conducted with `lm()` function in R [67].


Co-occurrence networks were used to identify potential interactions between protists and bacteria for each studied leafy green. Abundant protistan and bacterial ASVs (i.e., with average relative abundance  $>0.1\%$  and occurring in  $>30\%$  of samples) were selected for the construction of co-occurrence networks. A pairwise Spearman correlation matrix was calculated using the "psych" package of R (Spearman; P. adjust method: FDR). Spearman correlation coefficients of ( $\rho$ )  $>0.5$  (or  $< -0.5$ ) with  $p < 0.05$  were selected for the microbial network analysis to include a range of interaction strengths (not only strong interactions) [68]. Network properties were characterized and visualized using Gephi 0.9.2. The relationship between microbial nodes and quantified HPMGs was assessed using robust Spearman correlation coefficients of ( $\rho$ )  $>0.7$  (or  $< -0.7$ ) and  $p < 0.01$  [69, 70]. We focused on the association between microbial nodes and HPMGs and omitted the connections within the microbes or HPMGs. The roles of nodes in the bacteria-HPMG-protist network were assigned based on the method proposed by Guimera and Amaral (2005) [71]. Roles included: within-module degree ( $Z_i$ ) which measures how well a particular feature is connected to others in the same module, and among-module connectivity or participation coefficient ( $P_i$ ), which measures how a feature is linked to other modules in the network. Nodes were classified as peripherals, hubs, and connectors iterating their roles in the whole microbial network as previously described [72].

## RESULTS

### Phyllosphere protistan and bacterial communities across vegetable species

For all vegetable species, Alveolata (35.1%), Archaeplastida (32.6%), and Stramenopiles (17.2%) were the dominant protistan supergroup while *Colpoda* (phagotrophic; 16.1%), *Spumella* (phagotrophic), and *Desmodesmus* (phototrophic) were the Top 3 dominant genus (Fig. 1A). In terms of trophic groups, phototrophic protists (44.0%) characterized by Archaeplastida, and phagotrophic protists (44.8%) had comparable relative abundance (Fig. 1A). Colpodea (Ciliophora, Alveolata) was the main phagotrophic protistan class in Chinese chives, Chinese cabbages, and lettuces, while Filosa-Sarcomonadea (Cercozoa, Rhizaria) was the dominant phagotrophic protistan class in cabbages. Chrysophyceae (Ochrophyta; Stramenopiles) was the main phototrophic protistan classes in both Chinese cabbage and Cabbage samples, while Chlorophyceae (Chlorophyta; Archaeplastida) and Trebouxiophyceae (Chlorophyta; Archaeplastida) were the dominant phototrophic protistan classes in Chinese chives and lettuces, respectively (Supplementary Fig. S2). Animal parasite Apicomplexa was the most frequent parasite presenting in 29.2% of phyllosphere samples. Only one human parasite monocyte was found in one Chinese chive sample, with a relative abundance of 2.2% (Fig. 1A). There was no significant difference in protistan diversity or community composition between organic vs. conventional production systems (Fig. 1B, D). However, protistan richness (Fig. 1B;  $p < 0.05$ ) and community composition (Fig. 1C,  $p = 0.001$ ) differed significantly across the four analyzed vegetable species. Chinese chive had the highest protist diversity (Fig. 1B,  $p < 0.01$ ), with the highest abundance of Archaeplastida





**Fig. 1** Distribution and taxonomic composition of phyllosphere protists in the leafy greens. **A** Protistan community composition of vegetable species presenting the proportion of the different protist supergroups, genera, and trophic groups. **B** Diversity index of protistan community across different vegetable species and different production systems (OPLGs vs CPLGs). **C** Principal coordinate analysis showing the distribution pattern of phyllosphere protistan communities in the four studied vegetable species and **D** different production systems. Significant differences in protistan taxa and diversity (compared to other vegetables) across vegetable species are indicated with asterisks (sign test, \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ). “OPLGs” represents organically produced leafy greens and “CPLGs” represents conventionally produced leafy greens.

(Ochrophyta, phototrophic protists; Fig. 1A,  $p < 0.05$ ). Cabbage showed the lowest protistan richness (Fig. 1B) but higher relative abundances of Stramenopiles than other vegetables. Chinese cabbage had the highest relative abundances of phagotrophic protists and Alveolata (Fig. 1A,  $p < 0.05$ ). We identified no shared protistan ASVs among vegetable species, but each vegetable had its unique protistan ASVs (Supplementary Fig. S4A).


In OPLGs and CPLGs, core bacterial ASVs represented 45.2% and 51.9% of the total phyllosphere bacteria, respectively (Supplementary Fig. S4B). Proteobacteria and Bacteroidetes were the most dominant bacterial phyla accounting for more than 75% of phyllosphere bacteria, with *Chryseobacterium* and *Pseudomonas* being the two most abundant genera (Supplementary Fig. S5C). As with protists, bacterial community composition differed between vegetable species (Supplementary Fig. S5A,  $p = 0.001$ ). Chinese chive and lettuce had the highest bacterial richness (Supplementary Fig. S5B). The relative abundance of *Allorhizobium-Neorhizobium-Pararhizobium-Rhizobium* and *Curto-bacterium* was significantly higher in Chinese chive and Chinese cabbage than in other vegetables, respectively (Supplementary Fig. S5C,  $p < 0.05$ ). Production processes had a slight effect on the

phyllosphere bacterial community composition and diversity (Supplementary Figs. S5A and S5B).

Not surprisingly, phyllosphere microbial networks also differed across vegetable species (Supplementary Fig. S6A). Chinese chive had more negative edges and the most complex network (highest number of nodes, edges, and average degree) followed by lettuces, while neither cabbage nor Chinese cabbage networks had protistan nodes (Supplementary Fig. S6B).

### Correlations between human pathogens and phyllosphere protists

Twenty-six potential human pathogens including 61 marker genes were detected in 86 of 96 vegetable phyllosphere samples (Fig. 2). *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Cronobacter* spp. were the most prevalent and abundant pathogens occurring in 58.1%, 52.3%, 31.4% and 38.4% of all samples, respectively. Absolute abundance of *K. pneumoniae* *phoE*, *S. aureus* *tufA*, *P. aeruginosa* *regA*, and *Cronobacter* spp. ITS genes were  $1.1 \times 10^3$ – $4.0 \times 10^6$  copies/g,  $1.1 \times 10^3$ – $1.5 \times 10^5$  copies/g,  $1.2 \times 10^3$ – $5.1 \times 10^4$  copies/g, and  $1.7 \times 10^3$ – $3.1 \times 10^6$  copies/g, respectively. Four pathogenic



**Fig. 2 Absolute abundance of human pathogen marker genes in the vegetable phyllosphere.** Values presented in the heatmap were log-transformed. The bar plot on the left indicates the total absolute abundance and detection rate of each human pathogen marker gene (HPMG). The bar plot on the top shows the total abundance of HPMGs from each vegetable species, respectively. The red text represents pathogenic protists. Marker genes of *Klebsiella pneumoniae* and *Cronobacter* spp. showed the highest abundance in the phyllosphere of Chinese chives (sign test,  $**p < 0.01$ ). “OPLGs” represents organically produced leafy greens and “CPLGs” represents conventionally produced leafy greens.

protists: *Acanthamoeba* spp., *Cryptosporidium*, *Giardia lamblia*, and *Entamoeba histolytica* accounting for 1.8 % of the overall abundance of HPMGs, were only positive in less than ten samples (Fig. 2), of which *Acanthamoeba* spp. and *Giardia lamblia* had the relatively high abundance of  $1.6 \times 10^2$ – $3.0 \times 10^4$  copies/g and  $3.2 \times 10^2$ – $4.9 \times 10^4$  copies/g, respectively. There were also significant differences in HPMGs detected in different vegetable species (Supplementary Fig. S7A,  $P = 0.001$ ) but not between OPLGs and CPLGs (Supplementary Fig. S7B). The phyllosphere of Chinese chives had the greatest total abundance of HPMGs (ranging from  $2.7 \times 10^3$  to  $4.0 \times 10^6$  copies/g), followed by lettuce, cabbage, and Chinese cabbage (Fig. 2). *K. pneumoniae* and *Cronobacter* spp. ITS were significantly enriched in the phyllosphere of Chinese chives (Fig. 2,  $p < 0.01$ ).

The overall abundance of HPMGs was significantly and positively linked to the relative abundance (Supplementary Fig. S8A;  $p < 0.001$ ) and diversity (Supplementary Fig. S8C;  $p < 0.05$ ) of phagotrophic protists rather than bacteria (Supplementary Fig. S9). While the relative abundance of phototrophic protists negatively correlated with the overall abundance (Supplementary Fig. S8G;  $p < 0.05$ ) and diversity of HPMGs (Supplementary Fig. S8H;  $p < 0.05$ ). Specifically, the absolute abundance of Top4 prevalent HPMGs (*K. pneumoniae* *phoE*, *S. aureus* *tufA*, *P. aeruginosa* *regA*, and *Cronobacter* spp. ITS genes) positively correlated to the diversity and relative abundance of phagotrophic protists, but negatively correlated to the relative


abundance of phototrophic protists (Fig. 3A, B) rather than bacteria (Supplementary Fig. S9). In particular, the diversity of phagotrophic protists positively correlated with the absolute abundance of *K. pneumoniae* *phoE* (Fig. 3A;  $p < 0.001$ ). The absolute abundance of *P. aeruginosa* *regA* was significantly and positively associated with the relative abundance of phagotrophic protists (Fig. 3B;  $p < 0.05$ ). Random forest analyses statistically supported that diversity and the relative abundance of phagotrophic protists best predicted the absolute abundance of *K. pneumoniae* *phoE* ( $p < 0.01$ ), explaining more than half of the variation in the absolute abundance of *K. pneumoniae* *phoE*. Co-occurrence analysis identified 12 ASVs that were strongly and positively associated with HPMGs (Spearman correlation  $\geq 0.7$ ,  $p < 0.01$ ; Fig. 3D). The phagotrophic protist *Colpoda* (PASV982), which acted as a connector in the network (Supplementary Fig. S10) was closely related to the HPMGs of *Salmonella*, *L. pneumophila* and *V. cholerae* (Fig. 3D). Bacterial ASV708 assigned to *Chryseobacterium* was most tightly linked to pathogens, with up to 13 pathogens including 18 marker genes (Fig. 3D).

#### Isolated bacterial endosymbionts of phyllosphere protists

To gain deeper insight into the positive relations between phyllosphere protists and OHPs, we isolated eleven phyllosphere protists and their bacterial endosymbionts. All phyllosphere protist isolates (Supplementary Table S1) were phagotrophic protists, annotated as *Colpoda steini* PAPS-4, *Aspidisca lynceus*, *Spumella*







**Fig. 4** Maximum-likelihood phylogenetic trees of isolated phyllosphere protists and their bacterial endosymbionts. Sequences obtained in this study were marked in bold. Orange bars on the right and the top display the total number of endosymbionts bacteria isolated from phyllosphere protists, and the detection frequency of each endosymbiont bacteria, respectively.

$p < 0.001$ ), and adherence VFs (Type IV pili;  $p < 0.01$ ) than clinical genomes, respectively (Fig. 5B, Supplementary Figs. S15 and S16).

Furthermore, we analyzed the transferability potential for ARGs, MRGs, and VFGs carried by endosymbiont OHPs. Here, co-localizations of ARGs, MRGs, and MGEs were observed in both plasmids and chromosomal contigs. For example, on a plasmid contig of *O. anthropi* MN31484444, ARG *mdsB* was located near two insertion sequences (IS) *IS3* and *IS5* (<10 kb, Fig. 5C), meanwhile, MRGs *copC* and *cutO* were located near IS fragments *IS21* and *IS630*, respectively. MRGs *arsH* was observed near to *IS3* on a chromosomal contig of *P. nitroreducens* MN31484453. Besides, *ceoB* and *oqxB* genes encoding for multidrug antibiotic efflux pumps were identified on conjugative plasmid contigs of *A. xylooxidans* (Fig. 5C).


## DISCUSSION

The vegetable phyllosphere is an important habitat for protists [73, 74]. Here, although phyllosphere protists were variable between replicate samples, *Colpoda* were the most dominant and prevalent phyllosphere protists. This result is consistent with previous isolation-based evidence of *Colpoda* spp. dominance on the leaves [75] and amplicon-based evidence of *Colpoda* dominance on solanaceous plants [76]. Interestingly, we observed host-specific protists and no shared protistan ASV across vegetable species, suggesting that host selection of vegetable species largely shaped the distribution of protistan communities. This is potentially associated with the sensitivity to environmental changes of protists [77], as protists are likely to colonize selectively according to leaf surface factors, such as blade roughness, stomatal density, hydrophobicity, and level of epicuticular wax [78, 79]. In this study, cabbage harbored the lowest protistan diversity possibly attributed to a high level of hydrophobic waxy cuticle that reduces water on the surface and is thus unfavorable for protist colonization [80]. The species-specific protistan could also drive through bottom-up forces of bacterial communities, which are major prey for phagotrophic protists and are

significantly influenced by host selections [81]. Besides, in this study, production processes hardly influenced the alpha- or beta-diversity of phyllosphere protistan communities, which differed from soil protistans communities exhibiting distinct patterns between organic and chemically fertilized soils [82]. One explanation for this could be the multiple factors in the open nature of the phyllosphere [83], including unknown plant cultivation, transportation, and produce-packing conditions [84], which may weaken the influence of different production processes on phyllosphere protistan variations [85].

Host selection [86] and consequent different protistan communities and different microbial co-occurrence patterns may also affect the colonization of human pathogens in the phyllosphere, leading to different risks of human pathogens in vegetables. In this study, Chinese chives and lettuces might constitute a higher epidemiological risk than cabbage due to their higher content of OHPs. Specifically, *S. aureus*, *K. pneumoniae*, *P. aeruginosa*, and *Cronobacter* spp., were the prevalent OHPs found in all studied vegetable species. These results suggest that the ubiquitous detection of human pathogens in leafy greens is a vital health concern and that we should wash leafy greens repeatedly before eating them raw whether organically or conventionally grown [87]. The significant positive association between phyllosphere phagotrophic protists (both the diversity and the relative abundance) and the abundance of the Top 4 HPMGs, and the highest explanation of the phagotrophic protists on the abundance of *K. pneumoniae* *phoE* and *P. aeruginosa* *regA* both indicated a strong connection between phyllosphere phagotrophic protists and OHPs. This strong connection can be partially explained by the selective predation of phagotrophic protists on various human pathogens, including *K. pneumoniae*, *Pseudomonas aeruginosa*, and *Staphylococcus aureus* [15]. Phagotrophic protists can also influence the colonization of OHPs via regulating nutrient exchange [88] or pathogen-suppressing secondary metabolite genes [89].

Another way to link phagotrophic protists and OHPs is to become endosymbiosis. A key finding when isolating



**Fig. 5 Key features of endosymbiont genomes isolated from phyllosphere protists. A** Maximum-likelihood phylogenetic trees of endosymbiont genomes and reference genomes with 1000 trials. Sequences obtained in this study were marked in bold. Potential human opportunistic pathogens are marked with a red triangle. [E] and [C] represent the reference genomes from the environment and the clinic, respectively. The size of the purple circles indicates the genome size of each endosymbiont isolate. Heatmap denotes ARG types carried by the corresponding genomes, the purple, orange, and green bars display the total number of ARGs and VFGs carried by the corresponding isolates, respectively. **B** Comparative analysis of ARGs and VFGs carried by clinical genomes and three endosymbiont genomes (*A. xylooxidans*, *O. anthropi* and *S. maltophilia*). **C** Schematic of the genetic organization of ARGs, MRGs, and MGEs. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ .



phyllosphere protists and their endosymbionts was that phagotrophic protists *Colpoda steini*, HPMGs-associated protists in the network, internalize multiple antibiotic-resistant OHPs, including *P. nitroreducens*, *A. xylosoxidans* and *S. maltophilia*. *A. xylosoxidans* and *S. maltophilia* possessing the second highest number of ARGs and VFGs in this study, are common amoeba-resisting endosymbionts [90–92] associated with cystic fibrosis and respiratory infections [93, 94]. *P. nitroreducens*, never been reported as an endosymbiont of protists before, carried the highest number of ARGs and VFGs and has a close phylogenetic relationship with common protists endosymbiont *P. aeruginosa* [34, 95]. Hence, similar to *P. nitroreducens*, *P. aeruginosa* may establish a positive correlation with the relative abundance of phyllosphere phagotrophic protists through endosymbiosis, though we did not isolate it. In addition, four OHPs species only found within protists indicated that protists as a potential reservoir for OHP may transfer endosymbiont OHPs from other environments, resulting in the potential increase of OHPs in the phyllosphere. Moreover, all isolated protist-adapted endosymbiont OHPs have the potential to return to the environment even with more environmental persistence [96]. This may partly explain why some endosymbiont OHPs occurred in both free-living and isolated communities. Nonetheless, the findings of this study did not reveal the percentage of OHPs released from protists to the environment and the relative contribution of protistan predation on the variation of OHPs in the phyllosphere, which calls for more future research to be addressed. Given the limited culturable phyllosphere protists and the absence of obligate intracellular bacteria that cannot be cultivated without their hosts in this study, there may be more pathogenic endosymbionts OHPs within protists, indicating that more attention should be directed to the role of phagotrophic protists as important vectors of the transmission of OHPs in the vegetable phyllosphere.

Two main survival strategies of intracellular bacteria have been reported for digestion resistance [97]: escaping into the cytosol [98]; subverting antimicrobial mechanisms including removing toxic compounds, preventing phagosome-lysosome fusion, modulating phagosomal pH, etc [19]. In this study, the specific resistance and virulence traits of OHPs within protists may act as subvert antimicrobial mechanisms partially contributing to resisting digestion and facilitating the survival of the endosymbionts [97]. For instance, efflux pumps SmeDEF (ARGs) carried by OHPs *S. maltophilia* have been demonstrated to be induced by secondary metabolites of amoeba to directly expel toxic compounds secreted by *Acanthamoeba castellanii* [11], allowing the survival and intracellular growth of *S. maltophilia*, and contribute to intrinsic multidrug resistance of *S. maltophilia* [99]. The VF T6SS such as VgrG-1 found in *P. nitroreducens* were reported to defend against protist digestion by injecting antimicrobial toxins [100–102], or causing cytotoxic actin-crosslinking in the predatory amoeba *Dictyostelium discoideum* [103]. As for MRGs, arsenic (such as arsenic resistance operon *arsH*) and copper resistance genes (such as *copA*) carried by *A. xylosoxidans* may help adapt to copper and arsenic poisoning by *Dictyostelium* grazing and thus escape phagocytosis [17].

In our assay, most isolated endosymbionts OHPs carried both VFGs, MRGs, and ARGs, and the number of VFGs or MRGs and ARGs was positively correlated, indicating the co-occurrence profiles among them. The intimate co-occurrence and the high cotransfer potential among ARGs, VFGs, and MRGs have more often observed in genomes from pathogen species [104], such as *K. pneumoniae* [105] and enterococci [106], which potentially complicate the treatment of OHPs infections. Here, the residency of VFGs, MRGs, and ARGs on MGEs (plasmid fragments) from OHPs further indicated the transmission potential of virulence and resistance properties using mobile genetic elements via HGT. Moreover, HGT has been reported to increase greatly within phagotrophic protists contributing to the adaptation and

evolution of endosymbionts OHPs [107]. For example, the conjugation frequency of *E. coli* strains engulfed by the ciliate increased 2000- to 4000-fold after a full digestion cycle [108]. Similarly, protozoal predation enhances the transformation of a gene cassette in *Vibrio cholerae* by as much as 405-fold through SOS-regulated DNA integration [109]. These indicated that the HGT of ARGs, VFGs, and MRGs among endosymbiont OHPs may be further expedited under the predation pressure of protists [110], and thus contribute to the survival of more endosymbionts [111] and the emergence of resistant OHPs. In addition, endosymbionts *A. xylosoxidans* and *S. maltophilia* showed higher virulence potential including invasive adenylate cyclase/haemolysin, Resistance/nodulation/division (RND)-type efflux pump, attachment to host cells and biofilm formation, but lower antibiotic resistance potential than clinical genomes, which probably relate to HGT and less frequent exposure to antibiotics [105]. Overall, phyllosphere protists can be important vectors for the transmission of resistant OHPs and the resistance and virulence traits of OHPs may even be exchanged among endosymbionts OHPs which needs more attention.

## CONCLUSIONS

We revealed that phagotrophic protists in the phyllosphere of leafy greens can harbor multiple resistant OHPs and are closely related to OHP abundance. In these leafy greens, *Colpoda* was identified as the key phagotrophic protist of OHPs. ARGs, VFGs, and MRGs co-occur at high frequencies in endosymbionts OHPs with the potential for HGT, which contributes to the emergence of resistant OHPs in the phyllosphere of leafy greens and poses risks to consumers, and within the one health perspective, to the environment. Besides, host selection of vegetable species largely affects the distribution of OHPs and protistan communities. This work fills the knowledge gap regarding the association between phyllosphere protists and HPMGs and emphasizes that phyllosphere phagotrophic protists in leafy greens are more important in the transmission and evolution of OHPs than hitherto assumed. These findings advance our understanding of phyllosphere protists and are valuable for environmental health and food safety under the One Health framework.

## DATA AVAILABILITY

All raw sequencing data were deposited to the Sequence Read Archive (SRA) under the number PRJNA855267.

## REFERENCES

1. Ferreira G, Turk Z. An overview of organic vegetable production in the United States. USDA Economic Research Service. 2016;VGS-357-SA2. [https://www.ers.usda.gov/webdocs/outlooks/74639/60319\\_vgs-357-sa2\\_organicveg.pdf?v=1644.8](https://www.ers.usda.gov/webdocs/outlooks/74639/60319_vgs-357-sa2_organicveg.pdf?v=1644.8).
2. Zhou SYD, Zhu D, Giles M, Yang XR, Daniell T, Neilson R, et al. Phyllosphere of staple crops under pig manure fertilization, a reservoir of antibiotic resistance genes. *Environ Pollut*. 2019;252:227–35.
3. Lenzi A, Marvasi M, Baldi A. Agronomic practices to limit pre- and post-harvest contamination and proliferation of human pathogenic Enterobacteriaceae in vegetable produce. *Food Control*. 2021;119:107486.
4. Neil KP, Gwen B, Kathryn MCOJ, Eija T, Carlota M, Musser KA, et al. A novel vehicle for transmission of *Escherichia coli* O157:H7 to humans: multistate outbreak of *E. coli* O157:H7 infections associated with consumption of ready-to-bake commercial prepackaged cookie dough-United States, 2009. *Clin Infect Dis*. 2012;54:511–8.
5. Liu Q, Jin X, Feng X, Yang H, Fu C. Inactivation kinetics of *Escherichia coli* O157:H7 and *Salmonella Typhimurium* on organic carrot (*Daucus carota* L.) treated with low concentration electrolyzed water combined with short-time heat treatment. *Food Control*. 2019;106:106702.
6. Zhu B, Chen Q, Chen S, Zhu YG. Does organically produced lettuce harbor higher abundance of antibiotic resistance genes than conventionally produced? *Environ Int*. 2017;98:152–9.

7. Alsanius BW, von Essen E, Hartmann R, Vågsholm I, Doyle O, Schmutz U, et al. The “one health”-concept and organic production of vegetables and fruits. *Acta Hortic.* 2019;1242:1–14.
8. Castro-Ibáñez I, Gil MI, Allende A. Ready-to-eat vegetables: current problems and potential solutions to reduce microbial risk in the production chain. *LWT Food Sci Technol.* 2016;85:284–92.
9. Berger CN, Sodha SV, Shaw RK, Griffin PM, Pink D, Hand P, et al. Fresh fruit and vegetables as vehicles for the transmission of human pathogens. *Environ Microbiol.* 2010;12:2385–97.
10. Mogren L, Windstam S, Boqvist S, Vågsholm I, Söderqvist K, Rosberg AK, et al. The hurdle approach—A holistic concept for controlling food safety risks associated with pathogenic bacterial contamination of leafy green vegetables. A review. *Front Microbiol.* 2018;9:1965.
11. Denet E, Triadou S, Michalet S, Nazaret S, Favre-Bonté S. Growth of *Stenotrophomonas maltophilia* and expression of Sme efflux pumps encoding genes in the presence of supernatants from amoebal and bacterial co-cultures: towards the role of amoebal secondary metabolites. *Environ Microbiol Rep.* 2020;12:702–11.
12. Bashir I, War AF, Rafiq I, Reshi ZA, Rashid I, Shouche YS. Phyllosphere microbiome: diversity and functions. *Microbiol Res.* 2022;254:126888.
13. Michal S, Radka D, Vladimír B, Petr K, Monika M, Iva S. *Toxoplasma gondii* in vegetables from fields and farm storage facilities in the Czech Republic. *FEMS Microbiol Lett.* 2019;366:fnz170.
14. Maha B, Erica K, Hunter PR, Sitara SA. Risk factors for *Cryptosporidium* infection in low and middle income countries: a systematic review and meta-analysis. *PLoS Negl Trop Dis.* 2018;12:e0006553.
15. Bornier F, Zas E, Potheret D, Laaberki MH, Charpentier X. Environmental free-living amoebae can predate on diverse antibiotic-resistant human pathogens. *Appl Environ Microbiol.* 2021;87:e0074721.
16. Bonkowski M. Protozoa and plant growth: the microbial loop in soil revisited. *New Phytol.* 2004;162:617–31.
17. Hao X, Li X, Pal C, Hobman J, Larsson DG, Saquib Q, et al. Bacterial resistance to arsenic protects against protist killing. *Biometals.* 2017;30:307–11.
18. Hiltunen T, Becks L. Consumer co-evolution as an important component of the eco-evolutionary feedback. *Nat Commun.* 2014;5:5226.
19. Casadevall A. Evolution of intracellular pathogens. *Annu Rev Microbiol.* 2008;62:19–33.
20. Erken M, Lutz C, McDougald D. The rise of pathogens: Predation as a factor driving the evolution of human pathogens in the environment. *Microb Ecol.* 2013;65:860–8.
21. Casas V, Maloy S. Role of bacteriophage-encoded exotoxins in the evolution of bacterial pathogens. *Fut Microbiol.* 2011;6:1461–73.
22. Hao X, Luthje F, Ronn R, German NA, Li X, Huang F, et al. A role for copper in protozoan grazing – two billion years selecting for bacterial copper resistance. *Mol Microbiol.* 2016;102:628–41.
23. Van der Henst C, Scignari T, Maclachlan C, Blokesch M. An intracellular replication niche for *Vibrio cholerae* in the amoeba *Acanthamoeba castellanii*. *ISME J.* 2016;10:897–910.
24. Vieira A, Ramesh A, Seddon AM, Karlyshev AV. CmeABC multidrug efflux pump contributes to antibiotic resistance and promotes *Campylobacter jejuni* survival and multiplication in *Acanthamoeba polyphaga*. *Appl Environ Microbiol.* 2017;83:e01600–17.
25. Hamann E, Gruber-Vodicka H, Kleiner M, Tegetmeyer HE, Riedel D, Littmann S, et al. Environmental breviatea harbour mutualistic *Arcobacter epibionts*. *Nature.* 2016;534:254–8.
26. Hoque MM, Noorian P, Espinoza-Vergara G, Manuneedhi Cholan P, Kim M, Rahman MH, et al. Adaptation to an amoeba host drives selection of virulence-associated traits in *Vibrio cholerae*. *ISME J.* 2022;16:856–67.
27. Personnic N, Striednig B, Hilbi H. Quorum sensing controls persistence, resuscitation, and virulence of *Legionella* subpopulations in biofilms. *ISME J.* 2021;15:196–210.
28. Hilbi H, Weber SS, Ragaz C, Nyfeler Y, Urwyler S. Environmental predators as models for bacterial pathogenesis. *Environ Microbiol.* 2007;9:563–75.
29. Folkins MA, Dey R, Ashbolt NJ. Interactions between human reovirus and free-living amoebae: implications for enteric virus disinfection and aquatic persistence. *Environ Sci Technol.* 2020;54:10201–6.
30. Brandl MT, Rosenthal BM, Haxo AF, Berk SG. Enhanced survival of *salmonella enterica* in vesicles released by a soilborne *tetrahymena* species. *Appl Environ Microbiol.* 2005;71:1562–9.
31. Anacarso I, de Niederhausern S, Messi P, Guerrieri E, Iseppi R, Sabia C, et al. *Acanthamoeba polyphaga*, a potential environmental vector for the transmission of food-borne and opportunistic pathogens. *J Basic Microbiol.* 2012;52:261–8.
32. Bekele F, Tamirat T, Gellila B, Tsegaye Y. Parasitic contamination of raw vegetables and fruits collected from selected local markets in Arba Minch town, Southern Ethiopia. *Infect Dis Poverty.* 2017;6:19.
33. Rodrigues AC, da Silva MDC, Pereira RAS, Pinto LC. Prevalence of contamination by intestinal parasites in vegetables (*Lactuca sativa* L. and *Coriandrum sativum* L.) sold in markets in Belem, northern Brazil. *J Sci Food Agric.* 2020;100:2859–65.
34. Leong W, Poh WH, Williams J, Lutz C, Hoque MM, Poh YH, et al. Adaptation to an amoeba host leads to *Pseudomonas aeruginosa* isolates with attenuated virulence. *Appl Environ Microbiol.* 2022;88:e02322–21.
35. König L, Wentrup C, Schulz F, Wascher F, Escola S, Swanson MS, et al. Symbiont-mediated defense against *Legionella pneumophila* in Amoebae. *mBio.* 2019;10:e00333.
36. Darlison J, Mogren L, Rosberg AK, Gruden M, Minet A, Line C, et al. Leaf mineral content govern microbial community structure in the phyllosphere of spinach (*Spinacia oleracea*) and rocket (*Diplotaxis tenuifolia*). *Sci Total Environ.* 2019;675:501–12.
37. Stoeck T, Bass D, Nebel M, Christen R, Richards TA. Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol Ecol.* 2010;19:21–31.
38. Beckers B, Beeck MOD, Weyens N, Boerjan W, Vangronsveld J. Structural variability and niche differentiation in the rhizosphere and endosphere bacterial microbiome of field-grown poplar trees. *Microbiome.* 2017;5:25.
39. Bodenhausen N, Horton MW, Joy B. Bacterial communities associated with the leaves and the roots of *Arabidopsis thaliana*. *PLoS One.* 2013;8:e56329.
40. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol.* 2019;37:852–7.
41. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson A, Holmes SP. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods.* 2015;13:581–3.
42. Guillou L, Bachar D, Audic S, Bass D, Berney C, Bittner L, et al. The Protist Ribosomal Reference database (PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated taxonomy. *Nucleic Acids Res.* 2013;41:597–604.
43. Quast C, Pruesse E, Yilmaz P, Gerken J, Glckner FO. The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* 2012;41:590–6.
44. Li LJ, Lin CS, Huang XR, An XL, Li WJ, Su JQ, et al. Characterizing potential pathogens from intracellular bacterial community of protists in wastewater treatment plants. *Environ Int.* 2023;171:107723.
45. Singer D, Seppey CVW, Lentendu G, Dunthorn M, Bass D, Belbahri L, et al. Protist taxonomic and functional diversity in soil, freshwater and marine ecosystems. *Environ Int.* 2021;146:106262.
46. Xiong W, Jousset A, Li R, Delgado-Baquerizo M, Bahram M, Logares R, et al. A global overview of the trophic structure within microbiomes across ecosystems. *Environ Int.* 2021;151:106438.
47. Geisen S. The bacterial-fungal energy channel concept challenged by enormous functional versatility of soil protists. *Soil Biol Biochem.* 2016;102:22–5.
48. Miyaoka Y, Hatamoto M, Yamaguchi T, Syutsubo K. Eukaryotic community shift in response to organic loading rate of an aerobic trickling filter (down-flow hanging sponge reactor) treating domestic sewage. *Microb Ecol.* 2017;73:801–14.
49. Karnati SKR, Yu Z, Sylvester JT, Dehority BA, Morrison M, Firkins JL. Technical note: specific PCR amplification of protozoal 18S rDNA sequences from DNA extracted from ruminal samples of cows. *J Anim Sci.* 2003;81:812–5.
50. Rose JM, Caron DA, Sieracki ME, Poulton P. Counting heterotrophic nanoplanktonic protists in cultures and aquatic communities by flow cytometry. *Aquat Microb Ecol.* 2004;34:263–77.
51. Ronn R, Hao X, Luthje F, German N, Li X, Huang F, et al. Bacterial survival in *Dictyostelium*. *Bio Protoc.* 2017;7:e2376.
52. Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinform.* 2010;11:119.
53. Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, et al. MetaPhlan2 for enhanced metagenomic taxonomic profiling. *Nat Methods.* 2015;12:902–3.
54. Chen Q, An X, Li H, Su J, Ma Y, Zhu YG. Long-term field application of sewage sludge increases the abundance of antibiotic resistance genes in soil. *Environ Int.* 2016;92:93:1–10.
55. Chandan P, Johan BP, Christopher R, Erik K, Joakim L. BacMet: antibacterial biocide and metal resistance genes database. *Nucleic Acids Res.* 2014;42:D737–D43.
56. Zhao R, Yu K, Zhang J, Zhang G, Huang J, Ma L, et al. Deciphering the mobility and bacterial hosts of antibiotic resistance genes under antibiotic selection pressure by metagenomic assembly and binning approaches. *Water Res.* 2020;186:116318.
57. Liu B, Zheng D, Jin Q, Chen L, Yang J. VFDB 2019: a comparative pathogenomic platform with an interactive web interface. *Nucleic Acids Res.* 2019;47:D687–D92.

58. Krawczyk PS, Lipinski L, Dziembowski A. PlasFlow: predicting plasmid sequences in metagenomic data using genome signatures. *Nucleic Acids Res.* 2018;46:e35–e.
59. Liu M, Li X, Xie YZ, Bi DX, Sun JY, Li J, et al. ICEberg 2.0: an updated database of bacterial integrative and conjugative elements. *Nucleic Acids Res.* 2018;47:D660–D5.
60. Saenz JS, Marques TV, Barone RSC, Cyrino JEP, Kublik S, Nesme J, et al. Oral administration of antibiotics increased the potential mobility of bacterial resistance genes in the gut of the fish *Piaractus mesopotamicus*. *Microbiome.* 2019;7:24.
61. An XL, Wang JY, Pu Q, Li H, Pan T, Li HQ, et al. High-throughput diagnosis of human pathogens and fecal contamination in marine recreational water. *Environ Res.* 2020;190:109982.
62. Adl SM, Bass D, Lane CE, Lukes J, Schoch CL, Smirnov A, et al. Revisions to the classification, nomenclature, and diversity of eukaryotes. *J Eukaryot Microbiol.* 2019;66:4–119.
63. Freudenthal J, Ju F, Burgmann H, Dumack K. Microeukaryotic gut parasites in wastewater treatment plants: diversity, activity, and removal. *Microbiome.* 2022;10:27.
64. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker discovery and explanation. *Genome Biol.* 2011;12:R60.
65. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, Minchin PR, et al. R package 'vegan' version 2.5-1. *Community Ecol Package. Vers.* 2018;2:1–295.
66. Breiman L. Random forests. *Mach Learn.* 2001;45:5–32.
67. Team RDC. R: a language and environment for statistical computing. Vienna, Austria: R foundation for statistical computing; 2019.
68. de Vries FT, Griffiths RI, Bailey M, Craig H, Girlanda M, Gweon HS, et al. Soil bacterial networks are less stable under drought than fungal networks. *Nat Commun.* 2018;9:3033.
69. Xiong C, Singh BK, He JZ, Han YL, Li PP, Wan LH, et al. Plant developmental stage drives the differentiation in ecological role of the maize microbiome. *Microbiome.* 2021;9:171.
70. Wu S, Dong Y, Deng Y, Cui L, Zhuang X. Protistan consumers and phototrophs are more sensitive than bacteria and fungi to pyrene exposure in soil. *Sci Total Environ.* 2022;822:153539.
71. Guimera R, Nunes Amaral LA. Functional cartography of complex metabolic networks. *Nature.* 2005;433:895–500.
72. Shi Y, Delgado-Baquerizo M, Li Y, Yang Y, Zhu YG, Penuelas J, et al. Abundance of kinless hubs within soil microbial networks are associated with high functional potential in agricultural ecosystems. *Environ Int.* 2020;142:105869.
73. Flues S, Blokker M, Dumack K, Bonkowski M. Diversity of cercozoan species in the phyllosphere and rhizosphere of different plant species with a description of *Neocercomonas epiphylla* (Cercozoa, Rhizaria) a leaf-associated protist. *J Eukaryot Microbiol.* 2018;65:587–99.
74. Leveau JH. A brief from the leaf: latest research to inform our understanding of the phyllosphere microbiome. *Curr Opin Microbiol.* 2019;49:41–9.
75. Bamforth SS. Population dynamics of soil and vegetation protozoa. *Am Zool.* 1973;13:171–6.
76. Taerum SJ, Steven B, Gage D, Triplett LR. Dominance of ciliophora and chlorophyta among phyllosphere protists of solanaceous plants. *Phytobiomes J.* 2022;PBIOMES-04-22-0021F1.
77. Guo S, Xiong W, Hang X, Gao Z, Jiao Z, Liu H, et al. Protists as main indicators and determinants of plant performance. *Microbiome.* 2021;9:64.
78. Hunter PJ, Shaw RK, Berger CN, Frankel D PH. Older leaves of lettuce (*Lactuca* spp.) support higher levels of *Salmonella enterica* ser. *Senftenberg* attachment and show greater variation between plant accessions than do younger leaves. *FEMS Microbiol Lett.* 2015;362:fnv077.
79. Macarasin D, Patel J, Bauchaun G, Giron JA, Ravishankar S. Effect of spinach cultivar and bacterial adherence factors on survival of *Escherichia coli* O157:H7 on spinach leaves. *J Food Prot.* 2013;76:1829–37.
80. Luna-Guevara JJ, Arenas-Hernandez MMP, Martinez de la Pena C, Silva JL, Luna-Guevara ML. The role of pathogenic *E. coli* in fresh vegetables: Behavior, contamination factors, and preventive measures. *Int J Microbiol.* 2019;38:1–10.
81. Xiong C, Zhu YG, Wang JT, Singh B, Han LL, Shen JP, et al. Host selection shapes crop microbiome assembly and network complexity. *New Phytol.* 2021;229:1091–104.
82. Guo S, Xiong W, Xu H, Hang X, Liu H, Xun W, et al. Continuous application of different fertilizers induces distinct bulk and rhizosphere soil protist communities. *Eur J Soil Biol.* 2018;88:8–14.
83. Remus-Emsermann MNP, Schlechter RO. Phyllosphere microbiology: at the interface between microbial individuals and the plant host. *New Phytol.* 2018;218:1327–33.
84. Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. Plant-microbiome interactions: from community assembly to plant health. *Nat Rev Microbiol.* 2020;18:607–21.
85. Zhu Y-G, Xiong C, Wei Z, Chen Q-L, Ma B, Zhou S-Y-D, et al. Impacts of global change on the phyllosphere microbiome. *New Phytol.* 2022;234:1977–86.
86. Yang JW, Wu W, Chung CC, Chiang KP, Gong GC, Hsieh CH. Predator and prey biodiversity relationship and its consequences on marine ecosystem functioning-interplay between nanoflagellates and bacterioplankton. *ISME J.* 2018;12:1532–42.
87. Yemi GP, Delaquis P. Natural compounds with antibacterial activity against *Cronobacter* spp. in powdered infant formula: a review. *Front Nutr.* 2020;7:595964.
88. Geisen S, Hu S, Dela Cruz TEE, Veen GFC. Protists as catalyzers of microbial litter breakdown and carbon cycling at different temperature regimes. *ISME J.* 2021;15:618–21.
89. Sapp M, Ploch S, Fiore-Donno AM, Bonkowski M, Rose LE. Protists are an integral part of the *Arabidopsis thaliana* microbiome. *Environ Microbiol.* 2018;20:30–43.
90. Paes J, Kepler R, Goncalves RF, Berte FK, Virginio VG, Benitez LB, et al. Amoebal coculture and enrichment methods as a proposal for water quality control in Brazil. *Acta Trop.* 2021;223:106074.
91. Greub G, Scola BL, Raoult D, Massiliae IR, Massiliae CR. Amoebae-resisting bacteria isolated from human nasal dwabs by amoebal coculture. *Emerging Infect Dis.* 2004;10:470–7.
92. Vincent T, Gerald MD, Denyer SP, Jean-Yves M. Free-living amoebae and their intracellular pathogenic microorganisms: risks for water quality. *FEMS Microbiol Lett.* 2009;34:231–59.
93. Ridderberg W, Wang M, Norkov-Lauritsen N. Multilocus sequence analysis of isolates of *Achromobacter* from patients with cystic fibrosis reveals infecting species other than *Achromobacter xylosoxidans*. *J Clin Microbiol.* 2012;50:2688–94.
94. Gomes TS, Vaccaro L, Magnet A, Izquierdo F, Ollero D, Martinez-Fernandez C, et al. Presence and interaction of free-living amoebae and amoeba-resisting bacteria in water from drinking water treatment plants. *Sci Total Environ.* 2020;719:137080.
95. Timothy LT, Richard W, Egon O, Alan H. First report of *Pseudomonas nitroreducens* cultured from the lungs of a patient with pneumonia. *BMJ Case Rep.* 2021;14:e241327.
96. Amaro F, Martin-Gonzalez A. Microbial warfare in the wild—the impact of protists on the evolution and virulence of bacterial pathogens. *Int Microbiol.* 2021;24:559–71.
97. Strassmann JE, Shu L. Ancient bacteria-amoeba relationships and pathogenic animal bacteria. *PLoS Biol.* 2017;15:e2002460.
98. Gerstenmaier L, Pilla R, Herrmann L, Herrmann H, Prado M, Villafano GJ, et al. The autophagic machinery ensures nonlytic transmission of mycobacteria. *Proc Natl Acad Sci USA.* 2015;112:E687–E92.
99. Cho HH, Sung JY, Kwon KC, Koo SH. Expression of Sme efflux pumps and multilocus sequence typing in clinical isolates of *Stenotrophomonas maltophilia*. *Ann Lab Med.* 2012;32:38–43.
100. Flues S, Bass D, Bonkowski M. Grazing of leaf-associated Cercozoans (Protists: Rhizaria: Cercozoa) structures bacterial community composition and function. *Environ Microbiol.* 2017;19:3297–309.
101. George EE, Husnik F, Tashyreva D, Prokopchuk G, Horak A, Kwong WK, et al. Highly reduced genomes of protist endosymbionts show evolutionary convergence. *Curr Biol.* 2020;30:925–33.
102. French CT, Bulterys PL, Woodward CL, Tatters AO, Ng KR, Miller JF. Virulence from the rhizosphere: ecology and evolution of *Burkholderia pseudomallei*-complex species. *Curr Opin Microbiol.* 2020;54:18–32.
103. Joshi A, Kostiuik B, Rogers A, Teschler J, Pukatzki S, Yildiz FH. Rules of engagement: The type VI secretion system in *Vibrio cholerae*. *Trends Microbiol.* 2017;25:267–79.
104. Li LG, Xia Y, Zhang T. Co-occurrence of antibiotic and metal resistance genes revealed in complete genome collection. *ISME J.* 2017;11:651–62.
105. Derakhshan S, Najjar Peerayeh S, Bakhshi B. Association between presence of virulence genes and antibiotic resistance in clinical *Klebsiella Pneumoniae* isolates. *Lab Med.* 2016;47:306–11.
106. Biswas K, Sharma P, Joshi SR. Co-occurrence of antimicrobial resistance and virulence determinants in enterococci isolated from traditionally fermented fish products. *J Glob Antimicrob Res.* 2019;17:79–83.
107. Sun S, Noorian P, McDougald D. Dual role of mechanisms involved in resistance to predation by protozoa and virulence to humans. *Front Microbiol.* 2018;9:1017.
108. Schlimme W, Marchiani M, Hanselmann K, Bernard J. Gene transfer between bacteria within digestive vacuoles of protozoa. *FEMS Microbiol Ecol.* 1997;23:239–47.
109. Rahman MH, Mahbub KR, Espinoza-Vergara G, Ritchie A, Hoque MM, Noorian P, et al. Protozoal food vacuoles enhance transformation in *Vibrio cholerae* through SOS-regulated DNA integration. *ISME J.* 2022;16:1993–2001.



110. Shu HY, Fung CP, Liu YM, Wu KM, Chen YT, Li LH, et al. Genetic diversity of capsular polysaccharide biosynthesis in clinical isolates. *Microbiology*. 2009;155:4170–83.
111. Espinoza-Vergara G, Noorian P, Silva-Valenzuela CA, Raymond BBA, Allen C, Hoque MM, et al. *Vibrio cholerae* residing in food vacuoles expelled by protozoa are more infectious in vivo. *Nat Microbiol*. 2019;4:2466–74.

### AUTHOR CONTRIBUTIONS

CSL, LJJ, and YGZ designed the study. CSL, LJJ, and LYY performed research. CSL, KXR, SYDZ, AI, RN, XRY, EC and YGZ wrote and edited the paper. All authors have read and approved the contents of the manuscript.

### FUNDING

This work was supported by the National Natural Science Foundation of China (42090063, 32061143015, 32100331).

### COMPETING INTERESTS

The authors declare no competing interests.

### ADDITIONAL INFORMATION

**Supplementary information** The online version contains supplementary material available at <https://doi.org/10.1038/s43705-023-00302-z>.

**Correspondence** and requests for materials should be addressed to Yong-Guan Zhu.

**Reprints and permission information** is available at <http://www.nature.com/reprints>

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023