Editorial

Continuum

The word continuum has been around for a long time. Webster defines it as a continuous whole, quantity, or series. Although it is not a commonly used word, it is something with which we are all familiar. However, it is taking on new meaning in healthcare and particularly in rehabilitation care in the United States. Now most institutions are taking a look at the continuum of care as it relates to a specific disease entity, in this case, spinal cord injury.

The continuum of care has always been present in spinal cord injury and consisted of mainly three elements: acute care occurring immediately after the patient's injury; rehabilitation which starts anywhere from 2 to 6 weeks following the injury; and post discharge. In most areas of the world, that's still the scenario. However, with the advent of healthcare reform and managed care in the United States, new venues for care are being developed. This development is largely prompted by economic pressures for cost containment. Over 20 years ago when the model systems program started in the United States, the length of stay averaged around 150 days. With the emphasis on system of care, those lengths of hospital stay shortened to 90–100 days by the late 1980s. If one looks at length of stay statistics, focusing only on the last 12 to 18 months, the average length of in hospital stay is around 60 days.

So what has changed in the continuum? One new venue that has evolved is the subacute hospital stay. A patient participates in a subacute program when he or she is either too sick, too weak, or restricted because of bony management constraints to being able to receive more than 3 h of therapy a day. The care necessary is either less intense or can be provided by a lesser level of skilled paraprofessional. This creates a reduction in the cost compared to a full scale rehabilitation setting.

A second venue which has developed is the day hospital. A day hospital is a program in which the patient requires occupational and physical therapy on an intensive basis but either is capable of handling skin, bowel, and bladder and other nursing issues, or else has someone trained that can provide these in a home or apartment setting. In essence, the patient receives the same level of therapy care but does not require intensive skilled nursing care provided by a professional. Patients who qualify for this level of care are usually those that previously were kept in the hospital during the last 2–4 weeks to complete rehabilitation.

A third venue which has always been present but which is now receiving intensified interest is outpatient

therapy. A patient is appropriate for this level of care when their rehabilitation has been completed except for either occupational therapy or physical therapy. This level of care can usually be completed in 2–4 weeks with the patient receiving treatment either daily or, usually, three times per week.

It is important to remember that the patient still has the same rehabilitation goals depending on the level of injury and associated secondary problems. The goals are achieved in a variety of settings now rather than all inpatient. This has produced a significant reduction in the cost of providing rehabilitation, although having to do it in an outpatient setting may prolong the total time. The reason that these cost reductions are significant is that in the United States approximately two thirds of daily inpatient charges is attributed to nursing functions.

It is unlikely that economic pressures now being applied in the United States to reduce health costs have squeezed the last dollar out of the process. It is entirely conceivable that certain aspects of inpatient care, day hospital care, and certain outpatient care could be provided in the home. Changes will probably continue to occur for the next 3–5 years. These changes are proving to be stressful for the staff involved. It is also probable that it is stressful for the patient. No longer is there time to grieve or begin an adjustment process prior to being asked to participate in a very intense inpatient rehabilitation program and before being shunted from the protective environment of hospital situation. It remains to be determined when the best time frame is for the patient to be involved in a subacute setting. It is possible that in the years ahead, we may see patients going from the acute setting to the subacute setting, or even to home after a brief period of training before being admitted to a full scale inpatient rehabilitation program. At that juncture, a patient may be better able to appreciate the value of the inpatient rehabilitation process. With better understanding and motivation the patient will be able to make better use of the most intensive part of the process, the inpatient stay.

The continuum of care for spinal cord injury will be monitored and adjusted as it bends and is molded by economic pressures. It is in the best interest of the patient and the care providers that a focus on quality be maintained, no matter in which of the many venues of care the patient is located.

Dr David Apple MD
Associate Editor