EDITOR'S PAGE

Does regular standing improve bowel function in people with spinal cord injury?

JJ Wyndaele

Antwerp University Hospital, Antwerp, Belgium

E-mail: wyndaelejj@skynet.be

Dear *Spinal Cord* reader,

We wish you and yours a very happy 2015.

SC enters its 53th year and we are proud that it continues to be a fine journal, up to date, well recognized and valuable thanks to you all.

We are planning some further changes in the Journal which will be communicated in due course.

The January Issue 2015 contains some very good contributions:

Reviews: Ghizal Fatima *et al.* elucidate the importance of oxidative stress and antioxidants and its possible relationship with SCI. Zakrasek *et al.* explored the prevalence or incidence, risk factors, and costs of pressure ulcers among individuals with SCI, specifically in the context of the developing world, to highlight important targets for intervention and research. Allison and Ditor provide an overview of the many factors which contribute to the chronic inflammatory state typically observed following SCI.

Animal studies: Shouping *et al.* investigated whether Bosentan, an endothelin-A/-B dual receptor antagonist, could protect neurons after spinal cord ischemia reperfusion injury in rats and its underlying signaling pathway.

Classification systems: Chhabra *et al.* obtained, in an online survey, the opinion of experts on whether the currently available classification systems for thoracolumbar and subaxial cervical spine injuries meet their expectations with regard to the desired objectives of a good classification system and practical implementability.

Imaging: Öztürk *et al.* found, with ultrasonography, sciatic nerves to be smaller in subjects with SCI. Together with electrophysiological data, this preliminary finding could possibly be attributed to primary axonal loss.

Autonomic/cardiovascular function: Goh *et al.* detected a high incidence of reversed dipping and nocturnal hypertension in individuals with SCI and clinically significant blood pressure disorders. They postulate that elevated nocturnal blood pressure may contribute to nocturnal diuresis which might cause relative volume depletion and thereby contribute to daytime orthostatic hypotension. Malmqvist *et al.* investigated changes in heart rate variability patterns- and alterations after acute traumatic SCI. They describe different interesting findings possibly linked to pathophysiological mechanisms in different types of SC lesion. Serra-Anó *et al.* studied the differences in heart rate variability while sitting between able-bodied participants and paraplegic individuals and found alterations in the SCI group.

Urology: Shigemura *et al.* evaluated measures for preventing multidrug resistant *Pseudomonas aeruginosa* in catheter-associated urinary tract infection (UTI) in SCI individuals. Krassioukov *et al.* demonstrated that catheter re-use is intimately linked to UTI frequency and provides novel insight on bladder function and management in elite athletes with SCI.

Bowel: Morris *et al.* describe how SCI patients have the same risk of malignancy as the general population and are less likely to undergo screening colonoscopy. Moreover colonoscopy is often limited by poor bowel preparation and lower completion rates with a subsequent lower adenoma detection rate. Harvey *et al.* found that a 6-week standing programme does not reduce Time to First Stool after SCI.

Obesity: Wong *et al.* evaluated, in an international survey in medical staff from SCI centers, the knowledge, attitudes and practices towards obesity prevention and management and present the number of beds and dietitians available at each center. Claydon *et al.* found no need to change scaling powers to improve the relationships between the measures of adiposity and indices of obesity or cardiovascular disease risk in individuals with SCI. This manuscript is discussed in an editorial note by Nash.

AGAIN MANY THANKS TO AUTHORS, TO REVIEWERS AND TO THE EDITORIAL AND PUBLISHING STAFF. TOGETHER WE MAKE SC JOURNAL WHAT IT IS.