ERRATUM

Predictors of outcome for urethral injection of botulinum toxin to treat detrusor sphincter dyssynergia in men with spinal cord injury

JM Soler, JG Previnaire and N Hadiji

Spinal Cord (2016) **54**, 492; doi:10.1038/sc.2016.1

Correction to: *Spinal Cord* (2016) **54**, 452–456; doi:10.1038/sc.2015.224; published online 29 December 2015

Since the publication of this article, the authors have noticed an error in the data in Table 4. The correct version of Table 4 is

below. The PDF and online versions of the article have been amended.

The publisher apologizes for any inconvenience caused.

Table 4 Results of the main urodynamic data

	<i>P</i> Detrusor (cm H ₂ O) n = 36	DLPP (cm H ₂ O) n = 26	<i>P</i> Urethral (cm H ₂ O) n = 36	Bladder capacity (ml) n = 36
Before injection	77 (56) [0–240]	68 (27) [0–110]	135 (45) [60–200]	272 (170) [50–650]
Follow-Up	70 (38) [0–160]	64 (22) [0–100]	130 (30) [70–200]	277 (139) [35–600]
<i>P</i>	= 0.52	= 0.83	= 0.52	= 0.91

Abbreviations: DLPP, detrusor leak point pressure; *P* detrusor, maximum detrusor pressure; *P* urethral, maximum urethral pressure. Values are means, s.d. () and extremes [–].