SCIENTIFIC DATA

OPEN

Data Descriptor: An emissions-socioeconomic inventory of Chinese cities

Yuli Shan¹, Jianghua Liu², Zhu Liu^{1,3}, Shuai Shao² & Dabo Guan^{1,3,4}

Received: 27 September 2018

Accepted: 18 January 2019

Published: 26 February 2019

As the centre of human activity and being under the threat of climate change, cities are considered to be major components in the implementation of climate change mitigation and CO₂ emission reduction strategies. Inventories of cities' emissions serve as the foundation for the analysis of emissions characteristics and policymaking. China is the world's top energy consumer and CO₂ emitter, and it is facing great potential harm from climate change. Consequently, China is taking increasing responsibility in the fight against global climate change. Many energy/emissions control policies have been implemented in China, most of which are designed at the national level. However, cities are at different stages of industrialization and have distinct development pathways; they need specific control policies designed based on their current emissions characteristics. This study is the first to construct emissions inventories for 182 Chinese cities. The inventories are constructed using 17 fossil fuels and 47 socioeconomic sectors. These city-level emissions inventories have a scope and format consistent with China's national/provincial inventories. Some socioeconomic data of the cities, such as GDP, population, industrial structures, are included in the datasets as well. The dataset provides transparent, accurate, complete, comparable, and verifiable data support for further city-level emissions studies and low-carbon/sustainable development policy design. The dataset also offers insights for other countries by providing an emissions accounting method with limited data.

Design Type(s)	modeling and simulation objective • source-based data analysis objective • time series design • data integration objective
Measurement Type(s)	carbon dioxide emission
Technology Type(s)	computational modeling technique
Factor Type(s)	geographic location
Sample Characteristic(s)	China • manufacturing process • city

¹Water Security Research Centre and School of International Development, University of East Anglia, Norwich NR4 7TJ, UK. ²School of Urban and Regional Science, Shanghai University of Finance and Economics, Shanghai 200433, China. ³Department of Earth System Science, Tsinghua University, Beijing 100080, China. ⁴Center for Energy and Environmental Policy Research, Beijing Institute of Technology, Beijing 100081, China. Correspondence and requests for materials should be addressed to S.S. (email: shao.shuai@shufe.edu.cn) or to D.G. (email: dabo.guan@uea.ac.uk)

Background & Summary

Cities are considered to be major components in the implementation of climate change mitigation and CO₂ emission reduction strategies¹. Although a mention of “city” is lacking in the Paris Agreement or the Sustainable Development Goals, as all submissions focused on the national level, climate change actions should be taken at the city level².

Cities are the basic units for human activity³ and the main consumers of energy and emitters of CO₂ throughout the world^{4,5}. The CO₂ emissions from energy use in cities will grow by 1.8% per year between 2006 and 2030, with the share of global CO₂ emissions rising from 71 to 76%⁶. In China, urban energy use accounts for 85% of total emissions, which is higher than its share in the USA (80%) or Europe (69%)^{7,8}. The high energy demand and high CO₂ emissions of cities not only increase climate change concerns and environmental pressure but also increase residents' health problems through air pollution⁹. Both coastal and interior cities are facing danger from extreme weather, geological hazards, urban waterlogging, etc. Thus, cities are motivated to fight against climate change.

Although climate policies are usually designed at the national level, they are implemented at the city level. Without support from local city governments, national policies cannot be effectively executed. Considering that cities have different natural resource endowments and development pathways, each should have specific emission reduction actions that are designed based on that city's unique emission characteristics. In China, this is particularly true. There are over 330 cities in China, and they are at different stages of industrialization, with distinct development pathways. Therefore, cities are the key components in climate change policymaking, and many low-carbon projects and actions have been taken at the city level, such as the Local Governments for Sustainability (ICLEI) and the C40 Cities Climate Leadership Group (C40).

Understanding the emissions characteristics of cities is the foundation of any further city-level climate change actions. Compared to studies focused on national and provincial emissions accounts, far fewer have focused on city-level emissions, and those that do have methods limitations and geographical restrictions.

First, previous studies on city-level emissions have severe methodological weaknesses and limitations. Most previous city-level greenhouse gas emissions inventories were calculated using a bottom-up approach, i.e., by using energy consumption data from surveys of several sectors^{10–12}. The sectors were set differently between studies, making the cities' CO₂ emissions inconsistent and not comparable across studies, as well as inconsistent with the national and regional emission inventories. In addition, some studies used spatial and geographical analysis^{13,14}, night-time light imagery^{15,16}, or economic models^{17,18} to account for city emissions. These models can only estimate the overall CO₂ emissions of a city. They cannot exactly determine the contributions of emission sources (i.e., energy types or socioeconomic sectors).

Second, most of the previous studies on city-level emissions focused on megacities from developed countries with consistent and transparent data sources, especially US cities^{19–23}. Currently, city-level emissions are being studied from a more international perspective by analysing more global cities, especially cities from developing countries^{24–30}. Restricted by data availability, the CO₂ emissions from Chinese cities are far behind in their documentation. Sugar, *et al.*³¹ reported emissions for Beijing, Tianjin, and Shanghai in 2006 and compared the three cities' emissions with those of ten other global cities. Wang, *et al.*¹⁰ discussed the CO₂ emissions from 12 Chinese megacities, most of which are provincial capital cities. Dhakal⁸ examined the energy consumption and CO₂ emissions of all Chinese provincial cities. Zhou, *et al.*³² and Xu, *et al.*³³ account for the CO₂ emissions of specific city clusters, such as the Guangdong Bay cities and cities in the central plain. Ramaswami, *et al.*³⁴ in the cited study and a follow-up study developed a comprehensive emission database including the scope 1 and scope 2 CO₂ emissions of 233 prefecture-level and 637 county-level cities in China³⁵.

Thus, the previous assessments of city-level emissions either focused on total emissions (or combined emissions for several sectors) or on megacities with consistent and systematic energy statistics. Previous analyses of the bottom-up sector-based emissions of cities are inconsistent with national and regional emission inventories, making multi-scale emission studies unavailable. Additionally, such general emission data cannot support detailed city-level emission analysis and related emission reduction policy making.

The dataset in this study provides detailed emissions inventories for 182 Chinese cities. The inventories are constructed for 17 types of fossil fuel and 47 socioeconomic sectors that are consistent with the System of National Accounts. Additional socioeconomic indexes for the cities are included in the dataset. The dataset has been re-used in our latest study¹ and will facilitate further city-level emissions studies and low-carbon/sustainable development policy design.

Methods

City boundaries and emission scopes

This dataset provides the emissions and socioeconomic inventories of 182 Chinese cities; these cities cover 82% (33,880 billion yuan) of the country's GDP (41,303 billion yuan), 64% (860 million) of the population (1,341 million), and 35% (3.4 million km²) of the land area (9.6 million km²) in 2010³⁶. Most of the studied cities are located east of the Heihe-Tengchong line, where 96% of China's population lives on 43% of the land. The 182 cities are selected based on data availability.

The term ‘city’ here refers to administrative prefecture-level city rather than to a built-up city. Accordingly, the CO₂ emissions calculated in this dataset are Intergovernmental Panel on Climate Change (IPCC) administrative territorial CO₂ emissions, referring to emissions “taking place within national (including administered) territories and offshore areas over which the country has jurisdiction (page overview.5)”³⁷. We exclude the emissions induced by international aviation and shipping³⁸. Unlike production- or consumption-based emissions¹⁷, the administrative territorial scope quantifies the direct emissions induced by human activities within a regional boundary. That is, territorial emissions provide the data baseline for emission-related studies and regional carbon control.

The emission inventories include two components: CO₂ emitted from fossil fuel combustion (energy-related emissions) and CO₂ emitted from industrial production (process-related emissions). Process-related emissions refers to CO₂ emitted from industrial raw materials during chemical reactions, such as CO₂ escaping during calcium carbonate (CaCO₃) calcination in cement production.

The cities’ emissions inventories are uniform with China’s national and provincial emission inventories in scope, format, and data sources³⁹, making them comparable.

Emissions calculation and inventory construction

The energy-related emissions are calculated based on 17 fuels (shown in Table 1) and 47 socioeconomic sectors (shown in Table 2). The 17 types of fossil fuels are selected based on China’s official energy statistical system³⁶. There are 29 energy types used in the system: 26 are fossil fuels, one is electricity, one is heat, and one is other energy. As our study only accounts for the direct emissions from fossil fuel burning within one city boundary (the IPCC administrative territorial scope), the inventories exclude the indirect emissions induced by electricity and heat use. The CO₂ emissions related to electricity and heat generation, therefore, are calculated based on fuel inputs and allocated to the power plants. We also assume that there is no, or little, CO₂ emitted from other energy uses. Some of the fossil fuels share similar carbon content and have very low consumption volumes; we merge them in the emission accounts³⁹. The 47 socioeconomic sectors are set according to the System of National Accounts⁴⁰.

Energy-related CO₂ emissions are calculated based on the mass balance theory,⁴¹ see Equation 1.

$$CE_{ij} = AD_{ij} \times NCV_i \times CC_i \times O_{ij} \quad (1)$$

where CE_{ij} represents the CO₂ emissions induced by the combustion of fuel i in sector j , AD_{ij} (activity data) represents fossil fuel combustion by fuel and sector. The emission factor ($\text{ton CO}_2/\text{ton}$) is composed of a specific heat value factor- NCV_i (J/ton) multiplied by the carbon content per unit heat value- CC_i ($\text{ton CO}_2/\text{J}$) and oxygenation efficiency- O_{ij} (quantified as percentage). Specifically, NCV_i refers to the heat value produced per physical unit of fossil fuel i combusted, CC_i is the carbon content emitted per unit heat value when combusting per physical unit of fossil fuel i , while O_{ij} stands for the oxidation ratio of the fossil fuel combusted.

The emission factors (NCV_i , CC_i , and O_{ij}) have been published by international institutions, including the IPCC and the United Nations (UN; governmental agencies in China such as the National Bureau of Statistics of China (NBS) and the National Development and Reform Commission of China (NDRC);⁴² and previous studies such as the Multi-resolution Emission Inventory for China (MEIC)⁴³, Liu, *et al.*⁴⁴. Liu, *et al.*⁴⁴ re-evaluated the carbon content of raw coal samples from 4,243 state-owned Chinese coal mines and found that the emission factors for Chinese coal are, on average, 40% lower than the default values recommended by the IPCC. After comparing Liu, *et al.*⁴⁴ emissions factors with eight different sources, our previous study finds that Liu, *et al.*⁴⁴ emission factors are relatively lower than others (shown in Table 3 (available online only)). The seven sets of emission factors are collected from IPCC, NBS, NDRC, NC1994, NC2005, MEIC, UN-China, and UN-average. Generally, coal-related fuels have a larger range than oil- and gas-related fuels. Liu, *et al.*⁴⁴’s re-evaluated emission factors have already been widely used by many studies and institutions to calculate China’s emission inventory, including China’s third official emission inventory 2012⁴⁵. Thus, this study uses the above-mentioned updated emission factors. Table 1 gives the net calorific value (NCV_i) and carbon content (CC_i). Table 4 (available online only) shows the sector-specific oxygenation efficiency (O_{ij}), which considers sector discrepancies in technical level³⁹.

The process-related CO₂ emissions (CE_t) are calculated in Equation 2⁴¹. We include seven industrial processes, including cement production (for approximately 70% of the total process-related emissions in China^{45,46}), lime production (the 2nd largest emissions source⁴⁷), ammonia production, soda ash production, ferrochromium production, silicon metal production, and unclassified ferro-production. The process-related emissions are allocated to the corresponding sectors in the emission inventory. Cement and lime-related emissions are allocated to the sector “Non-metal Mineral Products”; ammonia and soda ash-related emissions are allocated to the sector “Raw Chemical Materials and Chemical Products”; Ferrochromium, silicon metal, and unclassified ferro-related emissions are allocated to the sector “Smelting and Pressing of Ferrous Metals”.

$$CE_t = AD_t \times EF_t \quad (2)$$

AD_t and EF_t in the equation refer to industrial production (activity data) and emission factors, respectively. The emission factors of industrial processes are collected from IPCC⁴¹ and NDRC⁴², as shown in Table 5.

No. (i) Unit	Fuels in China's Energy Statistics	Fuels in this study	NCV _i	CC _i
			PJ/10 ⁴ tonnes, 10 ⁸ m ³	tonne C/TJ
1	Raw coal	Raw coal	0.21	26.32
2	Cleaned coal	Cleaned coal	0.26	26.32
3	Other washed coal	Other washed coal	0.15	26.32
4	Briquettes	Briquette	0.18	26.32
		Gangue		
5	Coke	Coke	0.28	31.38
6	Coke oven gas	Coke over gas	1.61	21.49
7	Blast furnace gas	Other gas	0.83	21.49
	Converter gas			
	Other gas			
8	Other coking products	Other coking products	0.28	27.45
9	Crude Oil	Crude oil	0.43	20.08
10	Gasoline	Gasoline	0.44	18.9
11	Kerosene	Kerosene	0.44	19.6
12	Diesel oil	Diesel oil	0.43	20.2
13	Fuel oil	Fuel oil	0.43	21.1
14	Naphtha	Other petroleum products	0.51	17.2
	Lubricants			
	Paraffin			
	White spirit			
	Bitumen asphalt			
	Petroleum coke			
	Other petroleum products			
15	Liquefied petroleum gas (LPG)	LPG	0.47	20
16	Refinery gas	Refinery gas	0.43	20.2
17	Nature gas	Nature gas	3.89	15.32

Table 1. Fossil fuels in the city-level emissions inventories and emissions factors.

The cities' CO₂ emissions matrices (namely, inventories) are created as 19 columns and 48 rows. Seventeen fossil fuel-related emissions, process-related emissions and total emissions are represented by 19 columns, while 47 rows correspond to the 47 socioeconomic sectors. Each element of the matrices is identified as the CO₂ emissions from fossil fuel combustion/industrial production in the corresponding sector. An inventory of Beijing is given in Table 6 (available online only) as an example.

These methods on emission inventory construction are expanded version of descriptions in our related work³⁹. MATLAB R2014a is used to construct the cities' emission inventories. We provided the MATLAB code in the Supplementary Information. We also provided the activity data of the cities for additional data transparency and verifiability (see "China city-level Energy inventory, 2010", Data Citation 1). Researchers will be able to use the MATLAB code and energy inventories to recalculate the emission inventories for the cities or replicate to other cities.

Activity data collection

Fossil fuel combustion, i.e., the activity data for energy-related emission accounts, includes two parts: the energy inputs for electricity/heat generation and the total final consumption. Other inputs for energy transformation, such as coal cleaning or petroleum refineries, transfer the carbon element from one fuel to another. These processes emit little CO₂. Following our previous emissions inventories constructed for China and its provinces³⁹, fossil fuel combustion can be collected from a region's energy balance table (EBT) and final energy consumption can be captured by the industrial sector ($Energy_{ij}$). The EBT provides each fossil fuel's transformation and final consumption in farming, industry, construction, three service sectors, and households (rural and urban). As the entire industry sector consists of 40 sub-sectors, $Energy_{ij}$ presents the sectoral consumption of fossil fuel for the industry sector.

Generally, the EBT and $Energy_{ij}$ can be found in a city's statistical yearbook. However, due to the poor data quality of city-level statistics, not all cities' yearbooks publish the EBT or $Energy_{ij}$. We developed a series of methods in our previous study to estimate missing data⁴⁸:

No. (j)	Socioeconomic sectors	Category		
1	Farming, Forestry, Animal Husbandry, Fishery and Water Conservancy	The primary industry		
2	Coal Mining and Dressing	Energy production	Manufacturing industries	
3	Petroleum and Natural Gas Extraction	Energy production		
4	Ferrous Metals Mining and Dressing	Energy production		
5	Nonferrous Metals Mining and Dressing	Energy production		
6	Non-metal Minerals Mining and Dressing	Energy production		
7	Other Minerals Mining and Dressing	Energy production		
8	Logging and Transport of Wood and Bamboo	Light manufacturing		
9	Food Processing	Light manufacturing		
10	Food Production	Light manufacturing		
11	Beverage Production	Light manufacturing		
12	Tobacco Processing	Light manufacturing		
13	Textile Industry	Light manufacturing		
14	Garments and Other Fibre Products	Light manufacturing		
15	Leather, Furs, Down and Related Products	Light manufacturing		
16	Timber Processing, Bamboo, Cane, Palm Fibre & Straw Products	Light manufacturing		
17	Furniture Manufacturing	Light manufacturing		
18	Papermaking and Paper Products	Light manufacturing		
19	Printing and Record Medium Reproduction	Light manufacturing		
20	Cultural, Educational and Sports Articles	Light manufacturing		
21	Petroleum Processing and Coking	Energy production		
22	Raw Chemical Materials and Chemical Products	Heavy manufacturing		
23	Medical and Pharmaceutical Products	Light manufacturing		
24	Chemical Fibre	Heavy manufacturing		
25	Rubber Products	Heavy manufacturing		
26	Plastic Products	Heavy manufacturing		
27	Non-metal Mineral Products	Heavy manufacturing		
28	Smelting and Pressing of Ferrous Metals	Heavy manufacturing		
29	Smelting and Pressing of Nonferrous Metals	Heavy manufacturing		
30	Metal Products	Heavy manufacturing		
31	Ordinary Machinery	Heavy manufacturing		
32	Equipment for Special Purposes	Heavy manufacturing		
33	Transportation Equipment manufacturing	Heavy manufacturing		
34	Electric Equipment and Machinery	High-tech industry		
35	Electronic and Telecommunications Equipment	High-tech industry		
36	Instruments, Meters, Cultural and Office Machinery	High-tech industry		
37	Other Manufacturing Industry	High-tech industry		
38	Scrap and waste	High-tech industry		
39	Production and Supply of Electric Power, Steam and Hot Water	Energy production		
40	Production and Supply of Gas	Energy production		
41	Production and Supply of Tap Water	Heavy manufacturing		
42	Construction	Construction		
43	Transportation, Storage, Post and Telecommunication Services	Services sectors		
44	Wholesale, Retail Trade and Catering Services			
45	Other Service Sectors			
46	Urban Resident Energy Usage	Household		
47	Rural Resident Energy Usage			

Table 2. Sectors' definition of the emission inventories.

No.	Industry process	Emission factors	Allocation sectors
1	Ammonia production	1.5000	Raw Chemical Materials and Chemical Products
2	Soda Ash production	0.4150	Raw Chemical Materials and Chemical Products
3	Cement production	0.4985	Non-metal Mineral Products
4	Lime production	0.6830	Non-metal Mineral Products
5	Ferrochromium production	1.3000	Smelting and Pressing of Ferrous Metals
6	Silicon metal production	4.3000	Smelting and Pressing of Ferrous Metals
7	Ferro-unclassified production	4.0000	Smelting and Pressing of Ferrous Metals

Table 5. 7 industrial processes and emissions factors.

1. EBT: Very few cities have EBT in their statistical yearbooks. We scale down the corresponding provincial EBT to obtain the city table. We use each sector’s GDP to estimate farming, construction, and three service sectors, assuming that the city has the same farming/construction/service energy intensity as its province. We also use the urban/rural population to estimate the urban/rural household energy estimation on the premise that the city has the same per capita residential energy consumption as its province. The GDP and population data are collected from statistical yearbooks for the cities and their corresponding provinces.
2. $Energy_{ij}$: Some cities only provide $Energy_{ij}$ from enterprises of above-designated-size (ADS). ADS enterprises are defined as enterprises with prime operating revenue over 20 or 5 million yuan for different cities. ADS enterprises account for 50 to 90% (roughly) of one city’s total industrial output. We use the ADS industrial output ratio (calculated as the whole-industry output divided by the ADS enterprises’ output) to scale up ADS $Energy_{ij}$ and obtain sectoral fossil fuel consumption at the whole-industry scale.

As for cement production, the cities’ statistical yearbooks provide total cement production or production from ADS enterprises. We then scaled up the ADS cement production by the ADS industrial output ratio to obtain the total cement production.

The raw activity data are collected through a “crowd-sourcing” working mode implemented in the Applied Energy Summer School 2017 and 2018. Over 100 students joined the summer school and participated in data collection. The summer school will be held annually in the future, and more researchers will contribute to and update city-level data collection.

These methods on city-level data estimation and collection are expanded version of descriptions in our related work⁴⁸.

Socioeconomic indexes

This study collects several socioeconomic indexes for the 182 cities from the “China City Statistical Yearbook”⁴⁹, including:

1. population, in 10 thousand;
2. employed population, in 10 thousand;
3. employed population in sectors (primary industry; mining; manufacturing, electric power, gas and water production and supply; construction; transport, storage and post; information transmission, computer services and software industry; wholesale and retail trade; hotel and catering services; financial intermediation; real estate; leasing and business services; scientific research, technical services and geological exploration; water, environmental and public facilities management; resident services and other services; education; health, social security and social welfare; culture, sports and entertainment; public administration and social organization), in 10 thousand;
4. area, in square kilometres;
5. built up area, in square kilometres;
6. gross domestic product (GDP), in 10 thousand yuan;
7. primary industry, secondary industry, and tertiary industry’s share in GDP, in %;
8. industrial output, in 10 thousand yuan.

The socioeconomic indexes (as shown in Table 7 (available online only) and “China city-level socioeconomic inventory, 2010”, Data Citation 1) can be used to explore the drivers and characteristics of cities’ emissions.

Data Records

A total of 365 data records (emissions-socioeconomic inventories) are contained in the datasets. Of these,

- 182 are emissions inventories for cities (2010) [“China city-level emissions inventory, 2010”, Data Citation 1];
- 182 are energy inventories for cities (2010) [“China city-level energy inventory, 2010”, Data Citation 1];
- 1 is a socioeconomic inventory for cities (2010) [“China city-level socioeconomic inventory, 2010”, Data Citation 1];

The cities’ CO₂ emissions inventories are constructed at an IPCC territorial administrative scope, including both energy-related emissions (from fossil fuel combustion) and process-related emissions (from cement production). The socioeconomic inventory presents GDP, population, employed population (with structure), GDP (with structure), and area of the 182 cities.

Technical Validation

Uncertainties

CO₂ emissions inventories gather the contributions of economic activity to total CO₂ emissions for a given time period and area. Inventories are critical to many environmental decision-making processes and scientific goals. Policymaking and scientific research require reliable inventories to ensure the effectiveness of the policy process. In both types of applications, it is important to understand the uncertainty in emissions inventories. Additionally, uncertainty analysis can improve the accuracy of emissions accounts. Regarding the city-level CO₂ emissions inventories in this article, the literature shows that uncertainty regarding the process-related emissions in cement production is low. The inventories’ uncertainty mainly depends on energy-related emissions part^{44,50}. The contributing sources of uncertainty for energy-related emissions accounting are associated with emission factors, activity data and other estimation parameters (*Volume 1, Chapter 3, Page 6*)⁴¹. The uncertainty induced by emissions factors and energy activity data are both quantified for the cities’ emission inventories.

Uncertainties in activity data and emission factors. China’s energy data are of relatively poor quality compared with those of developed countries, especially city-level data. The literature also shows that the uncertainties range widely from sector to sector. The coefficient of variation (CV; the standard deviation divided by the mean) is used to quantify the uncertainty. According to a field survey led by previous studies, the fossil fuel consumed in China’s power generation sector has the lowest CV (5%)^{51,52}, compared with primary industry (30%)⁵³, other manufacturing sectors (10%), construction (10%)^{41,54}, transportation sector (16%)⁵⁵, and residential energy use (20%)⁴¹. The sources of uncertainties could lie in the opaqueness in China’s statistical systems, especially on the “*statistical approach on data collection, reporting and validation (Page 673)*”⁵⁶ and the dependence of China’s statistics departments on other government departments. Such uncertainties result in a large gap between China’s national fossil fuel consumption data and the aggregated provincial data. To cover the gap, China has adjusted its energy data three times since 2004, resulting in a gap between the latest national fossil fuel consumption data and provincial aggregated data of 5%⁵⁷. The gap between city-level aggregated energy consumption and the national overall data could be even larger.

Previous studies have debated China’s emission factors^{58–61}. The range of emission factors across different sources is as high as 40%. This study collects emission factors from Liu, *et al.*⁴⁴, which measured them based on a broad investigation of China’s fuel quality. Based on the statistical analysis of surveyed fuel quality, the CVs of coal-, oil-, and gas-related fuels are estimated as 3, 1, and 2%, respectively.

Monte Carlo simulations. Monte Carlo methods are used to simulate the uncertainties resulting from both fossil fuel combustion and emissions factors to estimate the overall uncertainty of the emissions⁴¹. Monte Carlo simulations select random values for the emission factor and activity data (fossil fuel consumption) from within their individual normal probability (density) functions and calculate the corresponding emission values (chapter 6 IPCC⁴¹). To perform Monte Carlo simulations, we first set up probability density functions for each input variable (emission factor and activity data). Both variables are assumed to follow a normal distribution⁴⁴. Then, we randomly sample both the activity data and the emission factors 20,000 times and obtain 20,000 CO₂ emission estimations. The uncertainties are obtained at a 97.5% confidence level and are calculated as the 97.5% confidence intervals of the estimates.

This article finds that the average uncertainties in the cities’ total CO₂ emissions range from −3.65 to 3.67% at a 97.5% confidence level (±47.5% confidence interval around the estimate). Hegang in Heilongjiang has the highest uncertainties in emissions of (−5.83, 5.86%), while Huizhou in Guangxi has the lowest value of (−0.91, 0.91%).

Limitations and future work

The cities’ emission inventories have some limitations that could lead to more uncertainty. Although these uncertainties may not be large enough to quantify, they are an indispensable component of the emission inventories’ uncertainties. First, this study only takes the energy-related and process-related emissions from seven industrial production processes into account in the emission accounts, and emissions emitted by other sources is missing, such as “agriculture”, “land-use change and forestry”, “waste”, and other industrial processes. Thus, the analysis is incomplete. In the future, we will expand the emission scope to achieve more complete inventories for cities. Second, the cities’ emission factors for

fossil fuels and industrial processes are substituted by national average emission factors during the process of accounting for cities' CO₂ emissions, resulting in inaccuracy. We hope that specific city-level emissions factors could be updated in the future to increase the accuracy of our results. If not, in our future research, we could employ provincial emission factors to obtain a more accurate emission inventory for the provinces. Third, due to the poor data quality for the cities, the EBTs of most cities are a downscaled version of the provincial table, assuming that the cities have the same sectoral energy intensity and per capita residential energy consumption with their provinces. Such assumptions bring additional uncertainties to cities' emission inventories. In the future, a consistent time-series emission inventory dataset for Chinese cities will be completed. We will integrate the bottom-up estimations (calculated based on survey data from enterprises)¹⁴ and satellite observations to achieve more emission accounts for these cities. More specifically, the high-resolution bottom-up emissions and satellite images can confirm some of the cities' emission sources (i.e. some super-emitting points). The night-light data will also be used to verify our top-down emissions inventories^{16,62}.

References

1. Shan, Y. *et al.* City-level climate change mitigation in China. *Science Advances* **4**, eaaq0390 (2018).
2. Tong, K. *et al.* The collective contribution of Chinese cities to territorial and electricity-related CO₂ emissions. *Journal of Cleaner Production* **189**, 910–921 (2018).
3. Van den Hoven, J., Lokhorst, G.-J. & Van de Poel, I. Engineering and the problem of moral overload. *Science and engineering ethics* **18**, 143–155 (2012).
4. Wang, Y., Song, Q., He, J. & Qi, Y. Developing low-carbon cities through pilots. *Climate Policy* **15**, S81–S103 (2015).
5. Satterthwaite, D. Cities' contribution to global warming: notes on the allocation of greenhouse gas emissions. *Environment and urbanization* **20**, 539–549 (2008).
6. IEA. *Cities, towns and renewable energy*. (International Energy Agency, 2009).
7. Dhakal, S. GHG emissions from urbanization and opportunities for urban carbon mitigation. *Current Opinion in Environmental Sustainability* **2**, 277–283, <http://doi.org/10.1016/j.cosust.2010.05.007> (2010).
8. Dhakal, S. Urban energy use and carbon emissions from cities in China and policy implications. *Energy Policy* **37**, 4208–4219, <http://doi.org/10.1016/j.enpol.2009.05.020> (2009).
9. Kan, H., Chen, B. & Hong, C. Health impact of outdoor air pollution in China: current knowledge and future research needs. *Environmental Health Perspectives* **117**, A187 (2009).
10. Wang, H., Zhang, R., Liu, M. & Bi, J. The carbon emissions of Chinese cities. *Atmospheric Chemistry and Physics* **12**, 6197–6206 (2012).
11. Glaeser, E. L. & Kahn, M. E. The greenness of cities: carbon dioxide emissions and urban development. *Journal of urban economics* **67**, 404–418 (2010).
12. Kennedy, C. *et al.* Methodology for inventorying greenhouse gas emissions from global cities. *Energy Policy* **38**, 4828–4837, <http://doi.org/10.1016/j.enpol.2009.08.050> (2010).
13. Cai, B. Characteristics analysis of CO₂ emissions of cities in China: Based on 0.1 degree grid dataset. *China Population, Resources and Environment* **22**, 151–157 (2012).
14. Cai, B. *et al.* China high resolution emission database (CHRED) with point emission sources, gridded emission data, and supplementary socioeconomic data. *Resources, Conservation and Recycling* **129**, 232–239 (2018).
15. Doll, C. N. H., Muller, J.-P. & Elvidge, C. D. Night-time imagery as a tool for global mapping of socioeconomic parameters and greenhouse gas emissions. *AMBIO: a Journal of the Human Environment* **29**, 157–162 (2000).
16. Meng, L., Graus, W., Worrell, E. & Huang, B. Estimating CO₂ (carbon dioxide) emissions at urban scales by DMSP/OLS (defense meteorological satellite program's operational linescan system) nighttime light imagery: Methodological challenges and a case study for China. *Energy* **71**, 468–478 (2014).
17. Mi, Z. *et al.* Consumption-based emission accounting for Chinese cities. *Applied Energy* **184**, 1073–1081, <http://doi.org/10.1016/j.apenergy.2016.06.094> (2016).
18. Auffhammer, M., Sun, W., Wu, J. & Zheng, S. The decomposition and dynamics of industrial carbon dioxide emissions for 287 Chinese cities in 1998–2009. *Journal of Economic Surveys* **30**, 460–481 (2016).
19. Ramaswami, A., Hillman, T., Janson, B., Reiner, M. & Thomas, G. A demand-centered, hybrid life-cycle methodology for city-scale greenhouse gas inventories. *Environmental science & technology* **42**, 6455–6461 (2008).
20. Hillman, T. & Ramaswami, A. Greenhouse gas emission footprints and energy use benchmarks for eight US cities. *Environmental science & technology* **44**, 1902–1910 (2010).
21. Parshall, L. *et al.* Modeling energy consumption and CO₂ emissions at the urban scale: methodological challenges and insights from the United States. *Energy Policy* **38**, 4765–4782 (2010).
22. Markolf, S. A., Matthews, H. S., Azevedo, I. L. & Hendrickson, C. An integrated approach for estimating greenhouse gas emissions from 100 US metropolitan areas. *Environmental Research Letters* **12**, 024003 (2017).
23. Fragkias, M., Lobo, J., Strumsky, D. & Seto, K. C. Does size matter? Scaling of CO₂ emissions and US urban areas. *PLoS One* **8**, e64727 (2013).
24. Ibrahim, N., Sugar, L., Hoornweg, D. & Kennedy, C. Greenhouse gas emissions from cities: comparison of international inventory frameworks. *Local Environment* **17**, 223–241 (2012).
25. D'Almeida Martins, R. & da Costa Ferreira, L. Climate change action at the city level: tales from two megacities in Brazil. *Management of Environmental Quality: An International Journal* **22**, 344–357 (2011).
26. Ali, G. & Nitivattananon, V. Exercising multidisciplinary approach to assess interrelationship between energy use, carbon emission and land use change in a metropolitan city of Pakistan. *Renewable and Sustainable Energy Reviews* **16**, 775–786 (2012).
27. Matese, A., Gioli, B., Vaccari, F., Zaldei, A. & Miglietta, F. Carbon dioxide emissions of the city center of Firenze, Italy: measurement, evaluation, and source partitioning. *Journal of Applied Meteorology and Climatology* **48**, 1940–1947 (2009).
28. Harvey, L. D. Tackling urban CO₂ emissions in Toronto. *Environment: Science and Policy for Sustainable Development* **35**, 16–44 (1993).
29. Gomi, K., Shimada, K. & Matsuoka, Y. A low-carbon scenario creation method for a local-scale economy and its application in Kyoto city. *Energy Policy* **38**, 4783–4796 (2010).
30. Creutzig, F., Baiocchi, G., Bierkandt, R., Pichler, P. P. & Seto, K. C. Global typology of urban energy use and potentials for an urbanization mitigation wedge. *Proc Natl Acad Sci USA* **112**, 6283–6288, <http://doi.org/10.1073/pnas.1315545112> (2015).
31. Sugar, L., Kennedy, C. & Leman, E. Greenhouse gas emissions from chinese cities. *Journal of Industrial Ecology* **16**, 552–563, <http://doi.org/10.1111/j.1530-9290.2012.00481.x> (2012).

32. Zhou, Y., Shan, Y., Liu, G. & Guan, D. Emissions and low-carbon development in Guangdong-Hong Kong-Macao Greater Bay Area cities. *Applied Energy* **228**, 1683–1692 (2018).
33. Xu, X. *et al.* Patterns of CO₂ emissions in 18 central Chinese cities from 2000 to 2014. *Journal of Cleaner Production* **172**, 529–540, <http://doi.org/10.1016/j.jclepro.2017.10.136> (2017).
34. Ramaswami, A., Jiang, D., Tong, K. & Zhao, J. Impact of the Economic Structure of Cities on Urban Scaling Factors: Implications for Urban Material and Energy Flows in China. *Journal of Industrial Ecology* **22**, 392–405 (2017).
35. Ramaswami, A. *et al.* Urban cross-sector actions for carbon mitigation with local health co-benefits in China. *Nature Climate Change* **7**, 736 (2017).
36. NBS. National Data. *National Bureau of Statistics*, <http://data.stats.gov.cn/easyquery.htm?cn=C01> (2018).
37. IPCC. *IPCC Guidelines for national greenhouse gas inventories* 1996. (Intergovernmental Panel on Climate Change, 1996).
38. Barrett, J. *et al.* Consumption-based GHG emission accounting: a UK case study. *Climate Policy* **13**, 451–470 (2013).
39. Shan, Y. *et al.* China CO₂ emission accounts 1997–2015. *Scientific Data* **5**, 170201, <http://doi.org/10.1038/sdata.2017.201> (2018).
40. National Administration for Quality Supervision and Inspection and Quarantine of China. *National industries classification (GB/T 4754-2011)*, http://www.stats.gov.cn/statsinfo/auto2073/201406/t20140606_564743.html (2011).
41. IPCC. *IPCC Guidelines for national greenhouse gas inventories* 2006. (Institute for Global Environmental Strategies (IGES), 2006).
42. NDRC. *Guidelines for provincial greenhouse gas inventories*, <http://www.cbcsd.org.cn/sjk/nengyuan/standard/home/20140113/download/shengjiwenshiqiti.pdf> (2011).
43. MEIC. National CO₂ emissions, <http://www.meicmodel.org/index.html> (2017).
44. Liu, Z. *et al.* Reduced carbon emission estimates from fossil fuel combustion and cement production in China. *Nature* **524**, 335–338 (2015).
45. NDRC. *First biennial update report on climate change*, <http://qhs.ndrc.gov.cn/dtj/201701/W020170123346264208002.pdf> (2016).
46. Shan, Y. *et al.* Peak cement-related CO₂ emissions and the changes in drivers in China. *Journal of Industrial Ecology*, in press, <http://doi.org/10.1111/jiec.12839> (2019).
47. Shan, Y., Liu, Z. & Guan, D. CO₂ emissions from China's lime industry. *Applied Energy* **166**, 245–252, <http://doi.org/10.1016/j.apenergy.2015.04.091> (2016).
48. Shan, Y. *et al.* Methodology and applications of city level CO₂ emission accounts in China. *Journal of Cleaner Production* **161**, 1215–1225, <http://doi.org/10.1016/j.jclepro.2017.06.075> (2017).
49. Bureau of Statistics. *China city statistical yearbook 2011 [Chinese Document]*. (China Statistics Press, 2011).
50. Zhao, Y., Nielsen, C. P., Lei, Y., McElroy, M. B. & Hao, J. Quantifying the uncertainties of a bottom-up emission inventory of anthropogenic atmospheric pollutants in China. *Atmospheric Chemistry and Physics* **11**, 2295–2308, <http://doi.org/10.5194/acp-11-2295-2011> (2011).
51. Zhao, Y. *et al.* Primary air pollutant emissions of coal-fired power plants in China: Current status and future prediction. *Atmospheric Environment* **42**, 8442–8452 (2008).
52. Wu, Y., Streets, D. G., Wang, S. X. & Hao, J. M. Uncertainties in estimating mercury emissions from coal-fired power plants in China. *Atmospheric Chemistry and Physics* **10**, 2937–2946 (2010).
53. Wang, S. X. & Zhang, C. Y. Spatial and temporal distribution of air pollutant emissions from open burning of crop residues in China. *Sciencepaper Online* **3**, 329–333 (2008).
54. Zhang, Q. *et al.* NO_x emission trends for China, 1995–2004: The view from the ground and the view from space. *Journal of Geophysical Research: Atmospheres* **112**, 1–18 (2007).
55. Karvosenoja, N. *et al.* Evaluation of the emissions and uncertainties of PM_{2.5} originated from vehicular traffic and domestic wood combustion in Finland. *Boreal environment research* **13**, 465–474 (2008).
56. Guan, D., Liu, Z., Geng, Y., Lindner, S. & Hubacek, K. The gigatonne gap in China's carbon dioxide inventories. *Nature Climate Change*, <http://doi.org/10.1038/nclimate1560> (2012).
57. Zheng, H. *et al.* How modifications of China's energy data affect carbon mitigation targets. *Energy policy* **116**, 337–343 (2018).
58. Olivier, J. G. J., Janssens-Maenhout, G., Muntean, M. & Peters, J. A. H. W. *Trends in global CO₂ emissions: 2015 report.* (PBL Publishers, 2015).
59. Le Quéré, C. *et al.* Global carbon budget 2015. *Earth System Science Data* **7**, 349–396 (2015).
60. Korsbakken, J. I., Peters, G. P. & Andrew, R. M. Uncertainties around reductions in China's coal use and CO₂ emissions. *Nature Climate Change* **6**, 687–690 (2016).
61. Jackson, R. B. *et al.* Reaching peak emissions. *Nature Climate Change* **6**, 7–10 (2016).
62. Shi, K. *et al.* Evaluating the ability of NPP-VIIRS nighttime light data to estimate the gross domestic product and the electric power consumption of China at multiple scales: A comparison with DMSP-OLS data. *Remote Sensing* **6**, 1705–1724 (2014).

Data Citation

1. Shan, Y., Liu, J., Liu, Z., Shao, S. & Guan, D. *Figshare* <https://doi.org/10.6084/m9.figshare.c.4237544.v1> (2018).

Acknowledgements

The authors acknowledge the efforts and “crowd-sourcing” work of the Applied Energy Summer School 2017 and 2018 held in Nanjing Normal University and Tsinghua University. This work was supported by the National Key R&D Programme of China (2016YFA0602604), the Natural Science Foundation of China (71533005, 71874097, 71503156, 91846301, 41629501, 41501605, 71873059, 71503156, 71773075, 71503168, and 71373153), the National Social Science Foundation of China (15CJY058), Chinese Academy of Engineering (2017-ZD-15-07), the UK Natural Environment Research Council (NE/N00714X/1 and NE/P019900/1), the Economic and Social Research Council (ES/L016028/1), the Royal Academy of Engineering (UK-CIAPP/425).

Author Contributions

Y.S. led the project, calculated and assembled the data, and prepared the manuscript. D.G. designed the research. J.L. collected the raw data and participated in the database construction. Z.L. and S.S. revised the manuscript.

Additional Information

Tables 3, 4, 6 and 7 are only available in the online version of this paper.

Supplementary information accompanies this paper at <http://www.nature.com/sdata>

Competing interests: The authors declare no competing interests.

How to cite this article: Rashid, H. *et al.* An emissions-socioeconomic inventory of Chinese cities. *Sci. Data*. 6:190027 <https://doi.org/10.1038/sdata.2019.27> (2019).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

The Creative Commons Public Domain Dedication waiver <http://creativecommons.org/publicdomain/zero/1.0/> applies to the metadata files made available in this article.

© The Author(s) 2019