that all medical staff personnel should receive simple instructions about management of dental trauma. Both Lin *et al.*² and Levin *et al.*³ also stated the importance of providing primary caregivers with the relevant education to improve their knowledge and ability of dealing with diagnosis and treatment of dental trauma. However, it is not only medical primary care providers who need continuing education and training in dental trauma. Hu *et al.*⁴ reported that general dental practitioners may also have limited knowledge in dental trauma management.

In conclusion, this audit revealed the majority of dento-alveolar trauma is likely to initially present outwith the dental hospital trauma clinic and highlights the need to ensure the provision of training in the management of dental trauma encompasses all centres likely to be presented with these injuries. Regular and repeat training to include rotational medical junior staff will be required and protocols for treatment and pathways of advice and referral should be clear and practical.

J. M. Phillips,
I. Shafi,
B. Kargul,
R. R. Welbury
Glasgow

1. Abu-Dawoud M, Al-Enezi B, Andersson L. Knowledge of emergency management of avulsed teeth among young physicians and dentists. *Dent Traumatol* 2007; **23**: 348-355.
2. Lin S, Levin L, Emodi O, Fuss Z, Peled M. Physician and emergency medical technicians' knowledge and experience regarding dental trauma. *Dent Traumatol* 2006; **22**: 124-126.
3. Levin L, Lin S, Emodi O, Gordon M, Peled M. Dento-alveolar and maxillofacial injuries - a survey of knowledge of the regimental aid providers in the Israeli army. *Dent Traumatol* 2007; **23**: 243-246.
4. Hu L W, Prisco C R, Bombana A C. Knowledge of Brazilian general dentists and endodontists about the emergency management of dento-alveolar trauma. *Dent Traumatol* 2006; **22**: 113-117.

DOI: 10.1038/sj.bdj.2011.292

NEEDLE RE-SHEATHING

Sir, we have recently received information from the local infection control specialist on the new EU directive banning re-sheathing of needles, which came into effect last year.

We run a large organisation where the dentists are split between using fully disposable syringes and metal syringes with disposable needles and resheath-

ing devices such as the Jenger. The directive is specifically called 'COUNCIL DIRECTIVE 2010/32/EU of 10 May 2010 implementing the Framework Agreement on prevention from sharp injuries in the hospital and health-care sector concluded by HOSPEEM and EPSU' which states, 'the practice of recapping shall be banned with immediate effect'.

I would like to pass on this information as it is as yet not to the best of my knowledge a well known directive within the profession (not surprising with the huge amount of bureaucracy we have suffered under as a profession as of late). Also perhaps the topic can be debated openly and advice for all members obtained as I know many of our own dentists will not be happy moving to fully disposable systems in favour of their favourite metal syringes.

R. Emanuel
Haywards Heath

DOI: 10.1038/sj.bdj.2011.293

ODONTOIATROS

Sir, I write further to K. Wilson's letter explaining the translation of the German word for 'dentist' (*BDJ* 2011; **210**: 53).

The same applies to Modern Greek, where the word for dentist is 'odontoiatros'. *Odonto* means of the teeth, and *iatros* is doctor. So a Greek dentist is also a tooth doctor.

D. Cooper
Halesowen

DOI: 10.1038/sj.bdj.2011.294

AN INTEGRAL LINK

Sir, at a dental conference held on 9-10 November 2009 in Göteborg, Sweden, at the initiative of the dental company Oral Care AB, a network of experts in geriatric dentistry was founded.

This interdisciplinary and autonomous network of expertise in areas such as pharmacology, pharmacokinetics, basic research, medicine and dentistry, will formulate guidelines for future geriatric dentistry. The group will work on quality assurance, quality development, as well as basic and applied research.

One of the objectives of this work is to promote dentistry as a natural and integral part of the chain of care of elderly and medically debilitated

people, in which dentistry is highly neglected today. Studies have shown that one in ten deaths from health care-related pneumonia in the elderly could be prevented by improving oral hygiene.

The number of elderly is growing rapidly in Europe and worldwide, but resources to provide good dental care are lacking to a large extent. This will inevitably lead to great suffering for this often multi-disease patient group, with malnutrition and worsened general health as the consequence. At the first meeting, the network therefore discussed how dentistry should be designed to rapidly meet this growing dental care need. It was also concluded that it is one thing to provide treatment of oral diseases, but equally important is to address the prevention of them.

At the 46th annual Swedish Dental Congress on 19 November 2010 in Göteborg, Sweden, Oral Care held an interdisciplinary symposium on elderly and geriatric dentistry, and provided informed on how IT support may be used in outreach geriatric dentistry.

About Oral Care AB

Oral Care is working with mobile dental care, where the main benefit for the patients is that both preventive measures and treatments can be given where the patient lives: at nursing homes or in private residences. Oral Care's complete preventive dentistry system also contributes to community benefits by decreasing the burden on the health care organisations by reducing the risk of morbidity. Oral Care conducts research in collaboration with leading academic institutions to develop tomorrow's geriatric dental care, with a strong focus on the links to medical benefits for the individual that adequate dental care among the elderly will lead to.

O. Johansson,
U. Fagerholm,
Å. Rundgren,
P. Sjögren
Sweden

DOI: 10.1038/sj.bdj.2011.295

REFORMING SURROUNDINGS

Sir, over the last few decades, the patient-clinician relationship has changed drastically. Today, the principle of patient